
## Le-Feng Chen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3785279/publications.pdf Version: 2024-02-01



LE-FENC CHEN

#	Article	IF	CITATIONS
1	Addition of Fibroblast-Stromal Cell Markers to Immune Synovium Pathotypes Better Predicts Radiographic Progression at 1 Year in Active Rheumatoid Arthritis. Frontiers in Immunology, 2021, 12, 778480.	2.2	3
2	Gasdermin-E Mediated Pyroptosis—A Novel Mechanism Regulating Migration, Invasion and Release of Inflammatory Cytokines in Rheumatoid Arthritis Fibroblast-like Synoviocytes. Frontiers in Cell and Developmental Biology, 2021, 9, 810635.	1.8	12
3	Neglected extra-articular manifestations in rheumatoid arthritis patients with normal body mass index: reduced skeletal muscle overlapping overfat. Therapeutic Advances in Chronic Disease, 2020, 11, 204062232097524.	1.1	3
4	Activation of the Peroxisome Proliferator–Activated Receptor γ Coactivator 1β/NFATc1 Pathway in Circulating Osteoclast Precursors Associated With Bone Destruction in Rheumatoid Arthritis. Arthritis and Rheumatology, 2019, 71, 1252-1264.	2.9	18
5	AB0579â€COMPARISON OF CLINICAL CHARACTERISTICS OF EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS BETWEEN RHEUMATOLOGY AND RESPIRATORY MEDICINE: A SINGLE CENTER, RETROSPECTIVE STUDY. , 2019, , .		0
6	THU0069â€RHEUMATOID CACHEXIA ASSESSED BY BODY COMPOSITION IS ASSOCIATED WITH WORSE DISEA IN RHEUMATOID ARTHRITIS PATIENTS WITH NORMAL BODY MASS INDEX. , 2019, , .	SE	0
7	SATO081â€BOTH OVERFAT AND MYOPENIA ARE ASSOCIATED WITH PHYSICAL DYSFUNCTION IN RHEUMATOID ARTHRITIS PATIENTS. , 2019, , .	)	0
8	Presence of hepatitis B virus in synovium and its clinical significance in rheumatoid arthritis. Arthritis Research and Therapy, 2018, 20, 130.	1.6	14
9	Short-course tocilizumab increases risk of hepatitis B virus reactivation in patients with rheumatoid arthritis: a prospective clinical observation. International Journal of Rheumatic Diseases, 2017, 20, 859-869.	0.9	89
10	Continuously elevated serum matrix metalloproteinase-3 for 3 ~ 6Âmonths predict one-year radiographi progression in rheumatoid arthritis: a prospective cohort study. Arthritis Research and Therapy, 2015, 17, 289.	c 1.6	34
11	Serum Matrix Metalloproteinase-3 as a Noninvasive Biomarker of Histological Synovitis for Diagnosis of Rheumatoid Arthritis. Mediators of Inflammation, 2014, 2014, 1-10.	1.4	57
12	Elevated Serum IgG4 Defines Specific Clinical Phenotype of Rheumatoid Arthritis. Mediators of Inflammation, 2014, 2014, 1-10.	1.4	30