

Per Pfeiffer

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3796294/per-pfeiffer-publications-by-citations.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140 papers	2,556 citations	26 h-index	47 g-index
153 ext. papers	3,385 ext. citations	4.7 avg, IF	5.01 L-index

#	Paper	IF	Citations
140	Encorafenib, Binimetinib, and Cetuximab in V600E-Mutated Colorectal Cancer. <i>New England Journal of Medicine</i> , 2019 , 381, 1632-1643	59.2	481
139	Efficacy and Safety of Pembrolizumab for Heavily Pretreated Patients With Advanced, Metastatic Adenocarcinoma or Squamous Cell Carcinoma of the Esophagus: The Phase 2 KEYNOTE-180 Study. <i>JAMA Oncology</i> , 2019 , 5, 546-550	13.4	225
138	Clinical trial enrollment, patient characteristics, and survival differences in prospectively registered metastatic colorectal cancer patients. <i>Cancer</i> , 2009 , 115, 4679-87	6.4	101
137	TAS-102 with or without bevacizumab in patients with chemorefractory metastatic colorectal cancer: an investigator-initiated, open-label, randomised, phase 2 trial. <i>Lancet Oncology</i> , 2020 , 21, 412-420	21.7	69
136	High BRAF Mutation Frequency and Marked Survival Differences in Subgroups According to KRAS/BRAF Mutation Status and Tumor Tissue Availability in a Prospective Population-Based Metastatic Colorectal Cancer Cohort. <i>PLoS ONE</i> , 2015 , 10, e0131046	3.7	64
135	Patterns of angiogenesis in nonsmall-cell lung carcinoma. <i>Cancer</i> , 2001 , 91, 1500-1509	6.4	64
134	Establishment and characterization of models of chemotherapy resistance in colorectal cancer: Towards a predictive signature of chemoresistance. <i>Molecular Oncology</i> , 2015 , 9, 1169-85	7.9	57
133	miR-345 in metastatic colorectal cancer: a non-invasive biomarker for clinical outcome in non-KRAS mutant patients treated with 3rd line cetuximab and irinotecan. <i>PLoS ONE</i> , 2014 , 9, e99886	3.7	54
132	A systematic review of salvage therapy to patients with metastatic colorectal cancer previously treated with fluorouracil, oxaliplatin and irinotecan +/- targeted therapy. <i>Cancer Treatment Reviews</i> , 2014 , 40, 701-15	14.4	53
131	Platinum-based cytotoxic therapy in basal cell carcinoma--a review of the literature. <i>Acta Oncologica</i> , 1996 , 35, 677-82	3.2	53
130	Persistent prevention of oxaliplatin-induced peripheral neuropathy using calmagafodipir (PledOx): a placebo-controlled randomised phase II study (PLIANT). <i>Acta Oncologica</i> , 2018 , 57, 393-402	3.2	52
129	Can we save the rectum by watchful waiting or transanal micro-surgery following (chemo) adjuvant therapy versus total mesorectal excision for early stage cancer (STAR-TREC study)? protocol for a multicentre, randomised feasibility study. <i>BMJ Open</i> , 2017 , 7, e019474	3	50
128	A randomized study of epirubicin at four different dose levels in advanced breast cancer. Feasibility of myelotoxicity prediction through single blood-sample measurement. <i>Cancer Chemotherapy and Pharmacology</i> , 1991 , 28, 465-9	3.5	48
127	Interleukin-6 and C-reactive protein as prognostic biomarkers in metastatic colorectal cancer. <i>Oncotarget</i> , 2016 , 7, 75013-75022	3.3	46
126	Peritoneal metastasis from pancreatic cancer treated with pressurized intraperitoneal aerosol chemotherapy (PIPAC). <i>Clinical and Experimental Metastasis</i> , 2017 , 34, 309-314	4.7	43
125	Prospective, single-center implementation and response evaluation of pressurized intraperitoneal aerosol chemotherapy (PIPAC) for peritoneal metastasis. <i>Therapeutic Advances in Medical Oncology</i> , 2018 , 10, 1758835918777036	5.4	42
124	Trends in cancer in the elderly population in Denmark, 1980-2012. <i>Acta Oncologica</i> , 2016 , 55 Suppl 1, 1-6	3.2	41

123	Clinical utility of KRAS status in circulating plasma DNA compared to archival tumour tissue from patients with metastatic colorectal cancer treated with anti-epidermal growth factor receptor therapy. <i>European Journal of Cancer</i> , 2015 , 51, 2678-85	7.5	40
122	Cetuximab in treatment of metastatic colorectal cancer: final survival analyses and extended RAS data from the NORDIC-VII study. <i>British Journal of Cancer</i> , 2017 , 116, 1271-1278	8.7	38
121	Tissue microRNAs as predictors of outcome in patients with metastatic colorectal cancer treated with first line Capecitabine and Oxaliplatin with or without Bevacizumab. <i>PLoS ONE</i> , 2014 , 9, e109430	3.7	37
120	Home parenteral nutrition increases fat free mass in patients with incurable gastrointestinal cancer. Results of a randomized controlled trial. <i>Clinical Nutrition</i> , 2019 , 38, 182-190	5.9	35
119	Cetuximab and irinotecan as third line therapy in patients with advanced colorectal cancer after failure of irinotecan, oxaliplatin and 5-fluorouracil. <i>Acta Oncologica</i> , 2007 , 46, 697-701	3.2	32
118	Prognostic role of carcinoembryonic antigen and carbohydrate antigen 19-9 in metastatic colorectal cancer: a BRAF-mutant subset with high CA 19-9 level and poor outcome. <i>British Journal of Cancer</i> , 2018 , 118, 1609-1616	8.7	31
117	Evaluation of Adjuvant Chemotherapy in Patients With Resected Pancreatic Cancer After Neoadjuvant FOLFIRINOX Treatment. <i>JAMA Oncology</i> , 2020 , 6, 1733-1740	13.4	29
116	Survival-associated heterogeneity of marker-defined perivascular cells in colorectal cancer. <i>Oncotarget</i> , 2016 , 7, 41948-41958	3.3	27
115	Optimizing the management of locally advanced pancreatic cancer with a focus on induction chemotherapy: Expert opinion based on a review of current evidence. <i>Cancer Treatment Reviews</i> , 2019 , 77, 1-10	14.4	26
114	Trends in colorectal cancer in the elderly in Denmark, 1980-2012. <i>Acta Oncologica</i> , 2016 , 55 Suppl 1, 29-39	3.2	25
113	Suramin in non-small cell lung cancer and advanced breast cancer. Two parallel phase II studies. <i>Acta Oncologica</i> , 1997 , 36, 171-4	3.2	24
112	Consequences of a high incidence of microsatellite instability and BRAF-mutated tumors: A population-based cohort of metastatic colorectal cancer patients. <i>Cancer Medicine</i> , 2019 , 8, 3623-3635	4.8	23
111	The potential diagnostic value of serum microRNA signature in patients with pancreatic cancer. <i>International Journal of Cancer</i> , 2016 , 139, 2312-24	7.5	23
110	A phase II study of Epirubicin in oxaliplatin-resistant patients with metastatic colorectal cancer and TOP2A gene amplification. <i>BMC Cancer</i> , 2016 , 16, 91	4.8	22
109	Risk factors for brain metastases in patients with metastatic colorectal cancer. <i>Acta Oncologica</i> , 2017 , 56, 639-645	3.2	20
108	Reduced-dose combination chemotherapy (S-1 plus oxaliplatin) versus full-dose monotherapy (S-1) in older vulnerable patients with metastatic colorectal cancer (NORDIC9): a randomised, open-label phase 2 trial. <i>The Lancet Gastroenterology and Hepatology</i> , 2019 , 4, 376-388	18.8	20
107	Short-time infusion of oxaliplatin (Eloxatin) in combination with capecitabine (Xeloda) in patients with advanced colorectal cancer. <i>Acta Oncologica</i> , 2003 , 42, 832-6	3.2	20
106	Trends in cancer of the liver, gall bladder, bile duct, and pancreas in elderly in Denmark, 1980-2012. <i>Acta Oncologica</i> , 2016 , 55 Suppl 1, 40-5	3.2	20

105	Clinical value of serum hyaluronan and propeptide of type III collagen in patients with pancreatic cancer. <i>International Journal of Cancer</i> , 2020 , 146, 2913-2922	7.5	19
104	Maintenance Therapy With Cetuximab Every Second Week in the First-Line Treatment of Metastatic Colorectal Cancer: The NORDIC-7.5 Study by the Nordic Colorectal Cancer Biomodulation Group. <i>Clinical Colorectal Cancer</i> , 2015 , 14, 170-6	3.8	18
103	FCGR polymorphisms and cetuximab efficacy in chemorefractory metastatic colorectal cancer: an international consortium study. <i>Gut</i> , 2015 , 64, 921-8	19.2	17
102	Pressurized IntraPeritoneal Aerosol Chemotherapy (PIPAC) as an outpatient procedure. <i>Pleura and Peritoneum</i> , 2018 , 3, 20180128	2	17
101	A systematic review of observational studies of trifluridine/tipiracil (TAS-102) for metastatic colorectal cancer. <i>Acta Oncologica</i> , 2019 , 58, 1149-1157	3.2	16
100	Pressurized IntraPeritoneal Aerosol Chemotherapy with one minute of electrostatic precipitation (ePIPAC) is feasible, but the histological tumor response in peritoneal metastasis is insufficient. <i>European Journal of Surgical Oncology</i> , 2020 , 46, 155-159	3.6	16
99	Treatment of peritoneal carcinomatosis with Pressurized IntraPeritoneal Aerosol Chemotherapy - PIPAC-OPC2. <i>Pleura and Peritoneum</i> , 2018 , 3, 20180108	2	16
98	Associations between primary tumor RAS, BRAF and PIK3CA mutation status and metastatic site in patients with chemo-resistant metastatic colorectal cancer. <i>Acta Oncologica</i> , 2018 , 57, 1057-1062	3.2	15
97	Cytotoxic treatment of metastatic breast cancer. Which drugs and drug combinations to use?. <i>Acta Oncologica</i> , 1992 , 31, 219-24	3.2	15
96	Plasma YKL-40 in patients with metastatic colorectal cancer treated with first line oxaliplatin-based regimen with or without cetuximab: RESULTS from the NORDIC VII Study. <i>PLoS ONE</i> , 2014 , 9, e87746	3.7	15
95	The effect of postoperative gemcitabine on overall survival in patients with resected pancreatic cancer: A nationwide population-based Danish register study. <i>Acta Oncologica</i> , 2019 , 58, 864-871	3.2	14
94	Bidirectional treatment of peritoneal metastasis with Pressurized IntraPeritoneal Aerosol Chemotherapy (PIPAC) and systemic chemotherapy: a systematic review. <i>BMC Cancer</i> , 2020 , 20, 105	4.8	14
93	High-dose radiotherapy and concurrent UFT plus l-leucovorin in locally advanced rectal cancer: a phase I trial. <i>Acta Oncologica</i> , 2005 , 44, 224-9	3.2	14
92	High RBM3 expression is associated with an improved survival and oxaliplatin response in patients with metastatic colorectal cancer. <i>PLoS ONE</i> , 2017 , 12, e0182512	3.7	14
91	CDX2: A Prognostic Marker in Metastatic Colorectal Cancer Defining a Better Mutated and a Worse Mutated Subgroup. <i>Frontiers in Oncology</i> , 2020 , 10, 8	5.3	13
90	Placebo-controlled phase II study of vitamin K3 cream for the treatment of cetuximab-induced rash. <i>Supportive Care in Cancer</i> , 2017 , 25, 2179-2185	3.9	12
89	Adjuvant Pressurized IntraPeritoneal Aerosol Chemotherapy (PIPAC) in resected high-risk colon cancer patients - study protocol for the PIPAC-OPC3 Trial. A prospective, controlled phase 2 Study. <i>Pleura and Peritoneum</i> , 2018 , 3, 20180107	2	12
88	Topoisomerase I copy number alterations as biomarker for irinotecan efficacy in metastatic colorectal cancer. <i>BMC Cancer</i> , 2017 , 17, 48	4.8	11

87	Molecular characterization of a large unselected cohort of metastatic colorectal cancers in relation to primary tumor location, rare metastatic sites and prognosis. <i>Acta Oncologica</i> , 2020 , 59, 417-426	3.2	11
86	Prognostic impact of Charlson's Age-Comorbidity Index and other risk factors in patients with pancreatic cancer. <i>European Journal of Cancer Care</i> , 2020 , 29, e13219	2.4	9
85	Pembrolizumab for patients with previously treated metastatic adenocarcinoma or squamous cell carcinoma of the esophagus: Phase 2 KEYNOTE-180 study.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 4049-4049	2.2	8
84	The changing face of treatment for metastatic colorectal cancer. <i>Expert Review of Anticancer Therapy</i> , 2019 , 19, 61-70	3.5	8
83	Tumor-specific genetic aberrations in cell-free DNA of gastroesophageal cancer patients. <i>Journal of Gastroenterology</i> , 2019 , 54, 108-121	6.9	8
82	Pre-treatment serum vitamin D deficiency is associated with increased inflammatory biomarkers and short overall survival in patients with pancreatic cancer. <i>European Journal of Cancer</i> , 2021 , 144, 72-80	7.5	8
81	Total cell-free DNA, carcinoembryonic antigen, and C-reactive protein for assessment of prognosis in patients with metastatic colorectal cancer. <i>Tumor Biology</i> , 2018 , 40, 1010428318811207	2.9	8
80	Initial treatment and survival in 4163 Danish patients with pancreatic cancer: A nationwide unselected real-world register study. <i>European Journal of Cancer</i> , 2020 , 129, 50-59	7.5	7
79	S-1 (Teysono) and gemcitabine in Caucasian patients with unresectable pancreatic adenocarcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2018 , 81, 573-578	3.5	7
78	Re: five-day oral etoposide treatment for advanced small-cell lung cancer: randomized comparison with intravenous chemotherapy. <i>Journal of the National Cancer Institute</i> , 1997 , 89, 1892-3	9.7	7
77	A pharmacokinetic study of prednimustine as compared with prednisolone plus chlorambucil in cancer patients. <i>Cancer Chemotherapy and Pharmacology</i> , 1991 , 28, 205-10	3.5	7
76	Angiogenesis Inhibitors for Colorectal Cancer. A Review of the Clinical Data. <i>Cancers</i> , 2021 , 13,	6.6	7
75	Can we predict toxicity and efficacy in older patients with cancer? Older patients with colorectal cancer as an example. <i>ESMO Open</i> , 2016 , 1, e000021	6	7
74	Personalized Clinical Decision Making in Gastrointestinal Malignancies: The Role of PET. <i>PET Clinics</i> , 2016 , 11, 273-83	2.2	7
73	Health-related quality of life in patients with metastatic colorectal cancer, association with systemic inflammatory response and RAS and BRAF mutation status. <i>European Journal of Cancer</i> , 2017 , 81, 26-35	7.5	6
72	Detection of free intraperitoneal tumour cells in peritoneal lavage fluid from patients with peritoneal metastasis before and after treatment with pressurised intraperitoneal aerosol chemotherapy (PIPAC). <i>Journal of Clinical Pathology</i> , 2019 , 72, 368-372	3.9	6
71	Randomized study comparing full dose monotherapy (S-1 followed by irinotecan) and reduced dose combination therapy (S-1/oxaliplatin followed by S-1/irinotecan) as initial therapy for older patients with metastatic colorectal cancer: NORDIC 9. <i>BMC Cancer</i> , 2017 , 17, 548	4.8	6
70	Intact and cleaved plasma soluble urokinase receptor in patients with metastatic colorectal cancer treated with oxaliplatin with or without cetuximab. <i>International Journal of Cancer</i> , 2015 , 137, 2470-7	7.5	6

69	Randomized study evaluating trifluridine/tipiracil (TAS-102) versus + trifluridine/tipiracil + bevacizumab as last-line therapy in patients with chemorefractory unresectable metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2019 , 37, 637-637	2.2	6
68	Spatial and phenotypic characterization of pancreatic cancer-associated fibroblasts after neoadjuvant treatment. <i>Histology and Histopathology</i> , 2020 , 35, 811-825	1.4	6
67	Experience with S-1 in older Caucasian patients with metastatic colorectal cancer (mCRC): Findings from an observational chart review. <i>Acta Oncologica</i> , 2016 , 55, 881-5	3.2	6
66	TIMP-1 is under regulation of the EGF signaling axis and promotes an aggressive phenotype in KRAS-mutated colorectal cancer cells: a potential novel approach to the treatment of metastatic colorectal cancer. <i>Oncotarget</i> , 2016 , 7, 59441-59457	3.3	5
65	Prognostic Value of Combined Detection of Serum IL6, YKL-40, and C-reactive Protein in Patients with Unresectable Pancreatic Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 176-184 ⁴		5
64	Trends in upper gastro-intestinal cancer among the elderly in Denmark, 1980-2012. <i>Acta Oncologica</i> , 2016 , 55 Suppl 1, 23-8	3.2	5
63	Treatment-related survival associations of claudin-2 expression in fibroblasts of colorectal cancer. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2018 , 472, 395-405 ^{5.1}		5
62	Mutational profiling and immunohistochemical analysis of a surgical series of ampullary carcinomas. <i>Journal of Clinical Pathology</i> , 2019 , 72, 762-770	3.9	4
61	Sustained-release metoclopramide plus methylprednisolone versus placebo plus methylprednisolone as antiemetic prophylaxis during non-cisplatin chemotherapy. A randomized double-blind cross-over trial. <i>Acta Oncologica</i> , 1996 , 35, 57-61	3.2	4
60	ABCG2 Protein Levels and Association to Response to First-Line Irinotecan-Based Therapy for Patients with Metastatic Colorectal Cancer. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	4
59	A randomized phase I/II study of everolimus, irinotecan, and cetuximab versus capecitabine and oxaliplatin in gemcitabine-resistant patients with pancreatic cancer.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 337-337	2.2	3
58	Current status of treatment of metastatic colorectal cancer with special reference to cetuximab and elderly patients. <i>OncoTargets and Therapy</i> , 2009 , 2, 17-27	4.4	3
57	Metastatic colorectal carcinomas with high SATB2 expression are associated with better prognosis and response to chemotherapy: a population-based Scandinavian study. <i>Acta Oncologica</i> , 2020 , 59, 284-290 ^{3.2}		3
56	Quality of Life in Vulnerable Older Patients with Metastatic Colorectal Cancer Receiving Palliative Chemotherapy-The Randomized NORDIC9-Study. <i>Cancers</i> , 2021 , 13,	6.6	3
55	Early F-FDG-PET/CT as a predictive marker for treatment response and survival in patients with metastatic colorectal cancer treated with irinotecan and cetuximab. <i>Acta Oncologica</i> , 2016 , 55, 1175-1182 ^{3.2}		3
54	Next-generation sequencing and histological response assessment in peritoneal metastasis from pancreatic cancer treated with PIPAC. <i>Journal of Clinical Pathology</i> , 2021 , 74, 19-24	3.9	3
53	Phase II study evaluating trifluridine/tipiracil + bevacizumab and capecitabine + bevacizumab in first-line unresectable metastatic colorectal cancer (mCRC) patients who are noneligible for intensive therapy (TASCO1): Results of the final analysis on the overall survival.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 14-14	2.2	3
52	S-1 in combination with docetaxel and oxaliplatin in patients with advanced gastro-esophageal adenocarcinoma: two parallel phase 1/2a studies. <i>Acta Oncologica</i> , 2017 , 56, 46-51	3.2	2

51	How to select colorectal cancer patients for personalized therapy. <i>EBioMedicine</i> , 2019 , 41, 36-37	8.8	2
50	Local recurrence of basal cell carcinoma and cisplatin containing chemotherapy. <i>Acta Oncologica</i> , 1997 , 36, 87	3.2	2
49	Maintenance therapy with biweekly cetuximab (C) in the first-line treatment of metastatic colorectal cancer (mCRC): The NORDIC 7.5 study (NCT00660582), by the Nordic Colorectal Cancer Biomodulation Group.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 3538-3538	2.2	2
48	The Global POLAR program: Two pivotal placebo-controlled studies of calmagafodipir used on top of modified FOLFOX6 to prevent chemotherapy-induced peripheral neuropathy (CIPN).. <i>Journal of Clinical Oncology</i> , 2019 , 37, TPS3616-TPS3616	2.2	2
47	QoL from TASCO1: Health-related quality of life of trifluridine/tipiracil-bevacizumab and capecitabine-bevacizumab as first-line treatments in metastatic colorectal cancer patients not eligible for intensive chemotherapy. Results from the TASCO1 phase II study.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 676-676	2.2	2
46	Feasibility of switching to S-1 after other fluoropyrimidine-related cardiotoxicity during chemotherapy for solid tumors.. <i>Journal of Clinical Oncology</i> , 2020 , 38, 7037-7037	2.2	2
45	Prognostic role of tumour-infiltrating lymphocytes and macrophages in relation to MSI, CDX2 and BRAF status: a population-based study of metastatic colorectal cancer patients. <i>British Journal of Cancer</i> , 2021 ,	8.7	2
44	Evaluation of the stage classification of anal cancer by the TNM 8th version versus the TNM 7th version. <i>Acta Oncologica</i> , 2020 , 59, 1016-1023	3.2	2
43	Predictive Value of Geriatric Oncology Screening and Geriatric Assessment in Older Patients with Solid Cancers: Protocol for a Danish prospective cohort study (PROGNOSIS-G8). <i>Journal of Geriatric Oncology</i> , 2021 , 12, 1270-1276	3.6	2
42	Frailty screening for predicting rapid functional decline, rapid progressive disease, and shorter overall survival in older patients with gastrointestinal cancer receiving palliative chemotherapy - a prospective, clinical study. <i>Journal of Geriatric Oncology</i> , 2021 , 12, 578-584	3.6	2
41	Primary tumor location and expression of mir-664 as a combined biomarker for bevacizumab effectiveness in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 3572-3572	2.2	1
40	Persistent prevention of CIPN using calmagafodipir (PledOx): Results from a placebo-controlled randomized phase II study (PLIANT) in patients with metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2016 , 34, 10018-10018	2.2	1
39	Pre-planned safety analysis of NORDIC 9: A randomized trial comparing full dose monotherapy (S-1) with reduced dose combination therapy (S-1/oxaliplatin) in older chemo-naïve patients with metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2017 , 35, 10032-10032	2.2	1
38	The prognostic value of serum CA 19-9 in patients with metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2017 , 35, e15131-e15131	2.2	1
37	The Global POLAR program: Calmagafodipir used on top of modified FOLFOX6 (5-FU/FA and oxaliplatin) to prevent chemotherapy induced peripheral neuropathy (CIPN).. <i>Journal of Clinical Oncology</i> , 2019 , 37, TPS722-TPS722	2.2	1
36	Chemotherapy for patients with non-resectable pancreatic cancer with additional chemo-radiotherapy for patients with potentially resectable tumours: Final Results.. <i>Journal of Clinical Oncology</i> , 2016 , 34, e15725-e15725	2.2	1
35	New use for old drugs: Epirubicin in colorectal cancer. <i>Acta Oncologica</i> , 2021 , 60, 954-956	3.2	1
34	Gene expression profiling of morphologic subtypes of pancreatic ductal adenocarcinoma using surgical and EUS-FNB specimens. <i>Pancreatology</i> , 2021 , 21, 530-543	3.8	1

33	Clinicopathological factors associated with tumour-specific mutation detection in plasma of patients with RAS-mutated or BRAF-mutated metastatic colorectal cancer. <i>International Journal of Cancer</i> , 2021 , 149, 1385-1397	7.5	1
32	Safety, tolerability and preliminary efficacy of CAN04, a first in class monoclonal antibody against IL1RAP, in combination with nab-paclitaxel and gemcitabine (NG) in subjects with pancreatic cancer.. <i>Journal of Clinical Oncology</i> , 2021 , 39, e16228-e16228	2.2	1
31	Cetuximab plus irinotecan administered biweekly with reduced infusion time to heavily pretreated patients with metastatic colorectal cancer and related RAS and BRAF mutation status. <i>International Journal of Cancer</i> , 2020 , 148, 2542	7.5	1
30	Predictive value of geriatric oncology screening and geriatric assessment of older patients with cancer: A randomized clinical trial protocol (PROGNOSIS-RCT). <i>Journal of Geriatric Oncology</i> , 2021 ,	3.6	1
29	-G12C Mutation in One Real-Life and Three Population-Based Nordic Cohorts of Metastatic Colorectal Cancer.. <i>Frontiers in Oncology</i> , 2022 , 12, 826073	5.3	1
28	Disparity in use of modern combination chemotherapy associated with facility type influences survival of 2655 patients with advanced pancreatic cancer. <i>Acta Oncologica</i> , 2021 , 1-9	3.2	1
27	Prognostic and diagnostic value of serum hyaluronan in patients with pancreatic carcinoma.. <i>Journal of Clinical Oncology</i> , 2018 , 36, e16249-e16249	2.2	0
26	Capecitabine in combination with panitumumab is not the treatment of choice in older patients with metastatic colorectal cancer. <i>Journal of Geriatric Oncology</i> , 2021 ,	3.6	0
25	TAS-102 plus bevacizumab in metastatic colorectal cancer - AuthorsSreply. <i>Lancet Oncology, The</i> , 2020 , 21, e227	21.7	
24	Response to the letter entitled: Re: Pre-treatment serum vitamin D deficiency is associated with increased inflammatory biomarkers and short overall survival in patients with pancreatic cancer: Analysis of the prognostic effect of serum vitamin D on pancreatic cancer: Several confounders. <i>European Journal of Cancer</i> , 2021 , 158, 349-358	7.5	
23	ISO-CC-005: A phase I/II study of Modufolin (MTHF) in combination with 5-FU, irinotecan, and oxaliplatin ± bevacizumab in patients with metastasizing colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 838-838	2.2	
22	Prognostic value of serum interleukin-6 and YKL-40 and systemic inflammatory response in patients with unresectable pancreatic cancer.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 267-267	2.2	
21	Prognostic significance of SATB1 expression in metastatic colorectal cancer: A Nordic prospective cohort study.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 707-707	2.2	
20	Expression of podocalyxin-like protein and epidermal growth factor receptor in metastatic colorectal cancer: Prognostic impact and relationship with response to cetuximab.. <i>Journal of Clinical Oncology</i> , 2018 , 36, e15587-e15587	2.2	
19	A randomized phase II study of second-line treatment with liposomal irinotecan, and S-1 versus liposomal irinotecan and 5-fluorouracil in gemcitabine-refractory metastatic pancreatic cancer patients.. <i>Journal of Clinical Oncology</i> , 2020 , 38, TPS4664-TPS4664	2.2	
18	Digitalized multiparametric analyses of tumor stroma for identification of low perivascular PDGFR expression and low vessel density as independent prognosis markers for stage IV CRC.. <i>Journal of Clinical Oncology</i> , 2014 , 32, e14525-e14525	2.2	
17	Benefit of EGFR-inhibition therapy for metastatic colorectal cancer patients with KRAS-mutated tumors and high plasma TIMP-1 level: Results from the NORDIC VII study.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3590-3590	2.2	
16	Efficacy and safety of S-1 and gemcitabine in an unselected Western cohort of patients with unresectable pancreatic cancer.. <i>Journal of Clinical Oncology</i> , 2015 , 33, e15258-e15258	2.2	

15	C-reactive protein and interleukin-6 as markers of systemic inflammatory response and as prognostic factors for metastatic colorectal cancer. Data from the randomized phase III NORDIC-VII study.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 3548-3548	2.2
14	Phase I dose-finding study of S-1 in combination with docetaxel and oxaliplatin (DOS) as first-line therapy in patients with advanced gastro-esophageal cancer.. <i>Journal of Clinical Oncology</i> , 2015 , 33, e15081-e15081	2.2
13	FOLFIRINOX for patients with borderline and never-resectable locally advanced pancreatic cancer, with the addition of chemoradiotherapy for potentially resectable patients: A phase II study.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 408-408	2.2
12	S-1 in combination with docetaxel and oxaliplatin (DOS) every 2 weeks (DOS2w) or every 3 weeks (DOS3w) in patients with advanced gastro-esophageal adenocarcinoma (aGEA): Final results of 2 parallel phase I studies.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 153-153	2.2
11	Outcome of cetuximab plus irinotecan in relation to RAS and BRAF mutational status in patients with colorectal cancer prior treated with a fluoropyrimidine, oxaliplatin and irinotecan.. <i>Journal of Clinical Oncology</i> , 2016 , 34, e15115-e15115	2.2
10	The prognostic value of serum IL-6 and YKL-40 in patients with metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2017 , 35, e15060-e15060	2.2
9	Plasma concentrations of YKL-40 in chemo-naïve patients with metastatic colorectal cancer treated with FLOX with or without cetuximab: Results from the NORDIC VII study.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 3548-3548	2.2
8	Biweekly cetuximab in combination with irinotecan as second-line treatment in patients with platinum-resistant gastroesophageal cancer (GEC).. <i>Journal of Clinical Oncology</i> , 2012 , 30, e14517-e14517 ²	2.2
7	Plasma levels of TIMP-1 in chemo-naïve patients with metastatic colorectal cancer treated with first-line FLOX with or without cetuximab: Results from the Nordic VII Study.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 392-392	2.2
6	Pretreatment plasma concentrations of YKL-40 and IL-6 in patients with pancreatic cancer: Potential diagnostic and prognostic biomarkers.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 164-164	2.2
5	Prognostic significance of tumor stromal and epithelial claudin 2 in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 3597-3597	2.2
4	Tumor perivascular PDGFBR as an independent prognostic factor in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 3571-3571	2.2
3	Trastuzumab with triple chemotherapy (DOC) in patients with advanced gastroesophageal cancer: A phase I dose-finding trial.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 108-108	2.2
2	Outcome after liver transplantation compared with chemotherapy in colorectal cancer patients with nonresectable liver-only disease.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 531-531	2.2
1	How to select cancer patients for immunotherapy. <i>EBioMedicine</i> , 2021 , 63, 103184	8.8