

Patrik Nasr

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3797900/publications.pdf>

Version: 2024-02-01

33
papers

5,448
citations

361296

20
h-index

414303

32
g-index

34
all docs

34
docs citations

34
times ranked

5716
citing authors

#	ARTICLE	IF	CITATIONS
1	Fibrosis stage is the strongest predictor for disease-specific mortality in NAFLD after up to 33 years of follow-up. <i>Hepatology</i> , 2015, 61, 1547-1554.	3.6	1,683
2	Increased risk of mortality by fibrosis stage in nonalcoholic fatty liver disease: Systematic review and meta-analysis. <i>Hepatology</i> , 2017, 65, 1557-1565.	3.6	1,294
3	Fibrosis stage but not NASH predicts mortality and time to development of severe liver disease in biopsy-proven NAFLD. <i>Journal of Hepatology</i> , 2017, 67, 1265-1273.	1.8	730
4	Association Between Fibrosis Stage and Outcomes of Patients With Nonalcoholic Fatty Liver Disease: A Systematic Review and Meta-Analysis. <i>Gastroenterology</i> , 2020, 158, 1611-1625.e12.	0.6	575
5	Risk for development of severe liver disease in lean patients with nonalcoholic fatty liver disease: A long-term follow-up study. <i>Hepatology Communications</i> , 2018, 2, 48-57.	2.0	200
6	Natural History of NAFLD/NASH. <i>Current Hepatology Reports</i> , 2017, 16, 391-397.	0.4	102
7	Natural history of nonalcoholic fatty liver disease: A prospective follow-up study with serial biopsies. <i>Hepatology Communications</i> , 2018, 2, 199-210.	2.0	102
8	Cardiovascular risk factors in non-alcoholic fatty liver disease. <i>Liver International</i> , 2019, 39, 197-204.	1.9	75
9	Accuracy of Noninvasive Scoring Systems in Assessing Risk of Death and Liver-Related Endpoints in Patients With Nonalcoholic Fatty Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2019, 17, 1148-1156.e4.	2.4	71
10	The European NAFLD Registry: A real-world longitudinal cohort study of nonalcoholic fatty liver disease. <i>Contemporary Clinical Trials</i> , 2020, 98, 106175.	0.8	71
11	Non-invasive tests accurately stratify patients with NAFLD based on their risk of liver-related events. <i>Journal of Hepatology</i> , 2022, 76, 1013-1020.	1.8	66
12	Low to moderate lifetime alcohol consumption is associated with less advanced stages of fibrosis in non-alcoholic fatty liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2017, 52, 159-165.	0.6	60
13	Elevated serum ferritin is associated with increased mortality in non-alcoholic fatty liver disease after 16 years of follow-up. <i>Liver International</i> , 2016, 36, 1688-1695.	1.9	54
14	Using a 3% Proton Density Fat Fraction as a Cut-Off Value Increases Sensitivity of Detection of Hepatic Steatosis, Based on Results From Histopathology Analysis. <i>Gastroenterology</i> , 2017, 153, 53-55.e7.	0.6	51
15	Moderate alcohol consumption is associated with advanced fibrosis in non-alcoholic fatty liver disease and shows a synergistic effect with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2021, 115, 154439.	1.5	41
16	Established and emerging factors affecting the progression of nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2020, 111, 154183.	1.5	39
17	SAF score and mortality in NAFLD after up to 41 years of follow-up. <i>Scandinavian Journal of Gastroenterology</i> , 2017, 52, 87-91.	0.6	32
18	The amount of liver fat predicts mortality and development of type 2 diabetes in non-alcoholic fatty liver disease. <i>Liver International</i> , 2020, 40, 1069-1078.	1.9	31

#	ARTICLE	IF	CITATIONS
19	Non-alcoholic fatty liver disease does not increase dementia risk although histology data might improve risk prediction. <i>JHEP Reports</i> , 2021, 3, 100218.	2.6	26
20	A Dynamic Aspartate to Alanine Aminotransferase Ratio Provides Valid Predictions of Incident Severe Liver Disease. <i>Hepatology Communications</i> , 2021, 5, 1021-1035.	2.0	23
21	Health Care Costs of Patients With Biopsy-Confirmed Nonalcoholic Fatty Liver Disease Are Nearly Twice Those of Matched Controls. <i>Clinical Gastroenterology and Hepatology</i> , 2020, 18, 1592-1599.e8.	2.4	21
22	Automated quantification of steatosis: agreement with stereological point counting. <i>Diagnostic Pathology</i> , 2017, 12, 80.	0.9	15
23	Biomarkers of liver fibrosis: prospective comparison of multimodal magnetic resonance, serum algorithms and transient elastography. <i>Scandinavian Journal of Gastroenterology</i> , 2020, 55, 848-859.	0.6	15
24	Low hepatic manganese concentrations in patients with hepatic steatosis – A cohort study of copper, iron and manganese in liver biopsies. <i>Journal of Trace Elements in Medicine and Biology</i> , 2021, 67, 126772.	1.5	15
25	Contrast-enhanced ultrasonography could be a non-invasive method for differentiating none or mild from severe fibrosis in patients with biopsy proven non-alcoholic fatty liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2016, 51, 1126-1132.	0.6	13
26	Non-invasive diagnosis and staging of non-alcoholic fatty liver disease. <i>Hormones</i> , 2022, 21, 349-368.	0.9	12
27	Modifiers of Liver-Related Manifestation in the Course of NAFLD. <i>Current Pharmaceutical Design</i> , 2020, 26, 1062-1078.	0.9	8
28	Repeated measurements of non-invasive fibrosis tests to monitor the progression of non-alcoholic fatty liver disease: A long-term follow-up study. <i>Liver International</i> , 2022, 42, 1545-1556.	1.9	6
29	Evaluating the prevalence and severity of NAFLD in primary care: the EPSONIP study protocol. <i>BMC Gastroenterology</i> , 2021, 21, 180.	0.8	5
30	Hepatic patatin-like phospholipase domain-containing 3 levels are increased in I148M risk allele carriers and correlate with NAFLD in humans. <i>Hepatology Communications</i> , 2022, 6, 2689-2701.	2.0	5
31	Serum levels of endotrophin are associated with nonalcoholic steatohepatitis. <i>Scandinavian Journal of Gastroenterology</i> , 2021, 56, 437-442.	0.6	4
32	Low awareness of non-alcoholic fatty liver disease in patients with type 2 diabetes in Swedish Primary Health Care. <i>Scandinavian Journal of Gastroenterology</i> , 2022, 57, 60-69.	0.6	3
33	Reply. <i>Hepatology</i> , 2016, 64, 310-311.	3.6	0