

# Jun Kameda

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3814472/publications.pdf>

Version: 2024-02-01

57  
papers

1,487  
citations

361413

20  
h-index

330143

37  
g-index

57  
all docs

57  
docs citations

57  
times ranked

1375  
citing authors

#	ARTICLE	IF	CITATIONS
1	Stick-slip behavior of a clayey crustal fault. <i>Physical Review Research</i> , 2022, 4, .	3.6	2
2	Rheological properties of halloysite soil slurry: a case study of weathered tephra involved in a shallow landslide triggered by the 2018 Eastern Iburi earthquake in Hokkaido, Japan. <i>Earth, Planets and Space</i> , 2022, 74, .	2.5	3
3	Rheological properties of concentrated allophane, halloysite, and kaolinite suspensions. <i>Applied Clay Science</i> , 2022, 226, 106557.	5.2	2
4	1-D inversion analysis of a shallow landslide triggered by the 2018 Eastern Iburi earthquake in Hokkaido, Japan. <i>Earth, Planets and Space</i> , 2021, 73, .	2.5	4
5	Mineralogical and physico-chemical properties of halloysite-bearing slip surface material from a landslide during the 2018 Eastern Iburi earthquake, Hokkaido. <i>Progress in Earth and Planetary Science</i> , 2021, 8, .	3.0	8
6	Influence of biopolymers on the rheological properties of seafloor sediments and the runout behavior of submarine debris flows. <i>Scientific Reports</i> , 2021, 11, 1493.	3.3	7
7	Generation of sintered fault rock and its implications for earthquake energetics and fault healing. <i>Communications Earth &amp; Environment</i> , 2020, 1, .	6.8	6
8	Cohesional Slip on a Plate Subduction Boundary During a Large Earthquake. <i>Geophysical Research Letters</i> , 2020, 47, e2020GL088395.	4.0	4
9	Fluidized landslides triggered by the liquefaction of subsurface volcanic deposits during the 2018 Iburi–Tobu earthquake, Hokkaido. <i>Scientific Reports</i> , 2019, 9, 13119.	3.3	33
10	Fault weakening caused by smectite swelling. <i>Earth, Planets and Space</i> , 2019, 71, .	2.5	10
11	Three-dimensional texture of natural pseudotachylyte: Pseudotachylyte formation mechanism in hydrous accretionary complex. <i>Island Arc</i> , 2018, 27, e12241.	1.1	0
12	Dehydroxylation Kinetics of Clay Minerals and Its Application to Friction Heating Along an Imbricate Thrust in an Accretionary Prism. <i>Geochemistry, Geophysics, Geosystems</i> , 2018, 19, 2991-3003.	2.5	10
13	Rheological properties of composite serpentine-brucite suspensions: Implications for mudflow behavior on forearc seamounts. <i>Marine Geology</i> , 2018, 403, 191-196.	2.1	9
14	Acoustic properties of deformed rocks in the Nobeoka thrust, in the Shimanto Belt, Kyushu, Southwest Japan. <i>Island Arc</i> , 2017, 26, e12198.	1.1	1
15	Temporal stress variations along a seismogenic megasplay fault in the subduction zone: An example from the Nobeoka Thrust, southwestern Japan. <i>Island Arc</i> , 2017, 26, e12193.	1.1	5
16	Alteration and dehydration of subducting oceanic crust within subduction zones: implications for dewatering step-down and plate-boundary seismogenesis. <i>Earth, Planets and Space</i> , 2017, 69, .	2.5	14
17	Opal-CT in chert beneath the toe of the Tohoku margin and its influence on the seismic aseismic transition in subduction zones. <i>Geophysical Research Letters</i> , 2017, 44, 687-693.	4.0	2
18	Sensitivity of Clay Suspension Rheological Properties to pH, Temperature, Salinity, and Smectite/Quartz Ratio. <i>Geophysical Research Letters</i> , 2017, 44, 9615-9621.	4.0	18

#	ARTICLE	IF	CITATIONS
19	Exchangeable cation composition of the smectite-rich plate boundary fault at the Japan Trench. <i>Geophysical Research Letters</i> , 2016, 43, 3112-3119.	4.0	7
20	Source and sink of fluid in pelagic siliceous sediments along a cold subduction plate boundary. <i>Tectonophysics</i> , 2016, 686, 146-157.	2.2	2
21	Hydrogeological responses to incoming materials at the erosional subduction margin, offshore Osa Peninsula, Costa Rica. <i>Geochemistry, Geophysics, Geosystems</i> , 2015, 16, 2725-2742.	2.5	11
22	Strength characteristics of Japan Trench borehole samples in the high-slip region of the 2011 Tohoku-Oki earthquake. <i>Earth and Planetary Science Letters</i> , 2015, 412, 35-41.	4.4	68
23	Pelagic smectite as an important factor in tsunamigenic slip along the Japan Trench. <i>Geology</i> , 2015, 43, 155-158.	4.4	65
24	Multiple damage zone structure of an exhumed seismogenic megasplay fault in a subduction zone - a study from the Nobeoka Thrust Drilling Project. <i>Earth, Planets and Space</i> , 2015, 67, .	2.5	15
25	Estimation of slip rate and fault displacement during shallow earthquake rupture in the Nankai subduction zone. <i>Earth, Planets and Space</i> , 2015, 67, .	2.5	15
26	Structure and lithology of the Japan Trench subduction plate boundary fault. <i>Tectonics</i> , 2015, 34, 53-69.	2.8	53
27	Frictional properties of sediments entering the Costa Rica subduction zone offshore the Osa Peninsula: implications for fault slip in shallow subduction zones. <i>Earth, Planets and Space</i> , 2014, 66, 72.	2.5	12
28	Changes in illite crystallinity within an ancient tectonic boundary thrust caused by thermal, mechanical, and hydrothermal effects: an example from the Nobeoka Thrust, southwest Japan. <i>Earth, Planets and Space</i> , 2014, 66, 116.	2.5	25
29	Quartz deposition and its influence on the deformation process of megathrusts in subduction zones. <i>Earth, Planets and Space</i> , 2014, 66, .	2.5	7
30	The influence of organic-rich shear zones on pelagic sediment deformation and seismogenesis in a subduction zone. <i>Journal of Mineralogical and Petrological Sciences</i> , 2014, 109, 228-238.	0.9	2
31	Structure and Composition of the Plate-Boundary Slip Zone for the 2011 Tohoku-Oki Earthquake. <i>Science</i> , 2013, 342, 1208-1211.	12.6	226
32	Low Coseismic Shear Stress on the Tohoku-Oki Megathrust Determined from Laboratory Experiments. <i>Science</i> , 2013, 342, 1211-1214.	12.6	220
33	Progress of illitization along an imbricate frontal thrust at shallow depths in an accretionary prism. <i>Tectonophysics</i> , 2013, 600, 41-51.	2.2	9
34	Hanging wall deformation of a seismogenic megasplay fault in an accretionary prism: The Nobeoka Thrust in southwestern Japan. <i>Journal of Structural Geology</i> , 2013, 52, 136-147.	2.3	25
35	Contrasts in physical properties between the hanging wall and footwall of an exhumed seismogenic megasplay fault in a subduction zone—An example from the Nobeoka Thrust Drilling Project. <i>Geochemistry, Geophysics, Geosystems</i> , 2013, 14, 5354-5370.	2.5	22
36	Importance of mechanochemical effects on fault slip behavior during earthquakes. <i>Geophysical Research Letters</i> , 2013, 40, 2988-2992.	4.0	24

#	ARTICLE	IF	CITATIONS
37	Tectonic mélange as fault rock of subduction plate boundary. <i>Tectonophysics</i> , 2012, 568-569, 25-38.	2.2	97
38	Silica diagenesis and its effect on interplate seismicity in cold subduction zones. <i>Earth and Planetary Science Letters</i> , 2012, 317-318, 136-144.	4.4	22
39	Runaway slip to the trench due to rupture of highly pressurized megathrust beneath the middle trench slope: The tsunamigenesis of the 2011 Tohoku earthquake off the east coast of northern Japan. <i>Earth and Planetary Science Letters</i> , 2012, 339-340, 32-45.	4.4	81
40	A new source of water in seismogenic subduction zones. <i>Geophysical Research Letters</i> , 2011, 38, n/a-n/a.	4.0	34
41	Smectite to chlorite conversion by frictional heating along a subduction thrust. <i>Earth and Planetary Science Letters</i> , 2011, 305, 161-170.	4.4	41
42	Low-grade metamorphism around the down-dip limit of seismogenic subduction zones: Example from an ancient accretionary complex in the Shimanto Belt, Japan. <i>Tectonophysics</i> , 2011, 502, 383-392.	2.2	16
43	Reproduction of thermal pressurization and fluidization of clay-rich fault gouges by high-velocity friction experiments and implications for seismic slip in natural faults. <i>Geological Society Special Publication</i> , 2011, 359, 267-285.	1.3	29
44	Smectite swelling in the Miura-Boso accretionary prism: Possible cause for incipient décollement zone formation. <i>Tectonophysics</i> , 2010, 494, 75-84.	2.2	8
45	Modification to the crystal structure of chlorite during early stages of its dissolution. <i>Physics and Chemistry of Minerals</i> , 2009, 36, 537-544.	0.8	15
46	Stacking faults with 180° layer rotation in celadonite, an Fe- and Mg-rich dioctahedral mica. <i>Clays and Clay Minerals</i> , 2008, 56, 612-621.	1.3	13
47	Polytype and morphological analyses of g <sub>1/4</sub> mbelite, a fibrous Mg-rich illite. <i>Clays and Clay Minerals</i> , 2007, 55, 453-466.	1.3	5
48	XRD and HRTEM analyses of stacking structures in sudoite, di-trioctahedral chlorite. <i>American Mineralogist</i> , 2007, 92, 1586-1592.	1.9	13
49	Novel 2:1 structure of phyllosilicates formed by annealing Fe <sup>3+</sup> , Mg-rich dioctahedral mica. <i>American Mineralogist</i> , 2007, 92, 1531-1534.	1.9	3
50	Morphological analyses of minute crystals by using stereo-photogrammetric scanning electron microscopy and electron back-scattered diffraction. <i>Journal of Microscopy</i> , 2007, 228, 358-365.	1.8	5
51	Stacking structures in pyrophyllite revealed by high-resolution transmission electron microscopy (HRTEM). <i>American Mineralogist</i> , 2006, 91, 1293-1299.	1.9	40
52	Stacking structure in disordered talc: Interpretation of its X-ray diffraction pattern by using pattern simulation and high-resolution transmission electron microscopy. <i>American Mineralogist</i> , 2006, 91, 1363-1370.	1.9	30
53	Dissolution of brucite on the (001) surface at neutral pH: <i>in situ</i> atomic force microscopy observations. <i>Clays and Clay Minerals</i> , 2006, 54, 598-604.	1.3	8
54	Morphological characteristics of ordered kaolinite: Investigation using electron back-scattered diffraction. <i>American Mineralogist</i> , 2005, 90, 1462-1465.	1.9	26

#	ARTICLE	IF	CITATIONS
55	H <sub>2</sub> generation during dry grinding of kaolinite. Journal of Colloid and Interface Science, 2004, 275, 225-228.	9.4	24
56	H <sub>2</sub> generation in wet grinding of granite and single-crystal powders and implications for H <sub>2</sub> concentration on active faults. Geophysical Research Letters, 2003, 30, .	4.0	29
57	Occurrences of Pseudotachylyte obtained from the Nojima fault at Nojima-Hirabayashi, Awaji Island, Japan. Journal of the Geological Society of Japan, 2002, 108, IX-X.	0.6	2