

Helene Eutamene

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3824031/publications.pdf>

Version: 2024-02-01

16
papers

909
citations

933447

10
h-index

996975

15
g-index

17
all docs

17
docs citations

17
times ranked

1412
citing authors

#	ARTICLE	IF	CITATIONS
1	Human milk oligosaccharides alleviate stress-induced visceral hypersensitivity and associated microbiota dysbiosis. <i>Journal of Nutritional Biochemistry</i> , 2022, 99, 108865.	4.2	7
2	Lactose and Fructo-oligosaccharides Increase Visceral Sensitivity in Mice via Glycation Processes, Increasing Mast Cell Density in Colonic Mucosa. <i>Gastroenterology</i> , 2020, 158, 652-663.e6.	1.3	36
3	The Infant-Derived <i>Bifidobacterium bifidum</i> Strain CNCM I-4319 Strengthens Gut Functionality. <i>Microorganisms</i> , 2020, 8, 1313.	3.6	10
4	The multicomponent medication Spascupreel attenuates stress-induced gut dysfunction in rats. <i>Neurogastroenterology and Motility</i> , 2020, 32, e13798.	3.0	3
5	Effects of thermized donkey milk with lysozyme activity on altered gut barrier in mice exposed to water-avoidance stress. <i>Journal of Dairy Science</i> , 2019, 102, 7697-7706.	3.4	14
6	Donkey milk consumption exerts anti-inflammatory properties by normalizing antimicrobial peptides levels in Paneth's cells in a model of ileitis in mice. <i>European Journal of Nutrition</i> , 2018, 57, 155-166.	3.9	40
7	The role of mucoprotectants in the management of gastrointestinal disorders. <i>Expert Review of Gastroenterology and Hepatology</i> , 2018, 12, 83-90.	3.0	21
8	5-oxoETE triggers nociception in constipation-predominant irritable bowel syndrome through MAS-related G protein-coupled receptor D. <i>Science Signaling</i> , 2018, 11, .	3.6	44
9	Evaluation of reticulated gelatin-hibiscus-propolis against intestinal commensal species commonly associated with urinary tract infections. <i>Future Microbiology</i> , 2017, 12, 505-513.	2.0	8
10	Modifications of mesenteric adipose tissue during moderate experimental colitis in mice. <i>Life Sciences</i> , 2014, 94, 1-7.	4.3	9
11	76 Congenitally Elevated Gut Permeability Is Linked to Basal Visceral Hyposensitivity but Stress-Induced Visceral Hypersensitivity in CA-MLCK Mice. <i>Gastroenterology</i> , 2014, 146, S-21.	1.3	1
12	Prevention of gut leakiness by a probiotic treatment leads to attenuated HPA response to an acute psychological stress in rats. <i>Psychoneuroendocrinology</i> , 2012, 37, 1885-1895.	2.7	495
13	Synergy between <i>Lactobacillus paracasei</i> and Its Bacterial Products to Counteract Stress-Induced Gut Permeability and Sensitivity Increase in Rats. <i>Journal of Nutrition</i> , 2007, 137, 1901-1907.	2.9	135
14	Acute Stress Modulates the Histamine Content of Mast Cells in the Gastrointestinal Tract Through Interleukin-1 and Corticotropin-Releasing Factor Release in Rats. <i>Journal of Physiology</i> , 2003, 553, 959-966.	2.9	62
15	Chronic ingestion of a potential food contaminant induces gastrointestinal inflammation in rats: role of nitric oxide and mast cells. <i>Digestive Diseases and Sciences</i> , 2000, 45, 1842-1849.	2.3	12
16	<i>Lactococcus lactis</i> NCDO2118 exerts visceral antinociceptive properties in rat via GABA production in the gastro-intestinal tract. <i>ELife</i> , 0, 11, .	6.0	12