

Patrocinio Molinero Hueso

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3826381/patrocinio-molinero-hueso-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

29
papers

642
citations

13
h-index

25
g-index

29
ext. papers

685
ext. citations

6
avg, IF

2.8
L-index

#	Paper	IF	Citations
29	Donor-specific circulating cell free DNA as a noninvasive biomarker of graft injury in heart transplantation. <i>Clinica Chimica Acta</i> , 2019 , 495, 590-597	6.2	18
28	Temporal expression patterns of the melatonergic system in the human thymus of children. <i>Molecular Metabolism</i> , 2019 , 28, 83-90	8.8	6
27	Noninvasive prenatal diagnosis by cell-free DNA screening for fetomaternal HPA-1a platelet incompatibility. <i>Transfusion</i> , 2018 , 58, 2272-2279	2.9	13
26	Non-invasive Prenatal Diagnosis of Feto-Maternal Platelet Incompatibility by Cold High Resolution Melting Analysis. <i>Advances in Experimental Medicine and Biology</i> , 2016 , 924, 67-70	3.6	5
25	Detection of p53 Mutations in Circulating DNA of Transplanted Hepatocellular Carcinoma Patients as a Biomarker of Tumor Recurrence. <i>Advances in Experimental Medicine and Biology</i> , 2016 , 924, 25-28	3.6	6
24	Screening of KRAS Mutation in Pre- and Post-Surgery Serum of Patients Suffering from Colon Cancer by COLD-PCR HRM. <i>Advances in Experimental Medicine and Biology</i> , 2016 , 924, 39-41	3.6	3
23	Role of early cell-free DNA levels decrease as a predictive marker of fatal outcome after severe traumatic brain injury. <i>Clinica Chimica Acta</i> , 2012 , 414, 12-7	6.2	67
22	Blocking of melatonin synthesis and MT(1) receptor impairs the activation of Jurkat T cells. <i>Cellular and Molecular Life Sciences</i> , 2010 , 67, 3163-72	10.3	20
21	Evidence of immune system melatonin production by two pineal melatonin deficient mice, C57BL/6 and Swiss strains. <i>Journal of Pineal Research</i> , 2009 , 47, 15-22	10.4	41
20	Treatment with testosterone or estradiol in melatonin treated females and males MRL/MpJ-Faslpr mice induces negative effects in developing systemic lupus erythematosus. <i>Journal of Pineal Research</i> , 2008 , 45, 204-11	10.4	10
19	Evidence for melatonin synthesis in the rat brain during development. <i>Journal of Pineal Research</i> , 2007 , 42, 240-6	10.4	53
18	Melatonin biosynthesis in the thymus of humans and rats. <i>Cellular and Molecular Life Sciences</i> , 2007 , 64, 781-90	10.3	59
17	Sex-dependent effect of melatonin on systemic erythematosus lupus developed in Mrl/Mpj-Faslpr mice: it ameliorates the disease course in females, whereas it exacerbates it in males. <i>Endocrinology</i> , 2006 , 147, 1717-24	4.8	27
16	Dual effect of melatonin as proinflammatory and antioxidant in collagen-induced arthritis in rats. <i>Journal of Pineal Research</i> , 2005 , 38, 93-9	10.4	29
15	Melatonin synthesis and melatonin-membrane receptor (MT1) expression during rat thymus development: role of the pineal gland. <i>Journal of Pineal Research</i> , 2005 , 39, 77-83	10.4	38
14	Melatonin prevents hyperhomocysteinemia and neural lipid peroxidation induced by methionine intake. <i>Current Neurovascular Research</i> , 2005 , 2, 175-8	1.8	12
13	Melatonin triggers Crohn's disease symptoms. <i>Journal of Pineal Research</i> , 2002 , 32, 277-8	10.4	31

12	Melatonin is responsible for the nocturnal increase observed in serum and thymus of thymosin alpha1 and thymulin concentrations: observations in rats and humans. <i>Journal of Neuroimmunology</i> , 2000 , 103, 180-8	3.5	49
11	Circadian variations in the rat serum total antioxidant status: correlation with melatonin levels. <i>Journal of Pineal Research</i> , 1998 , 25, 1-4	10.4	52
10	Nocturnal increases in the triiodothyronine/thyroxine ratio in the rat thymus and pineal gland follow increases of type II 5Xdeiodinase activity. <i>International Journal of Biochemistry and Cell Biology</i> , 1998 , 30, 235-41	5.6	7
9	Different experimental conditions which regulate type II 5Xdeiodinase mRNA in rat Harderian gland. <i>Life Sciences</i> , 1997 , 61, 181-92	6.8	3
8	Beta- and alpha-adrenergic mechanisms are involved in regulating type II thyroxine 5Xdeiodinase in rat thymus. <i>Life Sciences</i> , 1996 , 58, 1-8	6.8	5
7	Characterization of binding sites for beta-adrenergic agonists and vasoactive intestinal peptide in the rat harderian gland. <i>Microscopy Research and Technique</i> , 1996 , 34, 139-43	2.8	2
6	Expression of type II thyroxine 5Xdeiodinase from rat harderian gland in <i>Xenopus laevis</i> oocytes. <i>FEBS Letters</i> , 1994 , 354, 110-2	3.8	2
5	Thyroxine type II 5Xdeiodinase activity in pineal and Harderian gland is enhanced by hypothyroidism but is independent of serum thyroxine concentrations during hyperthyroidism. <i>International Journal of Biochemistry & Cell Biology</i> , 1993 , 25, 1041-6		5
4	In vivo activation of pineal N-acetyltransferase but not type II thyroxine 5Xdeiodinase by phenylephrine in young rats. <i>Neuroscience Letters</i> , 1991 , 127, 13-5	3.3	5
3	Decreased binding of vasoactive intestinal peptide to intestinal epithelial cells from hypothyroid rats. <i>Biochemical and Biophysical Research Communications</i> , 1989 , 162, 701-7	3.4	2
2	Interaction of vasoactive intestinal peptide (VIP) with rat lymphoid cells. <i>Peptides</i> , 1986 , 7, 177-81	3.8	60
1	The interaction of vasoactive intestinal peptide (VIP) with isolated bovine thyroid plasma membranes. <i>Biochemical and Biophysical Research Communications</i> , 1985 , 128, 1336-41	3.4	12