

# Faïk Errouissi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3833566/publications.pdf>

Version: 2024-02-01

13  
papers

673  
citations

1040056

9  
h-index

1281871

11  
g-index

13  
all docs

13  
docs citations

13  
times ranked

791  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | A Review on the Toxicity and Non-Target Effects of Macrocyclic Lactones in Terrestrial and Aquatic Environments. <i>Current Pharmaceutical Biotechnology</i> , 2012, 13, 1004-1060.   | 1.6 | 260       |
| 2  | Use of anthelmintics in herbivores and evaluation of risks for the non target fauna of pastures. <i>Veterinary Research</i> , 2002, 33, 547-562.  | 3.0 | 136       |
| 3  | Comparative effects of conventional and no-tillage management on some soil properties under Mediterranean semi-arid conditions in northwestern Tunisia. <i>Soil and Tillage Research</i> , 2010, 106, 247-253.  | 5.6 | 64        |
| 4  | Relative efficiency of extensive grazing vs. wild ungulates management for dung beetle conservation in a heterogeneous landscape from Southern Europe (Scarabaeinae, Aphodiinae, Geotrupinae). <i>Biological Conservation</i> , 2008, 141, 2879-2887.           | 4.1 | 47        |
| 5  | Effects of the Attractiveness for Dung Beetles of Dung Pat Origin and Size Along a Climatic Gradient. <i>Environmental Entomology</i> , 2004, 33, 45-53.  | 1.4 | 46        |
| 6  | Composition and Structure of Dung Beetle (Coleoptera: Aphodiidae, Geotrupidae, Scarabaeidae) Assemblages in Mountain Grasslands of the Southern Alps. <i>Annals of the Entomological Society of America</i> , 2004, 97, 701-709.                                | 2.5 | 37        |
| 7  | Soil invertebrates in durum wheat ( <i>Triticum durum</i> L.) cropping system under Mediterranean semi arid conditions: A comparison between conventional and no-tillage management. <i>Soil and Tillage Research</i> , 2011, 112, 122-132.                     | 5.6 | 34        |
| 8  | Seasonal occurrence and local coexistence within scarabaeid dung beetle guilds (Coleoptera: Tj ETQq0 0 0 rgBT /Ovgrlock 10, Tf 50 462   | 1.2 | 22        |
| 9  | Spatial and Temporal Variation in Species Composition, Diversity, and Structure of Mediterranean Dung Beetle Assemblages (Coleoptera: Scarabaeidae) Across a Bioclimatic Gradient. <i>Environmental Entomology</i> , 2012, 41, 785-801.                         | 1.4 | 11        |
| 10 | Dung beetle assemblages organization in two contrasted areas of the Mediterranean region: affinities and divergences. <i>Annales De La Societe Entomologique De France</i> , 2011, 47, 402-417.   | 0.9 | 8         |
| 11 | Consequences of habitat change in euromediterranean landscapes on the composition and diversity of dung beetle assemblages (Coleoptera, Scarabaeoidea). <i>Journal of Insect Conservation</i> , 2019, 23, 15-28.  | 1.4 | 8         |
| 12 | Diversity and structure of dung beetle assemblages under two contrasted habitats in Tunisia: oases vs. humid pastures. <i>Austral Entomology</i> , 2017, 56, 54-63.   | 1.4 | 0         |
| 13 | First data on the daily and seasonal activity patterns of <i>Stomoxys calcitrans</i> (Diptera: Muscidae) under Mediterranean semiarid climate in a dairy cattle farm in Tunisia. <i>International Journal of Tropical Insect Science</i> , 2022, 42, 1437-1447. | 1.0 | 0         |