

Soňa Legartová

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3837116/publications.pdf>

Version: 2024-02-01

32
papers

582
citations

567281

15
h-index

642732

23
g-index

33
all docs

33
docs citations

33
times ranked

843
citing authors

#	ARTICLE	IF	CITATIONS
1	The Highest Density of Phosphorylated Histone H1 Appeared in Prophase and Prometaphase in Parallel with Reduced H3K9me3, and HDAC1 Depletion Increased H1.2/H1.3 and H1.4 Serine 38 Phosphorylation. <i>Life</i> , 2022, 12, 798.	2.4	2
2	The SC-35 Splicing Factor Interacts with RNA Pol II and A-Type Lamin Depletion Weakens This Interaction. <i>Cells</i> , 2021, 10, 297.	4.1	2
3	A device for investigation of natural cell mobility and deformability. <i>Electrophoresis</i> , 2020, 41, 1238-1244.	2.4	3
4	N6-Adenosine Methylation in RNA and a Reduced m3G/TMG Level in Non-Coding RNAs Appear at Microirradiation-Induced DNA Lesions. <i>Cells</i> , 2020, 9, 360.	4.1	36
5	Cell differentiation and aging accompanied by depletion of the ACE2 protein. <i>Aging</i> , 2020, 12, 22495-22508.	3.1	11
6	DNA Damage Changes Distribution Pattern and Levels of HP1 Protein Isoforms in the Nucleolus and Increases Phosphorylation of HP1 ^{Î²} -Ser88. <i>Cells</i> , 2019, 8, 1097.	4.1	10
7	HDAC1 and HDAC3 underlie dynamic H3K9 acetylation during embryonic neurogenesis and in schizophrenia-like animals. <i>Journal of Cellular Physiology</i> , 2018, 233, 530-548.	4.1	61
8	Depletion of A-type lamins and <i>Lap2^{Î±}</i> reduces 53BP1 accumulation at UV _A -induced DNA lesions and <i>Lap2^{Î±}</i> protein is responsible for compactness of irradiated chromatin. <i>Journal of Cellular Biochemistry</i> , 2018, 119, 8146-8162.	2.6	10
9	H3K9me3 and H4K20me3 represent the epigenetic landscape for 53BP1 binding to DNA lesions. <i>Aging</i> , 2018, 10, 2585-2605.	3.1	27
10	PCNA is recruited to irradiated chromatin in late S-phase and is most pronounced in G2 phase of the cell cycle. <i>Protoplasma</i> , 2017, 254, 2035-2043.	2.1	15
11	Function of heterochromatin protein 1 during DNA repair. <i>Protoplasma</i> , 2017, 254, 1233-1240.	2.1	19
12	Mutations in the TP53 gene affected recruitment of 53BP1 protein to DNA lesions, but level of 53BP1 was stable after ^{Î³} -irradiation that depleted MDC1 protein in specific TP53 mutants. <i>Histochemistry and Cell Biology</i> , 2017, 148, 239-255.	1.7	13
13	An Endogenously Tagged Fluorescent Fusion Protein Library in Mouse Embryonic Stem Cells. <i>Stem Cell Reports</i> , 2017, 9, 1304-1314.	4.8	19
14	Advanced Confocal Microscopy Techniques to Study Protein-protein Interactions and Kinetics at DNA Lesions. <i>Journal of Visualized Experiments</i> , 2017, , .	0.3	7
15	Localized Movement and Levels of 53BP1 Protein Are Changed by ^{Î³} -irradiation in PML Deficient Cells. <i>Journal of Cellular Biochemistry</i> , 2016, 117, 2583-2596.	2.6	7
16	Advanced Image Acquisition and Analytical Techniques for Studies of Living Cells and Tissue Sections. <i>Microscopy and Microanalysis</i> , 2016, 22, 326-341.	0.4	4
17	The level and distribution pattern of HP1 ^{Î²} in the embryonic brain correspond to those of H3K9me1/me2 but not of H3K9me3. <i>Histochemistry and Cell Biology</i> , 2016, 145, 447-461.	1.7	7
18	Distinct kinetics of DNA repair protein accumulation at DNA lesions and cell cycle-dependent formation of ^{Î³} H2AX and NBS1-positive repair foci. <i>Biology of the Cell</i> , 2015, 107, 440-454.	2.0	24

#	ARTICLE	IF	CITATIONS
19	Localized movement and morphology of UBF1-positive nucleolar regions are changed by \hat{I}^3 -irradiation in G2 phase of the cell cycle. <i>Nucleus</i> , 2015, 6, 301-313.	2.2	9
20	Post-translational Modifications of Histones in Human Sperm. <i>Journal of Cellular Biochemistry</i> , 2015, 116, 2195-2209.	2.6	27
21	Coilin is rapidly recruited to UVA-induced DNA lesions and \hat{I}^3 -radiation affects localized movement of Cajal bodies. <i>Nucleus</i> , 2014, 5, 269-277.	2.2	22
22	Cell differentiation along multiple pathways accompanied by changes in histone acetylation status. <i>Biochemistry and Cell Biology</i> , 2014, 92, 85-93.	2.0	9
23	HP1 \hat{I}^2 -dependent recruitment of UBF1 to irradiated chromatin occurs simultaneously with CPDs. <i>Epigenetics and Chromatin</i> , 2014, 7, 39.	3.9	18
24	Nuclear Structures Surrounding Internal Lamin Invaginations. <i>Journal of Cellular Biochemistry</i> , 2014, 115, 476-487.	2.6	25
25	Basic nuclear processes affected by histone acetyltransferases and histone deacetylase inhibitors. <i>Epigenomics</i> , 2013, 5, 379-396.	2.1	28
26	Epigenetic aspects of HP1 exchange kinetics in apoptotic chromatin. <i>Biochimie</i> , 2013, 95, 167-179.	2.6	10
27	DNA-damage response in chromatin of ribosomal genes and the surrounding genome. <i>Gene</i> , 2013, 522, 156-167.	2.2	21
28	Acetylation-dependent nuclear arrangement and recruitment of BMI1 protein to UV-damaged chromatin. <i>Journal of Cellular Physiology</i> , 2012, 227, 1838-1850.	4.1	48
29	Effects of epigenetic-based anti-cancer drugs in leukaemia and multiple myeloma cells. <i>Cell Biology International</i> , 2011, 35, 1195-1203.	3.0	8
30	Recruitment of Oct4 Protein to UV-Damaged Chromatin in Embryonic Stem Cells. <i>PLoS ONE</i> , 2011, 6, e27281.	2.5	45
31	Chromocentre integrity and epigenetic marks. <i>Journal of Structural Biology</i> , 2010, 169, 124-133.	2.8	16
32	Nuclear organization of PML bodies in leukaemic and multiple myeloma cells. <i>Leukemia Research</i> , 2008, 32, 1866-1877.	0.8	19