

Qiuhe Ji

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3837295/publications.pdf>

Version: 2024-02-01

58
papers

4,692
citations

331538

21
h-index

161767

54
g-index

59
all docs

59
docs citations

59
times ranked

5873
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Prevalence of Diabetes among Men and Women in China. <i>New England Journal of Medicine</i> , 2010, 362, 1090-1101. | 13.9 | 2,685 |
| 2 | Standards of medical care for type 2 diabetes in China 2019. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3158. | 1.7 | 404 |
| 3 | Standards of care for type 2 diabetes in China. <i>Diabetes/Metabolism Research and Reviews</i> , 2016, 32, 442-458. | 1.7 | 236 |
| 4 | Primacy of the 3B Approach to Control Risk Factors for Cardiovascular Disease in Type 2 Diabetes Patients. <i>American Journal of Medicine</i> , 2013, 126, 925.e11-925.e22. | 0.6 | 174 |
| 5 | Acarbose compared with metformin as initial therapy in patients with newly diagnosed type 2 diabetes: an open-label, non-inferiority randomised trial. <i>Lancet Diabetes and Endocrinology</i> , 2014, 2, 46-55. | 5.5 | 134 |
| 6 | Impact of Waist Circumference and Body Mass Index on Risk of Cardiometabolic Disorder and Cardiovascular Disease in Chinese Adults: A National Diabetes and Metabolic Disorders Survey. <i>PLoS ONE</i> , 2013, 8, e57319. | 1.1 | 130 |
| 7 | The association between obstructive sleep apnea and metabolic syndrome: a systematic review and meta-analysis. <i>BMC Pulmonary Medicine</i> , 2015, 15, 105. | 0.8 | 98 |
| 8 | Type 2 diabetes mellitus unawareness, prevalence, trends and risk factors: National Health and Nutrition Examination Survey (NHANES) 1999-2010. <i>Journal of International Medical Research</i> , 2017, 45, 594-609. | 0.4 | 78 |
| 9 | Fasting triglycerides and glucose index is more suitable for the identification of metabolically unhealthy individuals in the Chinese adult population: A nationwide study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1050-1058. | 1.1 | 63 |
| 10 | BMI and waist circumference are associated with impaired glucose metabolism and type 2 diabetes in normal weight Chinese adults. <i>Journal of Diabetes and Its Complications</i> , 2014, 28, 470-476. | 1.2 | 43 |
| 11 | A phase 3, multicenter, randomized, allopurinol-controlled study assessing the safety and efficacy of oral febuxostat in Chinese gout patients with hyperuricemia. <i>International Journal of Rheumatic Diseases</i> , 2015, 18, 669-678. | 0.9 | 36 |
| 12 | Non-alcoholic fatty liver disease predicts type 2 diabetes mellitus, but not prediabetes, in Chinese adults: a five-year cohort study. <i>Liver International</i> , 2015, 35, 2401-2407. | 1.9 | 36 |
| 13 | The characteristics of newly diagnosed adult early-onset diabetes: a population-based cross-sectional study. <i>Scientific Reports</i> , 2017, 7, 46534. | 1.6 | 34 |
| 14 | Metabolic syndrome and chronic kidney disease in general Chinese adults: Results from the 2007-2008 China National Diabetes and Metabolic Disorders Study. <i>Clinica Chimica Acta</i> , 2014, 430, 115-120. | 0.5 | 31 |
| 15 | Association between Family History Risk Categories and Prevalence of Diabetes in Chinese Population. <i>PLoS ONE</i> , 2015, 10, e0117044. | 1.1 | 31 |
| 16 | Urban, semi-urban and rural difference in the prevalence of metabolic syndrome in Shaanxi province, northwestern China: a population-based survey. <i>BMC Public Health</i> , 2014, 14, 104. | 1.2 | 29 |
| 17 | Association between uric acid levels and cardiovascular outcomes and death in patients with type 2 diabetes: A subanalysis of EMPA-REG OUTCOME. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 1207-1214. | 2.2 | 29 |
| 18 | Effectiveness and safety of Bifidobacterium and berberine in human hyperglycemia and their regulatory effect on the gut microbiota: a multi-center, double-blind, randomized, parallel-controlled study. <i>Genome Medicine</i> , 2021, 13, 125. | 3.6 | 28 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Gender Differences in the Prevalence and Development of Metabolic Syndrome in Chinese Population with Abdominal Obesity. <i>PLoS ONE</i> , 2013, 8, e78270. | 1.1 | 27 |
| 20 | Obesity-Related Genomic Loci Are Associated with Type 2 Diabetes in a Han Chinese Population. <i>PLoS ONE</i> , 2014, 9, e104486. | 1.1 | 25 |
| 21 | Identification of autoimmune type 1 diabetes and multiple organ-specific autoantibodies in adult-onset non-insulin-requiring diabetes in China: A population-based multicentre nationwide survey. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 893-902. | 2.2 | 24 |
| 22 | <p>Prevalence and identification of type 1 diabetes in Chinese adults with newly diagnosed diabetes</p>. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2019, Volume 12, 1527-1541. | 1.1 | 23 |
| 23 | The Fecal Microbiota Is Already Altered in Normoglycemic Individuals Who Go on to Have Type 2 Diabetes. <i>Frontiers in Cellular and Infection Microbiology</i> , 2021, 11, 598672. | 1.8 | 23 |
| 24 | Efficacy and safety of metformin and sitagliptin based triple antihyperglycemic therapy (STRATEGY): a multicenter, randomized, controlled, non-inferiority clinical trial. <i>Science China Life Sciences</i> , 2017, 60, 225-238. | 2.3 | 20 |
| 25 | Recommendations for revision of Chinese diagnostic criteria for metabolic syndrome: A nationwide study. <i>Journal of Diabetes</i> , 2018, 10, 232-239. | 0.8 | 20 |
| 26 | Effectiveness and safety of bifidobacteria and berberine in people with hyperglycemia: study protocol for a randomized controlled trial. <i>Trials</i> , 2018, 19, 72. | 0.7 | 20 |
| 27 | Normal weight obesity and the risk of diabetes in Chinese people: a 9-year population-based cohort study. <i>Scientific Reports</i> , 2021, 11, 6090. | 1.6 | 17 |
| 28 | Association between Obstructive Sleep Apnea and Type 2 Diabetes Mellitus: A Dose-Response Meta-Analysis. <i>Evidence-based Complementary and Alternative Medicine</i> , 2021, 2021, 1-14. | 0.5 | 16 |
| 29 | Regional differences in diabetes prevalence and awareness between coastal and interior provinces in China: a population-based cross-sectional study. <i>BMC Public Health</i> , 2013, 13, 299. | 1.2 | 15 |
| 30 | Efficacy and safety of alogliptin in patients with type 2 diabetes mellitus: A multicentre randomized double-blind placebo-controlled Phase 3 study in mainland China, Taiwan, and Hong Kong. <i>Journal of Diabetes</i> , 2017, 9, 386-395. | 0.8 | 15 |
| 31 | Cigarette Smoking Is Associated with a Lower Prevalence of Newly Diagnosed Diabetes Screened by OGTT than Non-Smoking in Chinese Men with Normal Weight. <i>PLoS ONE</i> , 2016, 11, e0149234. | 1.1 | 15 |
| 32 | Body fat percentage cutoffs for risk of cardiometabolic abnormalities in the Chinese adult population: a nationwide study. <i>European Journal of Clinical Nutrition</i> , 2018, 72, 728-735. | 1.3 | 14 |
| 33 | Gender-differential effects on blood glucose levels between acarbose and metformin in Chinese patients with newly diagnosed type 2 diabetes: a sub-analysis of the MARCH trial. <i>Endocrine Journal</i> , 2021, 68, 69-79. | 0.7 | 10 |
| 34 | Evaluation of effectiveness of treatment paradigm for newly diagnosed type 2 diabetes patients in China: A nationwide prospective cohort study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 151-161. | 1.1 | 9 |
| 35 | Rfamide-related peptide-3 promotes alpha TC1 clone 6 cell survival likely via GPR147. <i>Peptides</i> , 2018, 107, 39-44. | 1.2 | 8 |
| 36 | Observational study evaluating the effectiveness of physician-targeted education for improving glycemic management of patients with type 2 diabetes (BEYOND II). <i>Journal of Diabetes</i> , 2020, 12, 66-76. | 0.8 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Three-component non-invasive risk score for undiagnosed diabetes in Chinese people: Development, validation and longitudinal evaluation. <i>Journal of Diabetes Investigation</i> , 2020, 11, 341-348. | 1.1 | 8 |
| 38 | Insulin resistance is not independently associated with chronic kidney disease in Chinese population: A population-based cross-sectional study. <i>Clinica Chimica Acta</i> , 2015, 448, 232-237. | 0.5 | 7 |
| 39 | Efficacy and safety of a needle-free injector in Chinese patients with type 2 diabetes mellitus treated with basal insulin: a multicentre, prospective, randomised, crossover study. <i>Expert Opinion on Drug Delivery</i> , 2019, 16, 995-1002. | 2.4 | 7 |
| 40 | Serum glycated albumin as good biomarker for predicting type 2 diabetes: A retrospective cohort study of China National Diabetes and Metabolic Disorders Survey. <i>Diabetes/Metabolism Research and Reviews</i> , 2021, , e3477. | 1.7 | 7 |
| 41 | Effectiveness of clinical alternatives to nerve conduction studies for screening for diabetic distal symmetrical polyneuropathy: A multi-center study. <i>Diabetes Research and Clinical Practice</i> , 2016, 115, 150-156. | 1.1 | 6 |
| 42 | Hypoglycaemia, Abnormal Lipids, and Cardiovascular Disease among Chinese with Type 2 Diabetes. <i>BioMed Research International</i> , 2015, 2015, 1-8. | 0.9 | 5 |
| 43 | Comparison of exenatide with biphasic insulin aspart 30 on glucose variability in type 2 diabetes: study protocol for a randomized controlled trial. <i>Trials</i> , 2016, 17, 160. | 0.7 | 5 |
| 44 | Differences between the perspectives of physicians and patients on the potential barriers to optimal diabetes control in China: a multicenter study. <i>BMC Health Services Research</i> , 2018, 18, 961. | 0.9 | 5 |
| 45 | Agreement Between the JDCCG, Revised NCEP-ATPIII, and IDF Definitions of Metabolic Syndrome in a Northwestern Chinese Population. <i>Diabetes Therapy</i> , 2018, 9, 1457-1468. | 1.2 | 5 |
| 46 | Screening strategy for islet autoantibodies in diabetes patients of different ages. <i>Diabetes Technology and Therapeutics</i> , 2021, , . | 2.4 | 5 |
| 47 | GAD65 Antibody Epitopes and Genetic Background in Latent Autoimmune Diabetes in Youth (LADY). <i>Frontiers in Immunology</i> , 2022, 13, 836952. | 2.2 | 5 |
| 48 | The Unmet Medical Needs of Current Injectable Antidiabetic Therapies in China: Patient and Health Care Professional Perspectives. <i>Clinical Therapeutics</i> , 2020, 42, 1549-1563. | 1.1 | 3 |
| 49 | Comparison of Blood Glucose Variability Between Exenatide and Biphasic Insulin Aspart 30 in Chinese Participants with Type 2 Diabetes Inadequately Controlled with Metformin Monotherapy: A Multicenter, Open-Label, Randomized Trial. <i>Diabetes Therapy</i> , 2020, 11, 2313-2328. | 1.2 | 3 |
| 50 | Efficacy and safety of metformin and sitagliptin-based dual and triple therapy in elderly Chinese patients with type 2 diabetes: Subgroup analysis of STRATEGY study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1532-1541. | 1.1 | 3 |
| 51 | Efficacy and safety of alogliptin versus acarbose in Chinese type 2 diabetes patients with high cardiovascular risk or coronary heart disease treated with aspirin and inadequately controlled with metformin monotherapy or drug-naïve: A multicentre, randomized, open-label, prospective study (ACADEMIC). <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 991-999. | 2.2 | 3 |
| 52 | Using noninvasive anthropometric indices to develop and validate a predictive model for metabolic syndrome in Chinese adults: a nationwide study. <i>BMC Endocrine Disorders</i> , 2022, 22, 53. | 0.9 | 3 |
| 53 | Islet Function Changes Among the Elderly Population. <i>Archives of Medical Research</i> , 2019, 50, 468-475. | 1.5 | 1 |
| 54 | Relationship between thyroid hormones and metabolic syndrome in a normal thyroid function population in Western China. <i>Chinese Medical Journal</i> , 2021, Publish Ahead of Print, 350-352. | 0.9 | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Pancreatic Safety of Once-Weekly Dulaglutide in Chinese Patients with Type 2 Diabetes Mellitus: Subgroup Analysis by Potential Influencing Factors. <i>Diabetes Therapy</i> , 2021, 12, 2677-2690. | 1.2 | 1 |
| 56 | Physical Activity, Sedentary Behavior, and the Risk of Cardiovascular Disease in Type 2 Diabetes Mellitus Patients: The MIDiab Study. <i>Engineering</i> , 2023, 20, 26-35. | 3.2 | 1 |
| 57 | Response to Serum cytokeratinâ€18 fragment levels predict development of type 2 diabetes mellitus in adult patients with NAFLD. <i>Liver International</i> , 2015, 35, 2622-2622. | 1.9 | 0 |
| 58 | Positive Selection of Natural Poly-Reactive B Cells in the Periphery Occurs Independent of Heavy Chain Allelic Inclusion. <i>PLoS ONE</i> , 2015, 10, e0125747. | 1.1 | 0 |