

Qiuhe Ji

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3837295/publications.pdf>

Version: 2024-02-01

58
papers

4,692
citations

331538

21
h-index

161767

54
g-index

59
all docs

59
docs citations

59
times ranked

5873
citing authors

#	ARTICLE	IF	CITATIONS
1	Physical Activity, Sedentary Behavior, and the Risk of Cardiovascular Disease in Type 2 Diabetes Mellitus Patients: The MIDiab Study. <i>Engineering</i> , 2023, 20, 26-35.	3.2	1
2	Efficacy and safety of alogliptin versus acarbose in Chinese type 2 diabetes patients with high cardiovascular risk or coronary heart disease treated with aspirin and inadequately controlled with metformin monotherapy or drug-naïve: A multicentre, randomized, open-label, prospective study (ACADEMIC). <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 991-999.	2.2	3
3	GAD65 Antibody Epitopes and Genetic Background in Latent Autoimmune Diabetes in Youth (LADY). <i>Frontiers in Immunology</i> , 2022, 13, 836952.	2.2	5
4	Using noninvasive anthropometric indices to develop and validate a predictive model for metabolic syndrome in Chinese adults: a nationwide study. <i>BMC Endocrine Disorders</i> , 2022, 22, 53.	0.9	3
5	The Fecal Microbiota Is Already Altered in Normoglycemic Individuals Who Go on to Have Type 2 Diabetes. <i>Frontiers in Cellular and Infection Microbiology</i> , 2021, 11, 598672.	1.8	23
6	Normal weight obesity and the risk of diabetes in Chinese people: a 9-year population-based cohort study. <i>Scientific Reports</i> , 2021, 11, 6090.	1.6	17
7	Relationship between thyroid hormones and metabolic syndrome in a normal thyroid function population in Western China. <i>Chinese Medical Journal</i> , 2021, Publish Ahead of Print, 350-352.	0.9	1
8	Serum glycated albumin as good biomarker for predicting type 2 diabetes: A retrospective cohort study of China National Diabetes and Metabolic Disorders Survey. <i>Diabetes/Metabolism Research and Reviews</i> , 2021, , e3477.	1.7	7
9	Effectiveness and safety of Bifidobacterium and berberine in human hyperglycemia and their regulatory effect on the gut microbiota: a multi-center, double-blind, randomized, parallel-controlled study. <i>Genome Medicine</i> , 2021, 13, 125.	3.6	28
10	Pancreatic Safety of Once-Weekly Dulaglutide in Chinese Patients with Type 2 Diabetes Mellitus: Subgroup Analysis by Potential Influencing Factors. <i>Diabetes Therapy</i> , 2021, 12, 2677-2690.	1.2	1
11	Association between Obstructive Sleep Apnea and Type 2 Diabetes Mellitus: A Dose-Response Meta-Analysis. <i>Evidence-based Complementary and Alternative Medicine</i> , 2021, 2021, 1-14.	0.5	16
12	Gender-differential effects on blood glucose levels between acarbose and metformin in Chinese patients with newly diagnosed type 2 diabetes: a sub-analysis of the MARCH trial. <i>Endocrine Journal</i> , 2021, 68, 69-79.	0.7	10
13	Screening strategy for islet autoantibodies in diabetes patients of different ages. <i>Diabetes Technology and Therapeutics</i> , 2021, , .	2.4	5
14	Observational study evaluating the effectiveness of physician-targeted education for improving glycemic management of patients with type 2 diabetes (BEYOND II). <i>Journal of Diabetes</i> , 2020, 12, 66-76.	0.8	8
15	Evaluation of effectiveness of treatment paradigm for newly diagnosed type 2 diabetes patients in China: A nationwide prospective cohort study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 151-161.	1.1	9
16	Three-component non-invasive risk score for undiagnosed diabetes in Chinese people: Development, validation and longitudinal evaluation. <i>Journal of Diabetes Investigation</i> , 2020, 11, 341-348.	1.1	8
17	The Unmet Medical Needs of Current Injectable Antidiabetic Therapies in China: Patient and Health Care Professional Perspectives. <i>Clinical Therapeutics</i> , 2020, 42, 1549-1563.	1.1	3
18	Comparison of Blood Glucose Variability Between Exenatide and Biphasic Insulin Aspart 30 in Chinese Participants with Type 2 Diabetes Inadequately Controlled with Metformin Monotherapy: A Multicenter, Open-Label, Randomized Trial. <i>Diabetes Therapy</i> , 2020, 11, 2313-2328.	1.2	3

#	ARTICLE	IF	CITATIONS
19	Association between uric acid levels and cardio-renal outcomes and death in patients with type 2 diabetes: A subanalysis of EMPA-REG OUTCOME. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 1207-1214.	2.2	29
20	Efficacy and safety of metformin and sitagliptin-based dual and triple therapy in elderly Chinese patients with type 2 diabetes: Subgroup analysis of STRATEGY study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1532-1541.	1.1	3
21	Efficacy and safety of a needle-free injector in Chinese patients with type 2 diabetes mellitus treated with basal insulin: a multicentre, prospective, randomised, crossover study. <i>Expert Opinion on Drug Delivery</i> , 2019, 16, 995-1002.	2.4	7
22	<p>Prevalence and identification of type 1 diabetes in Chinese adults with newly diagnosed diabetes</p>. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2019, Volume 12, 1527-1541.	1.1	23
23	Standards of medical care for type 2 diabetes in China 2019. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3158.	1.7	404
24	Islet Function Changes Among the Elderly Population. <i>Archives of Medical Research</i> , 2019, 50, 468-475.	1.5	1
25	Fasting triglycerides and glucose index is more suitable for the identification of metabolically unhealthy individuals in the Chinese adult population: A nationwide study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1050-1058.	1.1	63
26	Identification of autoimmune type 1 diabetes and multiple organ-specific autoantibodies in adult-onset non-insulin-requiring diabetes in China: A population-based multicentre nationwide survey. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 893-902.	2.2	24
27	Body fat percentage cutoffs for risk of cardiometabolic abnormalities in the Chinese adult population: a nationwide study. <i>European Journal of Clinical Nutrition</i> , 2018, 72, 728-735.	1.3	14
28	Recommendations for revision of Chinese diagnostic criteria for metabolic syndrome: A nationwide study. <i>Journal of Diabetes</i> , 2018, 10, 232-239.	0.8	20
29	Differences between the perspectives of physicians and patients on the potential barriers to optimal diabetes control in China: a multicenter study. <i>BMC Health Services Research</i> , 2018, 18, 961.	0.9	5
30	Agreement Between the JCDCC, Revised NCEP-ATPIII, and IDF Definitions of Metabolic Syndrome in a Northwestern Chinese Population. <i>Diabetes Therapy</i> , 2018, 9, 1457-1468.	1.2	5
31	Effectiveness and safety of bifidobacteria and berberine in people with hyperglycemia: study protocol for a randomized controlled trial. <i>Trials</i> , 2018, 19, 72.	0.7	20
32	RFamide-related peptide-3 promotes alpha TC1 clone 6 cell survival likely via GPR147. <i>Peptides</i> , 2018, 107, 39-44.	1.2	8
33	Efficacy and safety of metformin and sitagliptin based triple antihyperglycemic therapy (STRATEGY): a multicenter, randomized, controlled, non-inferiority clinical trial. <i>Science China Life Sciences</i> , 2017, 60, 225-238.	2.3	20
34	The characteristics of newly diagnosed adult early-onset diabetes: a population-based cross-sectional study. <i>Scientific Reports</i> , 2017, 7, 46534.	1.6	34
35	Type 2 diabetes mellitus unawareness, prevalence, trends and risk factors: National Health and Nutrition Examination Survey (NHANES) 1999-2010. <i>Journal of International Medical Research</i> , 2017, 45, 594-609.	0.4	78
36	Efficacy and safety of alogliptin in patients with type 2 diabetes mellitus: A multicentre randomized double-blind placebo-controlled Phase 3 study in mainland China, Taiwan, and Hong Kong. <i>Journal of Diabetes</i> , 2017, 9, 386-395.	0.8	15

#	ARTICLE	IF	CITATIONS
37	Effectiveness of clinical alternatives to nerve conduction studies for screening for diabetic distal symmetrical polyneuropathy: A multi-center study. <i>Diabetes Research and Clinical Practice</i> , 2016, 115, 150-156.	1.1	6
38	Standards of care for type 2 diabetes in China. <i>Diabetes/Metabolism Research and Reviews</i> , 2016, 32, 442-458.	1.7	236
39	Comparison of exenatide with biphasic insulin aspart 30 on glucose variability in type 2 diabetes: study protocol for a randomized controlled trial. <i>Trials</i> , 2016, 17, 160.	0.7	5
40	Cigarette Smoking Is Associated with a Lower Prevalence of Newly Diagnosed Diabetes Screened by OGTT than Non-Smoking in Chinese Men with Normal Weight. <i>PLoS ONE</i> , 2016, 11, e0149234.	1.1	15
41	The association between obstructive sleep apnea and metabolic syndrome: a systematic review and meta-analysis. <i>BMC Pulmonary Medicine</i> , 2015, 15, 105.	0.8	98
42	A phase 3, multicenter, randomized, allopurinol-controlled study assessing the safety and efficacy of oral febuxostat in Chinese gout patients with hyperuricemia. <i>International Journal of Rheumatic Diseases</i> , 2015, 18, 669-678.	0.9	36
43	Non-alcoholic fatty liver disease predicts type 2 diabetes mellitus, but not prediabetes, in Chinese men: a five-year cohort study. <i>Liver International</i> , 2015, 35, 2401-2407.	1.9	36
44	Response to Serum cytokeratin-18 fragment levels predict development of type 2 diabetes mellitus in adult patients with NAFLD. <i>Liver International</i> , 2015, 35, 2622-2622.	1.9	0
45	Positive Selection of Natural Poly-Reactive B Cells in the Periphery Occurs Independent of Heavy Chain Allelic Inclusion. <i>PLoS ONE</i> , 2015, 10, e0125747.	1.1	0
46	Hypoglycaemia, Abnormal Lipids, and Cardiovascular Disease among Chinese with Type 2 Diabetes. <i>BioMed Research International</i> , 2015, 2015, 1-8.	0.9	5
47	Insulin resistance is not independently associated with chronic kidney disease in Chinese population: A population-based cross-sectional study. <i>Clinica Chimica Acta</i> , 2015, 448, 232-237.	0.5	7
48	Association between Family History Risk Categories and Prevalence of Diabetes in Chinese Population. <i>PLoS ONE</i> , 2015, 10, e0117044.	1.1	31
49	Obesity-Related Genomic Loci Are Associated with Type 2 Diabetes in a Han Chinese Population. <i>PLoS ONE</i> , 2014, 9, e104486.	1.1	25
50	BMI and waist circumference are associated with impaired glucose metabolism and type 2 diabetes in normal weight Chinese adults. <i>Journal of Diabetes and Its Complications</i> , 2014, 28, 470-476.	1.2	43
51	Metabolic syndrome and chronic kidney disease in general Chinese adults: Results from the 2007-08 China National Diabetes and Metabolic Disorders Study. <i>Clinica Chimica Acta</i> , 2014, 430, 115-120.	0.5	31
52	Urban, semi-urban and rural difference in the prevalence of metabolic syndrome in Shaanxi province, northwestern China: a population-based survey. <i>BMC Public Health</i> , 2014, 14, 104.	1.2	29
53	Acarbose compared with metformin as initial therapy in patients with newly diagnosed type 2 diabetes: an open-label, non-inferiority randomised trial. <i>Lancet Diabetes and Endocrinology</i> , 2014, 2, 46-55.	5.5	134
54	Primacy of the 3B Approach to Control Risk Factors for Cardiovascular Disease in Type 2 Diabetes Patients. <i>American Journal of Medicine</i> , 2013, 126, 925.e11-925.e22.	0.6	174

#	ARTICLE	IF	CITATIONS
55	Regional differences in diabetes prevalence and awareness between coastal and interior provinces in China: a population-based cross-sectional study. BMC Public Health, 2013, 13, 299.	1.2	15
56	Impact of Waist Circumference and Body Mass Index on Risk of Cardiometabolic Disorder and Cardiovascular Disease in Chinese Adults: A National Diabetes and Metabolic Disorders Survey. PLoS ONE, 2013, 8, e57319.	1.1	130
57	Gender Differences in the Prevalence and Development of Metabolic Syndrome in Chinese Population with Abdominal Obesity. PLoS ONE, 2013, 8, e78270.	1.1	27
58	Prevalence of Diabetes among Men and Women in China. New England Journal of Medicine, 2010, 362, 1090-1101.	13.9	2,685