
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3840858/publications.pdf

Version: 2024-02-01



Ιμηγανιίμι

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Comparison of Retzius-sparing and conventional robot-assisted laparoscopic radical prostatectomy regarding continence and sexual function: an updated meta-analysis. Prostate Cancer and Prostatic Diseases, 2022, 25, 47-54. | 2.0 | 17 |
| 2 | Metabolomics analysis of human plasma reveals decreased production of trimethylamine Nâ€oxide retards the progression of chronic kidney disease. British Journal of Pharmacology, 2022, 179, 4344-4359. | 2.7 | 15 |
| 3 | Identification and characterization of isocitrate dehydrogenase 1 (IDH1) as a functional target of marine natural product grincamycin B. Acta Pharmacologica Sinica, 2021, 42, 801-813. | 2.8 | 5 |
| 4 | <i>Fusobacterium Nucleatum</i> Promotes the Development of Colorectal Cancer by Activating a Cytochrome P450/Epoxyoctadecenoic Acid Axis via TLR4/Keap1/NRF2 Signaling. Cancer Research, 2021, 81, 4485-4498. | 0.4 | 75 |
| 5 | The Functions of Cytochrome P450 ω-hydroxylases and the Associated Eicosanoids in Inflammation-Related Diseases. Frontiers in Pharmacology, 2021, 12, 716801. | 1.6 | 25 |
| 6 | Pleiotropic Functions of Cytochrome P450 Monooxygenase-Derived Eicosanoids in Cancer. Frontiers in Pharmacology, 2020, 11, 580897. | 1.6 | 11 |
| 7 | A retrospective analysis reveals a predictor of survival for the patient with paraquat intoxication. Clinica Chimica Acta, 2020, 511, 269-277. | 0.5 | 4 |
| 8 | A retrospective cross-sectional study of the associated factors of hyperuricemia in patients with chronic kidney disease. Journal of International Medical Research, 2020, 48, 030006052091922. | 0.4 | 1 |
| 9 | Higher BMP Expression in Tendon Stem/Progenitor Cells Contributes to the Increased Heterotopic Ossification in Achilles Tendon With Aging. Frontiers in Cell and Developmental Biology, 2020, 8, 570605. | 1.8 | 18 |
| 10 | <i>N</i> -Benzyl-linoleamide, a Constituent of <i>Lepidium meyenii</i> (Maca), Is an Orally Bioavailable Soluble Epoxide Hydrolase Inhibitor That Alleviates Inflammatory Pain. Journal of Natural Products, 2020, 83, 3689-3697. | 1.5 | 9 |
| 11 | Anti-inflammatory treatment with a soluble epoxide hydrolase inhibitor attenuates seizures and epilepsy-associated depression in the LiCl-pilocarpine post-status epilepticus rat model. Brain, Behavior, and Immunity, 2019, 81, 535-544. | 2.0 | 30 |
| 12 | Inhibition of soluble epoxide hydrolase attenuates a high-fat diet-mediated renal injury by activating PAX2 and AMPK. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 5154-5159. | 3.3 | 33 |
| 13 | Targeted Metabolomics Identifies the Cytochrome P450 Monooxygenase Eicosanoid Pathway as a Novel Therapeutic Target of Colon Tumorigenesis. Cancer Research, 2019, 79, 1822-1830. | 0.4 | 45 |
| 14 | Metabolic profiling of human plasma reveals the activation of 5-lipoxygenase in the acute attack of gouty arthritis. Rheumatology, 2019, 58, 345-351. | 0.9 | 21 |
| 15 | Lipidomic profiling reveals soluble epoxide hydrolase as a therapeutic target of obesity-induced colonic inflammation. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, 5283-5288. | 3.3 | 59 |
| 16 | Plasma profiling of amino acids distinguishes acute gout from asymptomatic hyperuricemia. Amino Acids, 2018, 50, 1539-1548. | 1.2 | 25 |
| 17 | Oxylipin profiling of human plasma reflects the renal dysfunction in uremic patients. Metabolomics, 2018, 14, 104. | 1.4 | 11 |
| 18 | Inhibition of Soluble Epoxide Hydrolase for Renal Health. Frontiers in Pharmacology, 2018, 9, 1551. | 1.6 | 23 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Lipidomic Profiling Reveals Soluble Epoxide Hydrolase As a Therapeutic Target of Obesityâ€induced Colonic Inflammation. FASEB Journal, 2018, 32, 560.1. | 0.2 | 0 |
| 20 | Inhibition of soluble epoxide hydrolase attenuates the kidney injury caused by ischemia/reperfusion in a murine model of acute kidney injury involved in GSKâ€3β phosphorylation. FASEB Journal, 2018, 32, 561.11. | 0.2 | 0 |
| 21 | A sensitive and accurate method to simultaneously measure uric acid and creatinine in human saliva by using LC–MS/MS. Bioanalysis, 2017, 9, 1751-1760. | 0.6 | 22 |
| 22 | Epoxide metabolites of arachidonate and docosahexaenoate function conversely in acute kidney injury involved in GSK3β signaling. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, 12608-12613. | 3.3 | 45 |
| 23 | A novel antimicrobial substance produced by Lactobacillus rhamnous LS8. Food Control, 2017, 73, 754-760. | 2.8 | 22 |
| 24 | Inhibition of mutant KrasG12D -initiated murine pancreatic carcinoma growth by a dual c-Raf and soluble epoxide hydrolase inhibitor t -CUPM. Cancer Letters, 2016, 371, 187-193. | 3.2 | 12 |
| 25 | Screening of soluble epoxide hydrolase inhibitory ingredients from traditional Chinese medicines for anti-inflammatory use. Journal of Ethnopharmacology, 2016, 194, 475-482. | 2.0 | 8 |
| 26 | Ornithine is a key mediator in hyperphosphatemia-mediated human umbilical vein endothelial cell apoptosis: Insights gained from metabolomics. Life Sciences, 2016, 146, 73-80. | 2.0 | 6 |
| 27 | Anti-phytopathogenic activity of sporothriolide, a metabolite from endophyte Nodulisporium sp. A21 in Ginkgo biloba. Pesticide Biochemistry and Physiology, 2016, 129, 7-13. | 1.6 | 37 |
| 28 | Inhibition of Chronic Pancreatitis and Murine Pancreatic Intraepithelial Neoplasia by a Dual Inhibitor of c-RAF and Soluble Epoxide Hydrolase in LSL-KrasG¹²D/Pdx-1-Cre Mice. Anticancer Research, 2016, 36, 27-37. | 0.5 | 14 |
| 29 | The Volume Ratio of Ground Glass Opacity in Early Lung CT Predicts Mortality in Acute Paraquat Poisoning. PLoS ONE, 2015, 10, e0121691. | 1.1 | 18 |
| 30 | In vitro and in vivo metabolism of N-adamantyl substituted urea-based soluble epoxide hydrolase inhibitors. Biochemical Pharmacology, 2015, 98, 718-731. | 2.0 | 14 |
| 31 | Biological evaluation of a novel sorafenib analogue, t-CUPM. Cancer Chemotherapy and Pharmacology, 2015, 75, 161-171. | 1.1 | 14 |
| 32 | Soluble Epoxide Hydrolase Inhibitor Attenuates Inflammation and Airway Hyperresponsiveness in Mice. American Journal of Respiratory Cell and Molecular Biology, 2015, 52, 46-55. | 1.4 | 45 |
| 33 | Dual inhibition of cyclooxygenase-2 and soluble epoxide hydrolase synergistically suppresses primary tumor growth and metastasis. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 11127-11132. | 3.3 | 84 |
| 34 | Optimized Inhibitors of Soluble Epoxide Hydrolase Improve in Vitro Target Residence Time and in Vivo Efficacy. Journal of Medicinal Chemistry, 2014, 57, 7016-7030. | 2.9 | 81 |
| 35 | Design, Synthesis, Antifungal, and Antioxidant Activities of (<i>E</i>)-6-((2-Phenylhydrazono)methyl)quinoxaline Derivatives. Journal of Agricultural and Food Chemistry, 2014, 62, 9637-9643. | 2.4 | 74 |
| 36 | Flavipin in Chaetomium globosum CDW7, an endophytic fungus from Ginkgo biloba, contributes to antioxidant activity. Applied Microbiology and Biotechnology, 2013, 97, 7131-7139. | 1.7 | 42 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Substituted phenyl groups improve the pharmacokinetic profile and anti-inflammatory effect of urea-based soluble epoxide hydrolase inhibitors in murine models. European Journal of Pharmaceutical Sciences, 2013, 48, 619-627. | 1.9 | 62 |
| 38 | Epoxygenated fatty acid profile and soluble epoxide hydrolase (sEH) activity in healthy and laminitic horses. Journal of Equine Veterinary Science, 2013, 33, 868. | 0.4 | 0 |
| 39 | Unique mechanistic insights into the beneficial effects of soluble epoxide hydrolase inhibitors in the prevention of cardiac fibrosis. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 5618-5623. | 3.3 | 85 |
| 40 | Anti-inflammatory Effects of ï‰-3 Polyunsaturated Fatty Acids and Soluble Epoxide Hydrolase Inhibitors in Angiotensin-Il–Dependent Hypertension. Journal of Cardiovascular Pharmacology, 2013, 62, 285-297. | 0.8 | 92 |
| 41 | Preparation of 20-HETE using multifunctional enzyme type 2-negative Starmerella bombicola. Journal of Lipid Research, 2013, 54, 3215-3219. | 2.0 | 12 |
| 42 | Epoxy metabolites of docosahexaenoic acid (DHA) inhibit angiogenesis, tumor growth, and metastasis. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 6530-6535. | 3.3 | 251 |
| 43 | Cardiac-generated prostanoids mediate cardiac myocyte apoptosis after myocardial ischaemia. Cardiovascular Research, 2012, 95, 336-345. | 1.8 | 26 |
| 44 | Biologically active ester derivatives as potent inhibitors of the soluble epoxide hydrolase. Bioorganic and Medicinal Chemistry Letters, 2012, 22, 5889-5892. | 1.0 | 10 |
| 45 | Development of pyrethroid-like fluorescent substrates for glutathione S-transferase. Analytical Biochemistry, 2012, 431, 77-83. | 1.1 | 8 |
| 46 | Soluble Epoxide Hydrolase Inhibitor Attenuates The Ovalbumin-Induced Murine Asthmatic Symptoms. , 2012, , . | | 0 |
| 47 | Substituted phenyl groups improve the pharmacokinetic profile of ureaâ€based soluble epoxide hydrolase inhibitors. FASEB Journal, 2012, 26, 849.1. | 0.2 | 1 |
| 48 | The anti-inflammatory effects of soluble epoxide hydrolase inhibitors are independent of leukocyte recruitment. Biochemical and Biophysical Research Communications, 2011, 410, 494-500. | 1.0 | 24 |
| 49 | Soluble Epoxide Hydrolase Inhibitors and Heart Failure. Cardiovascular Therapeutics, 2011, 29, 99-111. | 1.1 | 63 |
| 50 | Juvenile Hormone (JH) Esterase of the Mosquito Culex quinquefasciatus Is Not a Target of the JH Analog Insecticide Methoprene. PLoS ONE, 2011, 6, e28392. | 1.1 | 26 |
| 51 | Use of Metabolomic Profiling in the Study of Arachidonic Acid Metabolism in Cardiovascular Disease. Congestive Heart Failure, 2011, 17, 42-46. | 2.0 | 48 |
| 52 | Inhibition of soluble epoxide hydrolase contributes to the anti-inflammatory effect of antimicrobial triclocarban in a murine model. Toxicology and Applied Pharmacology, 2011, 255, 200-206. | 1.3 | 21 |
| 53 | Synthesis and Structureâ^'Activity Relationship Studies of Urea-Containing Pyrazoles as Dual Inhibitors of Cyclooxygenase-2 and Soluble Epoxide Hydrolase. Journal of Medicinal Chemistry, 2011, 54, 3037-3050. | 2.9 | 148 |
| 54 | Soluble Epoxide Hydrolase Is A Novel Therapeutic Target In Asthma By Modulating The Inflammatory Response. , 2010, , . | | 0 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Fumigaclavine C improves concanavalin A-induced liver injury in mice mainly via inhibiting TNF-α production and lymphocyte adhesion to extracellular matricesâ€. Journal of Pharmacy and Pharmacology, 2010, 56, 775-782. | 1.2 | 41 |
| 56 | Inhibition of soluble epoxide hydrolase enhances the anti-inflammatory effects of aspirin and 5-lipoxygenase activation protein inhibitor in a murine model. Biochemical Pharmacology, 2010, 79, 880-887. | 2.0 | 115 |
| 57 | Metabolic profiling of murine plasma reveals an unexpected biomarker in rofecoxib-mediated cardiovascular events. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 17017-17022. | 3.3 | 116 |
| 58 | 1-Aryl-3-(1-acylpiperidin-4-yl)urea Inhibitors of Human and Murine Soluble Epoxide Hydrolase: Structureâ^Activity Relationships, Pharmacokinetics, and Reduction of Inflammatory Pain. Journal of Medicinal Chemistry, 2010, 53, 7067-7075. | 2.9 | 148 |
| 59 | Incorporation of Piperazino Functionality into 1,3-Disubstituted Urea as the Tertiary Pharmacophore Affording Potent Inhibitors of Soluble Epoxide Hydrolase with Improved Pharmacokinetic Properties. Journal of Medicinal Chemistry, 2010, 53, 8376-8386. | 2.9 | 27 |
| 60 | Sorafenib has soluble epoxide hydrolase inhibitory activity, which contributes to its effect profile <i>in vivo</i> . Molecular Cancer Therapeutics, 2009, 8, 2193-2203. | 1.9 | 72 |
| 61 | Biotransformation of soybean isoflavones by a marine Streptomyces sp. 060524 and cytotoxicity of the products. World Journal of Microbiology and Biotechnology, 2009, 25, 115-121. | 1.7 | 25 |
| 62 | Pharmacokinetic optimization of four soluble epoxide hydrolase inhibitors for use in a murine model of inflammation. British Journal of Pharmacology, 2009, 156, 284-296. | 2.7 | 87 |
| 63 | Beneficial effects of soluble epoxide hydrolase inhibitors in myocardial infarction model: Insight gained using metabolomic approaches. Journal of Molecular and Cellular Cardiology, 2009, 47, 835-845. | 0.9 | 81 |
| 64 | Synthesis and biological evaluation of soluble epoxide hydrolase (sEH) inhibitors: tâ€AUCB and its derivatives. FASEB Journal, 2008, 22, 479.11. | 0.2 | 0 |
| 65 | Pharmacokinetic studies of four novel soluble epoxide hydrolase (sEH) inhibitors and antiâ€inflammatory efficacy of the most promising one tâ€AUCB. FASEB Journal, 2008, 22, 479.24. | 0.2 | 0 |
| 66 | Orally Bioavailable Potent Soluble Epoxide Hydrolase Inhibitors. Journal of Medicinal Chemistry, 2007, 50, 3825-3840. | 2.9 | 221 |
| 67 | Antifungal and new metabolites of Myrothecium sp. Z16, a fungus associated with white croaker Argyrosomus argentatus. Journal of Applied Microbiology, 2006, 100, 195-202. | 1.4 | 50 |
| 68 | Chaetominine, a Cytotoxic Alkaloid Produced by EndophyticChaetomiumsp. IFB-E015. Organic Letters, 2006, 8, 5709-5712. | 2.4 | 181 |
| 69 | New Resveratrol Oligomers from the Stem Bark ofHopea hainanensis. Helvetica Chimica Acta, 2005, 88, 2910-2917. | 1.0 | 25 |
| 70 | Anti-Helicobacter pylori substances from endophytic fungal cultures. World Journal of Microbiology and Biotechnology, 2005, 21, 553-558. | 1.7 | 48 |
| 71 | Structural Revision of Aspernigrin A, Reisolated fromCladosporiumherbarumIFB-E002. Journal of Natural Products, 2005, 68, 1106-1108. | 1.5 | 59 |
| 72 | In vitro anti-Helicobacter pylori action of 30 Chinese herbal medicines used to treat ulcer diseases. Journal of Ethnopharmacology, 2005, 98, 329-333. | 2.0 | 198 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Anti-Helicobacter pylori metabolites from Rhizoctonia sp. Cy064, an endophytic fungus in Cynodon dactylon. Fìtoterapìâ, 2004, 75, 451-456. | 1.1 | 66 |
| 74 | Aspergillus fumigatus CY018, an endophytic fungus in Cynodon dactylon as a versatile producer of new and bioactive metabolites. Journal of Biotechnology, 2004, 114, 279-287. | 1.9 | 132 |
| 75 | Leptosphaeric Acid, a Metabolite with a Novel Carbon Skeleton from Leptosphaeria sp. IV403, an Endophytic Fungus in Artemisia annua. Helvetica Chimica Acta, 2003, 86, 657-660. | 1.0 | 19 |
| 76 | Neoverataline A and B, two antifungal alkaloids with a novel carbon skeleton from Veratrum taliense. Tetrahedron, 2003, 59, 5743-5747. | 1.0 | 42 |
| 77 | Pinelloside, an antimicrobial cerebroside from Pinellia ternata. Phytochemistry, 2003, 64, 903-906. | 1.4 | 59 |
| 78 | Title is missing!. Helvetica Chimica Acta, 2002, 85, 2664-2667. | 1.0 | 36 |