

# Sarah J Charman

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3846397/publications.pdf>

Version: 2024-02-01

19  
papers

456  
citations

949033

11  
h-index

1113639

15  
g-index

20  
all docs

20  
docs citations

20  
times ranked

1040  
citing authors

#	ARTICLE	IF	CITATIONS
1	Physical Activity, Inactivity and Sleep in Patients with Significant Non-Alcoholic Fatty Liver Disease. <i>American Journal of the Medical Sciences</i> , 2022, 363, 80-83.	0.4	0
2	Movement as Medicine for Cardiovascular Disease Prevention: Pilot Feasibility Study of a Physical Activity Promotion Intervention for At-Risk Patients in Primary Care. <i>JMIR Cardio</i> , 2022, 6, e29035.	0.7	0
3	Feasibility of the cardiac output response to stress test in suspected heart failure patients. <i>Family Practice</i> , 2022, , .	0.8	1
4	Insights into heart failure hospitalizations, management, and services during and beyond COVID-19. <i>ESC Heart Failure</i> , 2021, 8, 175-182.	1.4	22
5	Overcoming barriers to engagement and adherence to a home-based physical activity intervention for patients with heart failure: a qualitative focus group study. <i>BMJ Open</i> , 2020, 10, e036382.	0.8	22
6	What are the Physiological Benefits of Increased Daily Number of Steps in Middle-Aged Women?. <i>American Journal of the Medical Sciences</i> , 2020, 360, 591-595.	0.4	0
7	Metabolic effects of bezafibrate in mitochondrial disease. <i>EMBO Molecular Medicine</i> , 2020, 12, e11589.	3.3	45
8	Analyzing walking speeds with ankle and wrist worn accelerometers in a cohort with myotonic dystrophy. <i>Disability and Rehabilitation</i> , 2019, 41, 2972-2978.	0.9	13
9	MSH3 modifies somatic instability and disease severity in Huntington's and myotonic dystrophy type 1. <i>Brain</i> , 2019, 142, 1876-1886.	3.7	114
10	Opportunities and challenges of a novel cardiac output response to stress (CORS) test to enhance diagnosis of heart failure in primary care: qualitative study. <i>BMJ Open</i> , 2019, 9, e028122.	0.8	3
11	Cardiac function is not associated with glucose control in older women. <i>Experimental Gerontology</i> , 2019, 116, 31-36.	1.2	0
12	Acceptability, Feasibility and Preliminary Evaluation of a Novel, Personalised, Home-Based Physical Activity Intervention for Chronic Heart Failure (Active-at-Home-HF): a Pilot Study. <i>Sports Medicine - Open</i> , 2019, 5, 45.	1.3	11
13	Physical education contributes to total physical activity levels and predominantly in higher intensity physical activity categories. <i>European Physical Education Review</i> , 2018, 24, 152-164.	1.2	22
14	A novel cardiac output response to stress test developed to improve diagnosis and monitoring of heart failure in primary care. <i>ESC Heart Failure</i> , 2018, 5, 703-712.	1.4	11
15	Cognitive behavioural therapy with optional graded exercise therapy in patients with severe fatigue with myotonic dystrophy type 1: a multicentre, single-blind, randomised trial. <i>Lancet Neurology</i> , The, 2018, 17, 671-680.	4.9	95
16	Associations between prolonged sedentary time and breaks in sedentary time with cardiometabolic risk in 10-14-year-old children: The HAPPY study. <i>Journal of Sports Sciences</i> , 2017, 35, 2164-2171.	1.0	36
17	Measuring Habitual Physical Activity in Neuromuscular Disorders: A Systematic Review. <i>Journal of Neuromuscular Diseases</i> , 2017, 4, 25-52.	1.1	28
18	The effect of percutaneous coronary intervention on habitual physical activity in older patients. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 248.	0.7	14

#	ARTICLE	IF	CITATIONS
19	Systematic development of a theory-informed multifaceted behavioural intervention to increase physical activity of adults with type 2 diabetes in routine primary care: Movement as Medicine for Type 2 Diabetes. <i>Implementation Science</i> , 2015, 11, 99.	2.5	19