

# Christine Masson

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/384904/christine-masson-publications-by-citations.pdf>

**Version:** 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

15  
papers

186  
citations

8  
h-index

13  
g-index

15  
ext. papers

207  
ext. citations

3.9  
avg, IF

1.74  
L-index

#	Paper	IF	Citations
15	Gamma-glutamyltranspeptidase-dependent glutathione catabolism results in activation of NF-kB. <i>Biochemical and Biophysical Research Communications</i> , <b>2000</b> , 276, 1062-7	3.4	34
14	Heritability for plasma VEGF concentration in the Stanislas family study. <i>Annals of Human Genetics</i> , <b>2007</b> , 71, 54-63	2.2	31
13	Effect of HMGCoA reductase inhibitors on cytochrome P450 expression in endothelial cell line. <i>Journal of Cardiovascular Pharmacology</i> , <b>2007</b> , 49, 306-15	3.1	20
12	Angiogenesis related genes NOS3, CD14, MMP3 and IL4R are associated to VEGF gene expression and circulating levels in healthy adults. <i>BMC Medical Genetics</i> , <b>2015</b> , 16, 90	2.1	19
11	Klotho KL-VS genotype is involved in blood pressure regulation. <i>Clinica Chimica Acta</i> , <b>2011</b> , 412, 1773-7	6.2	17
10	Human formyl peptide receptor 1 C32T SNP interacts with age and is associated with blood pressure levels. <i>Clinica Chimica Acta</i> , <b>2012</b> , 413, 34-8	6.2	14
9	Influence of pre-analytical variables on VEGF gene expression and circulating protein concentrations. <i>Biopreservation and Biobanking</i> , <b>2012</b> , 10, 454-61	2.1	14
8	Effect of acute and chronic psychostimulant drugs on redox status, AP-1 activation and pro-enkephalin mRNA in the human astrocyte-like U373 MG cells. <i>Neuropharmacology</i> , <b>2005</b> , 48, 673-84	5.5	10
7	Association between EGF and lipid concentrations: a benefit role in the atherosclerotic process?. <i>Clinica Chimica Acta</i> , <b>2009</b> , 402, 196-8	6.2	8
6	Functional epistatic interaction between rs6046G>A in F7 and rs5355C>T in SELE modifies systolic blood pressure levels. <i>PLoS ONE</i> , <b>2012</b> , 7, e40777	3.7	7
5	Genetic profiling of human cell lines used as in vitro model to study cardiovascular pathophysiology and pharmacotoxicology. <i>Cell Biology and Toxicology</i> , <b>2009</b> , 25, 561-71	7.4	5
4	Epigenome-wide association study in healthy individuals identifies significant associations with DNA methylation and PBMC extract VEGF-A concentration. <i>Clinical Epigenetics</i> , <b>2020</b> , 12, 79	7.7	3
3	Peripheral blood mononuclear cells extracts VEGF protein levels and VEGF mRNA: Associations with inflammatory molecules in a healthy population. <i>PLoS ONE</i> , <b>2019</b> , 14, e0220902	3.7	2
2	Lack of association between EGF 61A>G polymorphism and plasma EGF concentration in the STANISLAS family study. <i>Journal of Investigative Dermatology</i> , <b>2007</b> , 127, 969-70	4.3	1
1	A genetic determinant of VEGF-A levels is associated with telomere attrition. <i>Aging</i> , <b>2021</b> , 13, 23517-23526	3.6	1