

Kayhan Erturk

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/385048/publications.pdf>

Version: 2024-02-01

67
papers

455
citations

759233

12
h-index

839539

18
g-index

67
all docs

67
docs citations

67
times ranked

823
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Recurrence behavior in early-stage cutaneous melanoma: pattern, timing, survival, and influencing factors. <i>Melanoma Research</i> , 2017, 27, 134-139. | 1.2 | 36 |
| 2 | Tumor Infiltrating Lymphocytes (TILs) May be Only an Independent Predictor of Nodal Involvement but not for Recurrence and Survival in Cutaneous Melanoma Patients. <i>Cancer Investigation</i> , 2017, 35, 501-505. | 1.3 | 31 |
| 3 | Comparison of risk factors for cardiovascular disease in hemodialysis and peritoneal dialysis patients. <i>Clinics</i> , 2015, 70, 601-605. | 1.5 | 27 |
| 4 | Presence of histological regression as a prognostic factor in cutaneous melanoma patients. <i>Melanoma Research</i> , 2016, 26, 492-496. | 1.2 | 23 |
| 5 | Serum IGF-1 and IGFBP-3 levels as clinical markers for patients with lung cancer. <i>Biomedical Reports</i> , 2016, 4, 609-614. | 2.0 | 21 |
| 6 | Scalp melanoma is associated with high mitotic rate and is a poor prognostic factor for recurrence and outcome. <i>Melanoma Research</i> , 2017, 27, 387-390. | 1.2 | 21 |
| 7 | Clinical Significance of Circulating Serum Cellular Heat Shock Protein 90 (HSP90) Level in Patients with Cutaneous Malignant Melanoma. <i>Asian Pacific Journal of Cancer Prevention</i> , 2017, 18, 599-601. | 1.2 | 16 |
| 8 | Clinical significance of serum interleukin-29, interleukin-32, and tumor necrosis factor alpha levels in patients with gastric cancer. <i>Tumor Biology</i> , 2016, 37, 405-412. | 1.8 | 15 |
| 9 | Acral Lentiginous Melanoma Is Associated with Certain Poor Prognostic Histopathological Factors but May Not be Correlated with Nodal Involvement, Recurrence, and a Worse Survival. <i>Pathobiology</i> , 2018, 85, 227-231. | 3.8 | 15 |
| 10 | Serum nectin-2 and nectin-4 are diagnostic in lung cancer: which is superior?. <i>Wiener Klinische Wochenschrift</i> , 2019, 131, 419-426. | 1.9 | 15 |
| 11 | Serum activated leukocyte cell adhesion molecule and intercellular adhesion molecule-1 in patients with gastric cancer: Can they be used as biomarkers?. <i>Biomedicine and Pharmacotherapy</i> , 2016, 77, 86-91. | 5.6 | 14 |
| 12 | Elevated erythrocyte sedimentation rate is associated with metastatic disease and worse survival in patients with cutaneous malignant melanoma. <i>Molecular and Clinical Oncology</i> , 2017, 7, 1142-1146. | 1.0 | 14 |
| 13 | BRAF V600E mutation as a prognostic factor in cutaneous melanoma patients. <i>Dermatologic Therapy</i> , 2020, 33, e13270. | 1.7 | 14 |
| 14 | The efficacy of Pistacia Terebinthus soap in the treatment of cetuximab-induced skin toxicity. <i>Investigational New Drugs</i> , 2014, 32, 1295-1300. | 2.6 | 12 |
| 15 | Patient age and cutaneous malignant melanoma: Elderly patients are likely to have more aggressive histological features and poorer survival. <i>Molecular and Clinical Oncology</i> , 2017, 7, 1083-1088. | 1.0 | 12 |
| 16 | Clinical Significance of Serum Ykl-40 (Chitinase-3-Like-1 Protein) as a Biomarker in Melanoma: an Analysis of 112 Turkish Patients. <i>Asian Pacific Journal of Cancer Prevention</i> , 2017, 18, 1383-1387. | 1.2 | 10 |
| 17 | Histological lymphovascular invasion is associated with nodal involvement, recurrence, and survival in patients with cutaneous malignant melanoma. <i>International Journal of Dermatology</i> , 2017, 56, 166-170. | 1.0 | 9 |
| 18 | Nivolumab for metastatic uveal melanoma: a multicenter, retrospective study. <i>Melanoma Research</i> , 2021, 31, 449-455. | 1.2 | 9 |

| # | ARTICLE | IF | CITATIONS |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Primary tumour ulceration in cutaneous melanoma: its role on TNM stages.. Japanese Journal of Clinical Oncology, 2021, 51, 192-198. | 1.3 | 9 |
| 20 | Clinical Significance of Serum Galectin-3 Levels in Gastric Cancer Patients. Journal of Gastrointestinal Cancer, 2016, 47, 182-186. | 1.3 | 8 |
| 21 | Clinical significance of serum leptin level in patients with gastric cancer. European Cytokine Network, 2018, 29, 52-58. | 2.0 | 8 |
| 22 | De Novo and Nevus-Associated Melanomas: Different Histopathologic Characteristics but Similar Survival Rates. Pathology and Oncology Research, 2020, 26, 2483-2487. | 1.9 | 7 |
| 23 | Assessment of Anxiety and Depression Status in Turkish Cutaneous Melanoma Patients. Asian Pacific Journal of Cancer Prevention, 2017, 18, 369-373. | 1.2 | 7 |
| 24 | Early and late relapses of cutaneous melanoma patients. Postgraduate Medicine, 2019, 131, 207-211. | 2.0 | 6 |
| 25 | Limb melanomas: acral melanomas have worse survival. Journal of Dermatological Treatment, 2022, 33, 1630-1637. | 2.2 | 6 |
| 26 | Different mitotic rates are associated with different prognostic factors, relapses, and survival rates in melanoma. International Journal of Dermatology, 2022, 61, 472-479. | 1.0 | 6 |
| 27 | Clinical significance of serum claudin-1 levels in melanoma patients. Melanoma Research, 2016, 26, 377-381. | 1.2 | 5 |
| 28 | Cutaneous Melanoma in Association With Ichthyosis Vulgaris. Journal of Pediatric Hematology/Oncology, 2017, 39, 630-630. | 0.6 | 5 |
| 29 | Effect of biology on the outcome of female melanoma patients. Molecular and Clinical Oncology, 2017, 7, 1093-1100. | 1.0 | 5 |
| 30 | Anemia in Cutaneous Malignant Melanoma: Low Blood Hemoglobin Level is Associated with Nodal Involvement, Metastatic Disease, and Worse Survival. Nutrition and Cancer, 2018, 70, 236-240. | 2.0 | 5 |
| 31 | Coexistence of regression and tumor infiltrating lymphocytes is associated with more favorable survival in melanoma. Journal of Cancer Research and Clinical Oncology, 2021, 147, 2721-2729. | 2.5 | 5 |
| 32 | Clinical significance of serum protease activated receptor1 levels in patients with lung cancer. European Review for Medical and Pharmacological Sciences, 2016, 20, 243-9. | 0.7 | 5 |
| 33 | Cheek Cutaneous Melanomas. Annals of Plastic Surgery, 2019, 82, 407-410. | 0.9 | 4 |
| 34 | Palpebral cutaneous melanomas: a review of 17 cases from a tertiary center. International Journal of Dermatology, 2019, 58, 75-79. | 1.0 | 4 |
| 35 | Awareness on malignant melanoma and its prevention measures among Turkish cutaneous malignant melanoma patients: A tertiary cancer center experience. Dermatologic Therapy, 2020, 33, e14425. | 1.7 | 4 |
| 36 | Seasons influence diagnosis and outcome of cutaneous melanoma. Dermatologic Therapy, 2020, 33, e13625. | 1.7 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 37 | Clinical significance of serum Protease-Activated Receptor-1 (PAR-1) levels in patients with cutaneous melanoma. <i>BBA Clinical</i> , 2016, 5, 166-169. | 4.1 | 3 |
| 38 | Clinical significance of serum laminin levels in patients with lung cancer. <i>Biomedical Reports</i> , 2016, 4, 485-488. | 2.0 | 3 |
| 39 | Significance of serum neural precursor cell-expressed developmentally downregulated protein 9 in melanoma. <i>Molecular and Clinical Oncology</i> , 2017, 8, 204-208. | 1.0 | 3 |
| 40 | Digital clubbing as a first clinical presentation of pulmonary metastases in cutaneous melanoma. <i>Postgraduate Medicine</i> , 2018, 130, 278-279. | 2.0 | 3 |
| 41 | BRAF mutation status might contribute an effect on both disease-free and overall survival in stage III cutaneous melanomas treated with intermediate dose interferon-alpha. <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 84, 521-526. | 2.3 | 3 |
| 42 | Lymph node ratio has impact on relapse and outcome in patients with stage III melanoma. <i>International Journal of Clinical Oncology</i> , 2019, 24, 721-726. | 2.2 | 3 |
| 43 | Google searching as an indicator of population's interest in melanoma: A comparative study in Google Trends. <i>Dermatologic Therapy</i> , 2020, 33, e14421. | 1.7 | 3 |
| 44 | Inevitable hemodialysis for treating resistant hypertension in a patient with Leriche syndrome. <i>Clinics</i> , 2012, 67, 1483-1486. | 1.5 | 3 |
| 45 | Widespread finger skin metastases of melanoma. <i>Clinical Case Reports (discontinued)</i> , 2018, 6, 448-449. | 0.5 | 2 |
| 46 | Spitzoid cutaneous melanoma is associated with favorable clinicopathological factors and outcome. <i>Journal of Cosmetic Dermatology</i> , 2019, 18, 1841-1845. | 1.6 | 2 |
| 47 | Single-agent temozolomide may be an effective option for late adjuvant therapy in patients with melanoma. <i>Journal of Oncology Pharmacy Practice</i> , 2021, 27, 40-45. | 0.9 | 2 |
| 48 | Mitotic rate in node-positive stage III melanoma: it might be as important a prognostic factor as node number. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 873-878. | 1.3 | 2 |
| 49 | Digit melanomas are associated with poor prognostic factors and unfavorable survivals. <i>Journal of Cosmetic Dermatology</i> , 2022, 21, 2120-2129. | 1.6 | 2 |
| 50 | Is it solitary plasmacytoma or nonsecretory myeloma? A must-be-solved dilemma?. <i>Biomedicine and Pharmacotherapy</i> , 2016, 77, 27-29. | 5.6 | 1 |
| 51 | The course of stage III melanoma in accordance with the severity of node involvement. <i>Current Medical Research and Opinion</i> , 2019, 35, 1819-1824. | 1.9 | 1 |
| 52 | Number of Excised Lymph Nodes Has No Impact on Relapse and Survival in Patients With Stage III Melanoma. <i>Annals of Plastic Surgery</i> , 2019, 83, 455-458. | 0.9 | 1 |
| 53 | Trends in the characteristics of skin melanoma in accordance with time intervals: A single Turkish tertiary referral center experience. <i>Journal of Cancer Research and Therapeutics</i> , 2021, 17, 1119. | 0.9 | 1 |
| 54 | Cutaneous melanoma in vicenarians: Patients in their twenties and older patients show similar clinical behaviors and survival rates. <i>Journal of Cosmetic Dermatology</i> , 2020, 19, 2692-2696. | 1.6 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 55 | Serum 25-Hydroxyvitamin D Level Is Not Associated with Duration and Activity of Disease in Melanoma Patients. <i>Nutrition and Cancer</i> , 2021, 73, 1126-1129. | 2.0 | 0 |
| 56 | Serum folate and vitamin B12 levels in cutaneous melanoma. <i>Journal of Cosmetic Dermatology</i> , 2021, 20, 3007-3010. | 1.6 | 0 |
| 57 | Complementary and alternative medicine (CAM) in Turkish cutaneous melanoma patients: A prospective study from tertiary cancer center. <i>Journal of Oncology Pharmacy Practice</i> , 2022, 28, 282-286. | 0.9 | 0 |
| 58 | Auricular and periauricular melanomas have similar clinicopathologic factors and survival rates. <i>Journal of Cosmetic Dermatology</i> , 2021, , . | 1.6 | 0 |
| 59 | Unknown Primary Metastatic Melanoma Presented with Extensive Subcutaneous Masses and Lymph Node Enlargements. <i>Indian Journal of Surgery</i> , 0, , 1. | 0.3 | 0 |
| 60 | Larger Tumors Are Associated with Poorer Prognostic Factors in Cutaneous Melanoma. <i>Indian Journal of Surgery</i> , 0, , 1. | 0.3 | 0 |
| 61 | The role of adhesion molecules in hepatocellular carcinoma.. <i>Journal of Clinical Oncology</i> , 2015, 33, e15144-e15144. | 1.6 | 0 |
| 62 | Do serum nectin-2 levels have a prognostic effect in patients with colorectal cancer?. <i>Journal of Clinical Oncology</i> , 2015, 33, e14531-e14531. | 1.6 | 0 |
| 63 | Clinical significance of serum claudin-7 levels in patients with colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, e14533-e14533. | 1.6 | 0 |
| 64 | Diagnostic and prognostic roles of serum interleukin-32 (IL-32) levels in patients with pancreatic adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2016, 34, e15680-e15680. | 1.6 | 0 |
| 65 | Seasons Influence Diagnosis of Breast Cancer in Turkey. <i>Indian Journal of Surgery</i> , 0, , 1. | 0.3 | 0 |
| 66 | Trunk melanomas: no survival differences between lesion sites. <i>Postgraduate Medicine</i> , 2022, , . | 2.0 | 0 |
| 67 | Using Google as a Source of Information About Breast Cancer. <i>Indian Journal of Surgery</i> , 0, , . | 0.3 | 0 |