

Kayhan Erturk

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/385048/publications.pdf>

Version: 2024-02-01

67
papers

455
citations

759233

12
h-index

839539

18
g-index

67
all docs

67
docs citations

67
times ranked

823
citing authors

#	ARTICLE	IF	CITATIONS
1	Complementary and alternative medicine (CAM) in Turkish cutaneous melanoma patients: A prospective study from tertiary cancer center. <i>Journal of Oncology Pharmacy Practice</i> , 2022, 28, 282-286.	0.9	0
2	Limb melanomas: acral melanomas have worse survival. <i>Journal of Dermatological Treatment</i> , 2022, 33, 1630-1637.	2.2	6
3	Digit melanomas are associated with poor prognostic factors and unfavorable survivals. <i>Journal of Cosmetic Dermatology</i> , 2022, 21, 2120-2129.	1.6	2
4	Different mitotic rates are associated with different prognostic factors, relapses, and survival rates in melanoma. <i>International Journal of Dermatology</i> , 2022, 61, 472-479.	1.0	6
5	Trunk melanomas: no survival differences between lesion sites. <i>Postgraduate Medicine</i> , 2022, , .	2.0	0
6	Single-agent temozolomide may be an effective option for late adjuvant therapy in patients with melanoma. <i>Journal of Oncology Pharmacy Practice</i> , 2021, 27, 40-45.	0.9	2
7	Serum 25-Hydroxyvitamin D Level Is Not Associated with Duration and Activity of Disease in Melanoma Patients. <i>Nutrition and Cancer</i> , 2021, 73, 1126-1129.	2.0	0
8	Serum folate and vitamin B12 levels in cutaneous melanoma. <i>Journal of Cosmetic Dermatology</i> , 2021, 20, 3007-3010.	1.6	0
9	Trends in the characteristics of skin melanoma in accordance with time intervals: A single Turkish tertiary referral center experience. <i>Journal of Cancer Research and Therapeutics</i> , 2021, 17, 1119.	0.9	1
10	Coexistence of regression and tumor infiltrating lymphocytes is associated with more favorable survival in melanoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 2721-2729.	2.5	5
11	Mitotic rate in node-positive stage III melanoma: it might be as important a prognostic factor as node number. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 873-878.	1.3	2
12	Nivolumab for metastatic uveal melanoma: a multicenter, retrospective study. <i>Melanoma Research</i> , 2021, 31, 449-455.	1.2	9
13	Auricular and periauricular melanomas have similar clinicopathologic factors and survival rates. <i>Journal of Cosmetic Dermatology</i> , 2021, , .	1.6	0
14	Primary tumour ulceration in cutaneous melanoma: its role on TNM stages.. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 192-198.	1.3	9
15	Google searching as an indicator of population's interest in melanoma: A comparative study in Google Trends. <i>Dermatologic Therapy</i> , 2020, 33, e14421.	1.7	3
16	Awareness on malignant melanoma and its prevention measures among Turkish cutaneous malignant melanoma patients: A tertiary cancer center experience. <i>Dermatologic Therapy</i> , 2020, 33, e14425.	1.7	4
17	Seasons influence diagnosis and outcome of cutaneous melanoma. <i>Dermatologic Therapy</i> , 2020, 33, e13625.	1.7	4
18	De Novo and Nevus-Associated Melanomas: Different Histopathologic Characteristics but Similar Survival Rates. <i>Pathology and Oncology Research</i> , 2020, 26, 2483-2487.	1.9	7

#	ARTICLE	IF	CITATIONS
19	BRAF V600E mutation as a prognostic factor in cutaneous melanoma patients. <i>Dermatologic Therapy</i> , 2020, 33, e13270.	1.7	14
20	Cutaneous melanoma in vicenarians: Patients in their twenties and older patients show similar clinical behaviors and survival rates. <i>Journal of Cosmetic Dermatology</i> , 2020, 19, 2692-2696.	1.6	0
21	Serum nectin-2 and nectin-4 are diagnostic in lung cancer: which is superior?. <i>Wiener Klinische Wochenschrift</i> , 2019, 131, 419-426.	1.9	15
22	The course of stage III melanoma in accordance with the severity of node involvement. <i>Current Medical Research and Opinion</i> , 2019, 35, 1819-1824.	1.9	1
23	BRAF mutation status might contribute an effect on both disease-free and overall survival in stage III cutaneous melanomas treated with intermediate dose interferon-alpha. <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 84, 521-526.	2.3	3
24	Spitzoid cutaneous melanoma is associated with favorable clinicopathological factors and outcome. <i>Journal of Cosmetic Dermatology</i> , 2019, 18, 1841-1845.	1.6	2
25	Lymph node ratio has impact on relapse and outcome in patients with stage III melanoma. <i>International Journal of Clinical Oncology</i> , 2019, 24, 721-726.	2.2	3
26	Cheek Cutaneous Melanomas. <i>Annals of Plastic Surgery</i> , 2019, 82, 407-410.	0.9	4
27	Number of Excised Lymph Nodes Has No Impact on Relapse and Survival in Patients With Stage III Melanoma. <i>Annals of Plastic Surgery</i> , 2019, 83, 455-458.	0.9	1
28	Palpebral cutaneous melanomas: a review of 17 cases from a tertiary center. <i>International Journal of Dermatology</i> , 2019, 58, 75-79.	1.0	4
29	Early and late relapses of cutaneous melanoma patients. <i>Postgraduate Medicine</i> , 2019, 131, 207-211.	2.0	6
30	Widespread finger skin metastases of melanoma. <i>Clinical Case Reports (discontinued)</i> , 2018, 6, 448-449.	0.5	2
31	Anemia in Cutaneous Malignant Melanoma: Low Blood Hemoglobin Level is Associated with Nodal Involvement, Metastatic Disease, and Worse Survival. <i>Nutrition and Cancer</i> , 2018, 70, 236-240.	2.0	5
32	Digital clubbing as a first clinical presentation of pulmonary metastases in cutaneous melanoma. <i>Postgraduate Medicine</i> , 2018, 130, 278-279.	2.0	3
33	Clinical significance of serum leptin level in patients with gastric cancer. <i>European Cytokine Network</i> , 2018, 29, 52-58.	2.0	8
34	Acral Lentiginous Melanoma Is Associated with Certain Poor Prognostic Histopathological Factors but May Not be Correlated with Nodal Involvement, Recurrence, and a Worse Survival. <i>Pathobiology</i> , 2018, 85, 227-231.	3.8	15
35	Recurrence behavior in early-stage cutaneous melanoma: pattern, timing, survival, and influencing factors. <i>Melanoma Research</i> , 2017, 27, 134-139.	1.2	36
36	Scalp melanoma is associated with high mitotic rate and is a poor prognostic factor for recurrence and outcome. <i>Melanoma Research</i> , 2017, 27, 387-390.	1.2	21

#	ARTICLE	IF	CITATIONS
37	Tumor Infiltrating Lymphocytes (TILs) May be Only an Independent Predictor of Nodal Involvement but not for Recurrence and Survival in Cutaneous Melanoma Patients. <i>Cancer Investigation</i> , 2017, 35, 501-505.	1.3	31
38	Histological lymphovascular invasion is associated with nodal involvement, recurrence, and survival in patients with cutaneous malignant melanoma. <i>International Journal of Dermatology</i> , 2017, 56, 166-170.	1.0	9
39	Cutaneous Melanoma in Association With Ichthyosis Vulgaris. <i>Journal of Pediatric Hematology/Oncology</i> , 2017, 39, 630-630.	0.6	5
40	Elevated erythrocyte sedimentation rate is associated with metastatic disease and worse survival in patients with cutaneous malignant melanoma. <i>Molecular and Clinical Oncology</i> , 2017, 7, 1142-1146.	1.0	14
41	Effect of biology on the outcome of female melanoma patients. <i>Molecular and Clinical Oncology</i> , 2017, 7, 1093-1100.	1.0	5
42	Significance of serum neural precursor cell-expressed developmentally downregulated protein 9 in melanoma. <i>Molecular and Clinical Oncology</i> , 2017, 8, 204-208.	1.0	3
43	Patient age and cutaneous malignant melanoma: Elderly patients are likely to have more aggressive histological features and poorer survival. <i>Molecular and Clinical Oncology</i> , 2017, 7, 1083-1088.	1.0	12
44	Assessment of Anxiety and Depression Status in Turkish Cutaneous Melanoma Patients. <i>Asian Pacific Journal of Cancer Prevention</i> , 2017, 18, 369-373.	1.2	7
45	Clinical Significance of Circulating Serum Cellular Heat Shock Protein 90 (HSP90) Level in Patients with Cutaneous Malignant Melanoma. <i>Asian Pacific Journal of Cancer Prevention</i> , 2017, 18, 599-601.	1.2	16
46	Clinical Significance of Serum Ykl-40 (Chitinase-3-Like-1 Protein) as a Biomarker in Melanoma: an Analysis of 112 Turkish Patients. <i>Asian Pacific Journal of Cancer Prevention</i> , 2017, 18, 1383-1387.	1.2	10
47	Clinical significance of serum claudin-1 levels in melanoma patients. <i>Melanoma Research</i> , 2016, 26, 377-381.	1.2	5
48	Presence of histological regression as a prognostic factor in cutaneous melanoma patients. <i>Melanoma Research</i> , 2016, 26, 492-496.	1.2	23
49	Clinical significance of serum Protease-Activated Receptor-1 (PAR-1) levels in patients with cutaneous melanoma. <i>BBA Clinical</i> , 2016, 5, 166-169.	4.1	3
50	Clinical Significance of Serum Galectin-3 Levels in Gastric Cancer Patients. <i>Journal of Gastrointestinal Cancer</i> , 2016, 47, 182-186.	1.3	8
51	Clinical significance of serum laminin levels in patients with lung cancer. <i>Biomedical Reports</i> , 2016, 4, 485-488.	2.0	3
52	Is it solitary plasmacytoma or nonsecretory myeloma? A must-be-solved dilemma?. <i>Biomedicine and Pharmacotherapy</i> , 2016, 77, 27-29.	5.6	1
53	Serum IGF-1 and IGFBP-3 levels as clinical markers for patients with lung cancer. <i>Biomedical Reports</i> , 2016, 4, 609-614.	2.0	21
54	Serum activated leukocyte cell adhesion molecule and intercellular adhesion molecule-1 in patients with gastric cancer: Can they be used as biomarkers?. <i>Biomedicine and Pharmacotherapy</i> , 2016, 77, 86-91.	5.6	14

#	ARTICLE	IF	CITATIONS
55	Clinical significance of serum interleukin-29, interleukin-32, and tumor necrosis factor alpha levels in patients with gastric cancer. <i>Tumor Biology</i> , 2016, 37, 405-412.	1.8	15
56	Diagnostic and prognostic roles of serum interleukin-32 (IL-32) levels in patients with pancreatic adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2016, 34, e15680-e15680.	1.6	0
57	Clinical significance of serum protease activated receptor1 levels in patients with lung cancer. <i>European Review for Medical and Pharmacological Sciences</i> , 2016, 20, 243-9.	0.7	5
58	Comparison of risk factors for cardiovascular disease in hemodialysis and peritoneal dialysis patients. <i>Clinics</i> , 2015, 70, 601-605.	1.5	27
59	The role of adhesion molecules in hepatocellular carcinoma.. <i>Journal of Clinical Oncology</i> , 2015, 33, e15144-e15144.	1.6	0
60	Do serum nectin-2 levels have a prognostic effect in patients with colorectal cancer?. <i>Journal of Clinical Oncology</i> , 2015, 33, e14531-e14531.	1.6	0
61	Clinical significance of serum claudin-7 levels in patients with colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, e14533-e14533.	1.6	0
62	The efficacy of Pistacia Terebinthus soap in the treatment of cetuximab-induced skin toxicity. <i>Investigational New Drugs</i> , 2014, 32, 1295-1300.	2.6	12
63	Inevitable hemodialysis for treating resistant hypertension in a patient with Leriche syndrome. <i>Clinics</i> , 2012, 67, 1483-1486.	1.5	3
64	Unknown Primary Metastatic Melanoma Presented with Extensive Subcutaneous Masses and Lymph Node Enlargements. <i>Indian Journal of Surgery</i> , 0, , 1.	0.3	0
65	Larger Tumors Are Associated with Poorer Prognostic Factors in Cutaneous Melanoma. <i>Indian Journal of Surgery</i> , 0, , 1.	0.3	0
66	Seasons Influence Diagnosis of Breast Cancer in Turkey. <i>Indian Journal of Surgery</i> , 0, , 1.	0.3	0
67	Using Google as a Source of Information About Breast Cancer. <i>Indian Journal of Surgery</i> , 0, , .	0.3	0