Olivier Barbier

List of Publications by Citations

Source: https://exaly.com/author-pdf/3864203/olivier-barbier-publications-by-citations.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

108 papers

4,813 citations

41 h-index 66 g-index

111 ext. papers

5,313 ext. citations

5.5 avg, IF

5.18 L-index

#	Paper	IF	Citations
108	Peroxisome proliferator-activated receptors: regulation of transcriptional activities and roles in inflammation. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2003 , 85, 267-73	5.1	233
107	Farnesoid X receptor agonists suppress hepatic apolipoprotein CIII expression. <i>Gastroenterology</i> , 2003 , 125, 544-55	13.3	210
106	Glucose regulates the expression of the farnesoid X receptor in liver. <i>Diabetes</i> , 2004 , 53, 890-8	0.9	195
105	Inactivation of androgens by UDP-glucuronosyltransferase enzymes in humans. <i>Trends in Endocrinology and Metabolism</i> , 2003 , 14, 473-9	8.8	186
104	FXR induces the UGT2B4 enzyme in hepatocytes: a potential mechanism of negative feedback control of FXR activity. <i>Gastroenterology</i> , 2003 , 124, 1926-40	13.3	158
103	Acute antiinflammatory properties of statins involve peroxisome proliferator-activated receptor-alpha via inhibition of the protein kinase C signaling pathway. <i>Circulation Research</i> , 2006 , 98, 361-9	15.7	147
102	PPAR alpha inhibits vascular smooth muscle cell proliferation underlying intimal hyperplasia by inducing the tumor suppressor p16INK4a. <i>Journal of Clinical Investigation</i> , 2005 , 115, 3228-38	15.9	126
101	Long noncoding RNA MEG3 induces cholestatic liver injury by interaction with PTBP1 to facilitate shp mRNA decay. <i>Hepatology</i> , 2017 , 65, 604-615	11.2	118
100	Treatment with camu camu () prevents obesity by altering the gut microbiota and increasing energy expenditure in diet-induced obese mice. <i>Gut</i> , 2019 , 68, 453-464	19.2	117
99	Peroxisome proliferator-activated receptor alpha induces hepatic expression of the human bile acid glucuronidating UDP-glucuronosyltransferase 2B4 enzyme. <i>Journal of Biological Chemistry</i> , 2003 , 278, 32852-60	5.4	106
98	Tissue-specific, inducible, and hormonal control of the human UDP-glucuronosyltransferase-1 (UGT1) locus. <i>Journal of Biological Chemistry</i> , 2005 , 280, 37547-57	5.4	105
97	Peroxisome proliferator-activated receptor alpha (PPARalpha) turnover by the ubiquitin-proteasome system controls the ligand-induced expression level of its target genes. <i>Journal of Biological Chemistry</i> , 2002 , 277, 37254-9	5.4	104
96	The UDP-glucuronosyltransferase 1A9 enzyme is a peroxisome proliferator-activated receptor alpha and gamma target gene. <i>Journal of Biological Chemistry</i> , 2003 , 278, 13975-83	5.4	98
95	H19 promotes cholestatic liver fibrosis by preventing ZEB1-mediated inhibition of epithelial cell adhesion molecule. <i>Hepatology</i> , 2017 , 66, 1183-1196	11.2	97
94	Expression of the human UGT1 locus in transgenic mice by 4-chloro-6-(2,3-xylidino)-2-pyrimidinylthioacetic acid (WY-14643) and implications on drug metabolism through peroxisome proliferator-activated receptor alpha activation. <i>Drug Metabolism</i>	4	96
93	Human UDP-glucuronosyltransferase (UGT)1A3 enzyme conjugates chenodeoxycholic acid in the liver. <i>Hepatology</i> , 2006 , 44, 1158-70	11.2	94
92	Characterization of UDP-glucuronosyltransferases active on steroid hormones. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1999 , 69, 413-23	5.1	93

(2006-2012)

91	Metabolomic profiling of 17 bile acids in serum from patients with primary biliary cirrhosis and primary sclerosing cholangitis: a pilot study. <i>Digestive and Liver Disease</i> , 2012 , 44, 303-10	3.3	92	
90	UDP-glucuronosyltransferase 2B15 (UGT2B15) and UGT2B17 enzymes are major determinants of the androgen response in prostate cancer LNCaP cells. <i>Journal of Biological Chemistry</i> , 2007 , 282, 3346	6- 3 347	4 ⁸³	
89	Intestinal glucuronidation protects against chemotherapy-induced toxicity by irinotecan (CPT-11). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 19143-8	11.5	78	
88	Bcl2 is a critical regulator of bile acid homeostasis by dictating Shp and lncRNA H19 function. <i>Scientific Reports</i> , 2016 , 6, 20559	4.9	75	
87	Inactivation by UDP-glucuronosyltransferase enzymes: the end of androgen signaling. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2008 , 109, 247-53	5.1	73	
86	Prospective evaluation of ursodeoxycholic acid withdrawal in patients with primary sclerosing cholangitis. <i>Hepatology</i> , 2014 , 60, 931-40	11.2	72	
85	Profiling circulating and urinary bile acids in patients with biliary obstruction before and after biliary stenting. <i>PLoS ONE</i> , 2011 , 6, e22094	3.7	66	
84	Inactivation of androgens by UDP-glucuronosyltransferases in the human prostate. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2008 , 22, 259-70	6.5	59	
83	Effect of n-3 fatty acids on the expression of inflammatory genes in THP-1 macrophages. <i>Lipids in Health and Disease</i> , 2016 , 15, 69	4.4	58	
82	The liver X-receptor alpha controls hepatic expression of the human bile acid-glucuronidating UGT1A3 enzyme in human cells and transgenic mice. <i>Hepatology</i> , 2006 , 44, 368-78	11.2	58	
81	Arctic berry extracts target the gut-liver axis to alleviate metabolic endotoxaemia, insulin resistance and hepatic steatosis in diet-induced obese mice. <i>Diabetologia</i> , 2018 , 61, 919-931	10.3	56	
80	Expression of hepatic Fibroblast Growth Factor 19 is enhanced in Primary Biliary Cirrhosis and correlates with severity of the disease. <i>Scientific Reports</i> , 2015 , 5, 13462	4.9	56	
79	Hepatic expression of the UGT1A9 gene is governed by hepatocyte nuclear factor 4alpha. <i>Molecular Pharmacology</i> , 2005 , 67, 241-9	4.3	56	
78	Transcriptional regulation of human Rev-erbalpha gene expression by the orphan nuclear receptor retinoic acid-related orphan receptor alpha. <i>Journal of Biological Chemistry</i> , 2002 , 277, 49275-81	5.4	55	
77	Transcriptomic and metabolomic signatures of an n-3 polyunsaturated fatty acids supplementation in a normolipidemic/normocholesterolemic Caucasian population. <i>Journal of Nutritional Biochemistry</i> , 2013 , 24, 54-61	6.3	54	
76	Lipid-activated transcription factors control bile acid glucuronidation. <i>Molecular and Cellular Biochemistry</i> , 2009 , 326, 3-8	4.2	50	
<i>75</i>	Cellular specific expression of the androgen-conjugating enzymes UGT2B15 and UGT2B17 in the human prostate epithelium. <i>Endocrine Research</i> , 2004 , 30, 717-25	1.9	49	
74	Coordinate regulation of hepatic bile acid oxidation and conjugation by nuclear receptors. Molecular Pharmaceutics, 2006 , 3, 212-22	5.6	47	

73	Multiple roles for UDP-glucuronosyltransferase (UGT)2B15 and UGT2B17 enzymes in androgen metabolism and prostate cancer evolution. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2015 , 145, 187-92	5.1	44
72	Isoform-specific regulation of uridine diphosphate-glucuronosyltransferase 2B enzymes in the human prostate: differential consequences for androgen and bioactive lipid inactivation. <i>Endocrinology</i> , 2006 , 147, 5431-42	4.8	43
71	Inactivation of the pure antiestrogen fulvestrant and other synthetic estrogen molecules by UDP-glucuronosyltransferase 1A enzymes expressed in breast tissue. <i>Molecular Pharmacology</i> , 2006 , 69, 908-20	4.3	43
70	Cellular localization of uridine diphosphoglucuronosyltransferase 2B enzymes in the human prostate by in situ hybridization and immunohistochemistry. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2000 , 85, 4819-26	5.6	43
69	N-glycosylation and residue 96 are involved in the functional properties of UDP-glucuronosyltransferase enzymes. <i>Biochemistry</i> , 2000 , 39, 11540-52	3.2	42
68	Activators of the farnesoid X receptor negatively regulate androgen glucuronidation in human prostate cancer LNCAP cells. <i>Biochemical Journal</i> , 2008 , 410, 245-53	3.8	41
67	Genome-wide association study of the plasma triglyceride response to an n-3 polyunsaturated fatty acid supplementation. <i>Journal of Lipid Research</i> , 2014 , 55, 1245-53	6.3	38
66	Trans Fatty Acids Suppress TNF-Induced Inflammatory Gene Expression in Endothelial (HUVEC) and Hepatocellular Carcinoma (HepG2) Cells. <i>Lipids</i> , 2017 , 52, 315-325	1.6	36
65	Androgen glucuronidation: an unexpected target for androgen deprivation therapy, with prognosis and diagnostic implications. <i>Cancer Research</i> , 2013 , 73, 6963-71	10.1	35
64	Differential expression of the androgen-conjugating UGT2B15 and UGT2B17 enzymes in prostate tumor cells during cancer progression. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, E428-	-32 ⁶	35
63	Isolation and characterization of a simian UDP-glucuronosyltransferase UGT2B18 active on 3-hydroxyandrogens. <i>Journal of Molecular Biology</i> , 1998 , 275, 785-94	6.5	35
62	Transgenic B PUFA enrichment alters morphology and gene expression profile in adipose tissue of obese mice: Potential role for protectins. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 666-76	12.7	33
61	A pharmacogenomics study of the human estrogen glucuronosyltransferase UGT1A3. <i>Pharmacogenetics and Genomics</i> , 2007 , 17, 481-95	1.9	33
60	Serum Autotaxin is a Marker of the Severity of Liver Injury and Overall Survival in Patients with Cholestatic Liver Diseases. <i>Scientific Reports</i> , 2016 , 6, 30847	4.9	32
59	Differences in metabolomic and transcriptomic profiles between responders and non-responders to an n-3 polyunsaturated fatty acids (PUFAs) supplementation. <i>Genes and Nutrition</i> , 2013 , 8, 411-23	4.3	31
58	The cynomolgus monkey (Macaca fascicularis) is the best animal model for the study of steroid glucuronidation. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2003 , 85, 235-45	5.1	31
57	Genomic and non-genomic interactions of PPARalpha with xenobiotic-metabolizing enzymes. <i>Trends in Endocrinology and Metabolism</i> , 2004 , 15, 324-30	8.8	30
56	Distribution of uridine diphosphate-glucuronosyltransferase (UGT) expression and activity in cynomolgus monkey tissues: evidence for differential expression of steroid-conjugating UGT enzymes in steroid target tissues. <i>Endocrinology</i> , 2000 , 141, 2472-80	4.8	30

(2010-2003)

55	Human uridine diphosphate-glucuronosyltransferase UGT2B7 conjugates mineralocorticoid and glucocorticoid metabolites. <i>Endocrinology</i> , 2003 , 144, 2659-68	4.8	29	
54	Cloning and characterization of a simian UDP-glucuronosyltransferase enzyme UGT2B20, a novel C19 steroid-conjugating protein. <i>Biochemical Journal</i> , 1999 , 337, 567-574	3.8	29	
53	Crypt Organoid Culture as an in Vitro Model in Drug Metabolism and Cytotoxicity Studies. <i>Drug Metabolism and Disposition</i> , 2017 , 45, 748-754	4	28	•
52	Calcitrol (1alpha,25-dihydroxyvitamin D3) inhibits androgen glucuronidation in prostate cancer cells. <i>Molecular Cancer Therapeutics</i> , 2008 , 7, 380-90	6.1	28	
51	Association between Metabolite Profiles, Metabolic Syndrome and Obesity Status. <i>Nutrients</i> , 2016 , 8,	6.7	27	
50	UGT2B17 Expedites Progression of Castration-Resistant Prostate Cancers by Promoting Ligand-Independent AR Signaling. <i>Cancer Research</i> , 2016 , 76, 6701-6711	10.1	27	
49	Molecular cloning, expression and characterization of a monkey steroid UDP-glucuronosyltransferase, UGT2B19, that conjugates testosterone. <i>FEBS Journal</i> , 1999 , 260, 701-8		26	
48	TReP-132 is a novel progesterone receptor coactivator required for the inhibition of breast cancer cell growth and enhancement of differentiation by progesterone. <i>Molecular and Cellular Biology</i> , 2006 , 26, 7632-44	4.8	25	
47	The Human UDP-glucuronosyltransferase UGT2A1 and UGT2A2 enzymes are highly active in bile acid glucuronidation. <i>Drug Metabolism and Disposition</i> , 2013 , 41, 1616-20	4	24	
46	Enzymatic production of bile Acid glucuronides used as analytical standards for liquid chromatography-mass spectrometry analyses. <i>Molecular Pharmaceutics</i> , 2006 , 3, 293-302	5.6	23	
45	Role of glucuronidation for hepatic detoxification and urinary elimination of toxic bile acids during biliary obstruction. <i>PLoS ONE</i> , 2013 , 8, e80994	3.7	23	
44	The hepatokine Tsukushi is released in response to NAFLD and impacts cholesterol homeostasis. JCI Insight, 2019 , 4,	9.9	22	
43	Plasma estrone sulfate assay in men: Comparison of radioimmunoassay, mass spectrometry coupled to gas chromatography (GC-MS), and liquid chromatography-tandem mass spectrometry (LC-MS/MS). <i>Clinica Chimica Acta</i> , 2010 , 411, 1208-13	6.2	21	
42	TReP-132 controls cell proliferation by regulating the expression of the cyclin-dependent kinase inhibitors p21WAF1/Cip1 and p27Kip1. <i>Molecular and Cellular Biology</i> , 2005 , 25, 4335-48	4.8	21	
41	A Study of the Differential Effects of Eicosapentaenoic Acid (EPA) and Docosahexaenoic Acid (DHA) on Gene Expression Profiles of Stimulated Thp-1 Macrophages. <i>Nutrients</i> , 2017 , 9,	6.7	20	
40	PPARalpha L162V polymorphism alters the potential of n-3 fatty acids to increase lipoprotein lipase activity. <i>Molecular Nutrition and Food Research</i> , 2010 , 54, 543-50	5.9	20	
39	Cardiometabolic risk factors are influenced by Stearoyl-CoA Desaturase (SCD) -1 gene polymorphisms and n-3 polyunsaturated fatty acid supplementation. <i>Molecular Nutrition and Food Research</i> , 2014 , 58, 1079-86	5.9	19	
38	Regulation of endobiotics glucuronidation by ligand-activated transcription factors: physiological function and therapeutic potential. <i>Drug Metabolism Reviews</i> , 2010 , 42, 110-22	7	19	

37	Reduced Myelination and Increased Glia Reactivity Resulting from Severe Neonatal Hyperbilirubinemia. <i>Molecular Pharmacology</i> , 2016 , 89, 84-93	4.3	18
36	The human UGT1A3 enzyme conjugates norursodeoxycholic acid into a C23-ester glucuronide in the liver. <i>Journal of Biological Chemistry</i> , 2010 , 285, 1113-21	5.4	18
35	Extensive splicing of transcripts encoding the bile acid-conjugating enzyme UGT2B4 modulates glucuronidation. <i>Pharmacogenetics and Genomics</i> , 2010 , 20, 195-210	1.9	18
34	Whey protein hydrolysate and branched-chain amino acids downregulate inflammation-related genes in vascular endothelial cells. <i>Nutrition Research</i> , 2017 , 38, 43-51	4	17
33	Nuclear receptors and endobiotics glucuronidation: the good, the bad, and the UGT. <i>Drug Metabolism Reviews</i> , 2013 , 45, 34-47	7	16
32	UGT2B23, a novel uridine diphosphate-glucuronosyltransferase enzyme expressed in steroid target tissues that conjugates androgen and estrogen metabolites. <i>Endocrinology</i> , 1999 , 140, 5538-48	4.8	16
31	Glucuronidation of the nonsteroidal antiestrogen EM-652 (SCH 57068), by human and monkey steroid conjugating UDP-glucuronosyltransferase enzymes. <i>Molecular Pharmacology</i> , 2001 , 59, 636-45	4.3	15
30	Effect of S-adenosyl-L-methionine on liver biochemistry and quality of life in patients with primary biliary cholangitis treated with ursodeoxycholic acid. A prospective, open label pilot study. <i>Journal of Gastrointestinal and Liver Diseases</i> , 2018 , 27, 273-279	1.4	15
29	Liver Expression of Sulphotransferase 2A1 Enzyme Is Impaired in Patients with Primary Sclerosing Cholangitis: Lack of the Response to Enhanced Expression of PXR. <i>Journal of Immunology Research</i> , 2015 , 2015, 571353	4.5	14
28	Modulation of the biomarkers of inflammation and oxidative stress by ruminant trans fatty acids and dairy proteins in vascular endothelial cells (HUVEC). <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2017 , 126, 64-71	2.8	13
27	Enantiomer selective glucuronidation of the non-steroidal pure anti-androgen bicalutamide by human liver and kidney: role of the human UDP-glucuronosyltransferase (UGT)1A9 enzyme. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2013 , 113, 92-102	3.1	13
26	Treatment with a novel agent combining docosahexaenoate and metformin increases protectin DX and IL-6 production in skeletal muscle and reduces insulin resistance in obese diabetic db/db mice. <i>Diabetes, Obesity and Metabolism,</i> 2017 , 19, 313-319	6.7	12
25	Differences in transcriptional activation by the two allelic (L162V Polymorphic) variants of PPAR after Omega-3 fatty acids treatment. <i>PPAR Research</i> , 2009 , 2009, 369602	4.3	12
24	The Hepatokine TSK does not affect brown fat thermogenic capacity, body weight gain, and glucose homeostasis. <i>Molecular Metabolism</i> , 2019 , 30, 184-191	8.8	10
23	PPAREA Master Regulator of Bilirubin Homeostasis. PPAR Research, 2014, 2014, 747014	4.3	10
22	Urinary Elimination of Bile Acid Glucuronides under Severe Cholestatic Situations: Contribution of Hepatic and Renal Glucuronidation Reactions. <i>Canadian Journal of Gastroenterology and Hepatology</i> , 2018 , 2018, 8096314	2.8	8
21	Isolation and characterization of the monkey UGT2B30 gene that encodes a uridine diphosphate-glucuronosyltransferase enzyme active on mineralocorticoid, glucocorticoid, androgen and oestrogen hormones. <i>Biochemical Journal</i> , 2002 , 365, 213-22	3.8	8
20	Associations between branched chain amino acid levels, obesity and cardiometabolic complications. <i>Integrative Obesity and Diabetes</i> , 2015 , 1,		8

19	Low Endrenergic receptor level may promote development of castration resistant prostate cancer and altered steroid metabolism. <i>Oncotarget</i> , 2016 , 7, 1878-94	3.3	8
18	Cloning and characterization of a simian UDP-glucuronosyltransferase enzyme UGT2B20, a novel C19 steroid-conjugating protein. <i>Biochemical Journal</i> , 1999 , 337, 567	3.8	8
17	Selective and sensitive quantification of the cytochrome P450 3A4 protein in human liver homogenates through multiple reaction monitoring mass spectrometry. <i>Proteomics</i> , 2016 , 16, 2827-28	3 7 .8	7
16	Dairy Product Consumption Interacts with Glucokinase (GCK) Gene Polymorphisms Associated with Insulin Resistance. <i>Journal of Personalized Medicine</i> , 2017 , 7,	3.6	6
15	N-3 Polyunsaturated Fatty Acids Stimulate Bile Acid Detoxification in Human Cell Models. <i>Canadian Journal of Gastroenterology and Hepatology</i> , 2018 , 2018, 6031074	2.8	6
14	The androgen-conjugating uridine diphosphoglucuronosyltransferase-2B enzymes are differentially expressed temporally and spatially in the monkey follicle throughout the menstrual cycle. <i>Endocrinology</i> , 2001 , 142, 2499-507	4.8	6
13	Heterozygous Inactivation of the Nuclear Receptor PXR/NR1I2 in a Patient With Anabolic Steroid-Induced Intrahepatic Cholestasis. <i>Hepatitis Monthly</i> , 2016 , 16, e35953	1.8	5
12	Differential Role of Liver X Receptor (LXR) and LXR in the Regulation of UDP-Glucuronosyltransferase 1A1 in Humanized Mice. <i>Drug Metabolism and Disposition</i> , 2020 , 48, 255-2	263	4
11	Dietary fat and low fiber in purified diets differently impact the gut-liver axis to promote obesity-linked metabolic impairments. <i>American Journal of Physiology - Renal Physiology</i> , 2021 , 320, G10	0∮4 ¹ -Gʻ	1033
10	Surgical support during the terrorist attacks in Paris, November 13, 2015: Experience at Bigin Military Teaching Hospital. <i>Journal of Trauma and Acute Care Surgery</i> , 2017 , 82, 1122-1128	3.3	3
9	Reply:. <i>Hepatology</i> , 2007 , 45, 1084-1085	11.2	2
8	Fish oil replacement prevents, while docosahexaenoic acid-derived protectin DX mitigates end-stage-renal-disease in atherosclerotic diabetic mice. <i>FASEB Journal</i> , 2021 , 35, e21559	0.9	2
7	The Androgen-Conjugating Uridine Diphosphoglucuronosyltransferase-2B Enzymes Are Differentially Expressed Temporally and Spatially in the Monkey Follicle throughout the Menstrual Cyc	le	1
6	Intestinal UGT1A1 and protection against Irinotecan-induced toxicity in a novel UGT1A1 tissue-specific humanized mouse model. <i>Drug Metabolism and Disposition</i> , 2021 ,	4	1
5	High-Fat Diet Modulates Hepatic Amyloid hand Cerebrosterol Metabolism in the Triple Transgenic Mouse Model of Alzheimer's Disease. <i>Hepatology Communications</i> , 2021 , 5, 446-460	6	1
4	Bile acids during pregnancy: Trimester variations and associations with glucose homeostasis. <i>Health Science Reports</i> , 2021 , 4, e243	2.2	O
3	Adjunct Fenofibrate Up-regulates Bile Acid Glucuronidation and Improves Treatment Response For Patients With Cholestasis. <i>Hepatology Communications</i> , 2021 , 5, 2035-2051	6	О
2	Liquid chromatography coupled to tandem mass spectrometry methods for the selective and sensitive determination of 24S-hydroxycholesterol, its sulfate, and/or glucuronide conjugates in plasma <i>Journal of Mass Spectrometry</i> , 2022 , 57, e4827	2.2	O

Well-Tolerated Amphotericin B Derivatives That Effectively Treat Visceral Leishmaniasis. *ACS Infectious Diseases*, **2021**, 7, 2472-2482

5.5