

Diane M O brien

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/386647/diane-m-obrien-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

78
papers

2,491
citations

29
h-index

49
g-index

81
ext. papers

2,907
ext. citations

3.9
avg, IF

5.06
L-index

#	Paper	IF	Citations
78	Habitual Intake of Marine-derived n-3 Polyunsaturated Fatty Acids is Inversely Associated with a Cardiometabolic Inflammatory Profile in Yup'ik Alaska Native People. <i>Journal of Nutrition</i> , 2021 ,	4.1	3
77	Investigating the performance of 24-h urinary sucrose and fructose as a biomarker of total sugars intake in US participants - a controlled feeding study. <i>American Journal of Clinical Nutrition</i> , 2021 , 114, 721-730	7	2
76	Metabolic Characterization of Meat, Fish, and Soda Intake in Males: Secondary Results from a Randomized Inpatient Pilot Study. <i>Obesity</i> , 2021 , 29, 995-1002	8	
75	Validation of dried blood spot sampling for determining trophic positions of Arctic char using nitrogen stable isotope analyses of amino acids. <i>Rapid Communications in Mass Spectrometry</i> , 2021 , 35, e8992	2.2	1
74	Seasonal variation in added sugar or sugar sweetened beverage intake in Alaska native communities: an exploratory study. <i>International Journal of Circumpolar Health</i> , 2021 , 80, 1920779	1.7	1
73	The Breath Carbon Isotope Ratio Reflects Short-term Added-Sugar Intake in a Dose-Response, Crossover Feeding Study of 12 Healthy Adults. <i>Journal of Nutrition</i> , 2021 , 151, 628-635	4.1	2
72	The carbon isotope ratios of nonessential amino acids identify sugar-sweetened beverage (SSB) consumers in a 12-wk inpatient feeding study of 32 men with varying SSB and meat exposures. <i>American Journal of Clinical Nutrition</i> , 2021 , 113, 1256-1264	7	0
71	Characterizing Vegetable and Fruit Intake in a Remote Alaska Native Community Using Reflection Spectroscopy and 24-Hour Recalls. <i>Journal of Nutrition Education and Behavior</i> , 2021 , 53, 712-718	2	2
70	Does feeding type matter? Contribution of organic matter sources to benthic invertebrates on the Arctic Chukchi Sea shelf. <i>Food Webs</i> , 2021 , 29, e00205	1.8	1
69	Reply to Mantha OL et al. <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 1110-1111	7	
68	Dried Blood Spot Sampling of Landlocked Arctic Char (<i>Salvelinus alpinus</i>) for Estimating Mercury Exposure and Stable Carbon Isotope Fingerprinting of Essential Amino Acids. <i>Environmental Toxicology and Chemistry</i> , 2020 , 39, 893-903	3.8	2
67	Perspective: Dietary Biomarkers of Intake and Exposure-Exploration with Omics Approaches. <i>Advances in Nutrition</i> , 2020 , 11, 200-215	10	35
66	The Carbon Isotope Ratios of Serum Amino Acids in Combination with Participant Characteristics can be Used to Estimate Added Sugar Intake in a Controlled Feeding Study of US Postmenopausal Women. <i>Journal of Nutrition</i> , 2020 , 150, 2764-2771	4.1	3
65	Sources of primary production to Arctic bivalves identified using amino acid stable carbon isotope fingerprinting. <i>Isotopes in Environmental and Health Studies</i> , 2019 , 55, 366-384	1.5	8
64	The Nitrogen Isotope Ratio Is a Biomarker of Yup'ik Traditional Food Intake and Reflects Dietary Seasonality in Segmental Hair Analyses. <i>Journal of Nutrition</i> , 2019 , 149, 1960-1966	4.1	4
63	Strengthening adolescents' connection to their traditional food system improves diet quality in remote Alaska Native communities: results from the Neqa Elicarvigmun Pilot Study. <i>Translational Behavioral Medicine</i> , 2019 , 9, 952-961	3.2	7
62	Associations of plasma, RBCs, and hair carbon and nitrogen isotope ratios with fish, meat, and sugar-sweetened beverage intake in a 12-wk inpatient feeding study. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1306-1315	7	14

61	Morphological characteristics of harbor seal (<i>Phoca vitulina</i>) whiskers and their use in dietary reconstruction using stable isotope ratios. <i>Canadian Journal of Zoology</i> , 2018 , 96, 1255-1263	1.5	4
60	Serum Nitrogen and Carbon Stable Isotope Ratios Meet Biomarker Criteria for Fish and Animal Protein Intake in a Controlled Feeding Study of a Women's Health Initiative Cohort. <i>Journal of Nutrition</i> , 2018 , 148, 1931-1937	4.1	17
59	Stable isotope analysis of CO in breath indicates metabolic fuel shifts in torpid arctic ground squirrels. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2017 , 209, 10-15	2.6	3
58	Declines in traditional marine food intake and vitamin D levels from the 1960s to present in young Alaska Native women. <i>Public Health Nutrition</i> , 2017 , 20, 1738-1745	3.3	21
57	Bi-cultural dynamics for risk and protective factors for cardiometabolic health in an Alaska Native (Yup'ik) population. <i>PLoS ONE</i> , 2017 , 12, e0183451	3.7	2
56	Resource partitioning between Pacific walruses and bearded seals in the Alaska Arctic and sub-Arctic. <i>Oecologia</i> , 2017 , 184, 385-398	2.9	7
55	Dietary and genetic influences on hemostasis in a Yup'ik Alaska Native population. <i>PLoS ONE</i> , 2017 , 12, e0173616	3.7	4
54	Polymorphisms in stearoyl coa desaturase and sterol regulatory element binding protein interact with N-3 polyunsaturated fatty acid intake to modify associations with anthropometric variables and metabolic phenotypes in Yup'ik people. <i>Molecular Nutrition and Food Research</i> , 2016 , 60, 2642-2653	5.9	3
53	Linkage and association analysis of circulating vitamin D and parathyroid hormone identifies novel loci in Alaska Native Yup'ik people. <i>Genes and Nutrition</i> , 2016 , 11, 23	4.3	3
52	Genetics, Diet, and Season Are Associated with Serum 25-Hydroxycholecalciferol Concentration in a Yup'ik Study Population from Southwestern Alaska. <i>Journal of Nutrition</i> , 2016 , 146, 318-25	4.1	22
51	Feeding ecologies of key bivalve and polychaete species in the Bering Sea as elucidated by fatty acid and compound-specific stable isotope analyses. <i>Marine Ecology - Progress Series</i> , 2016 , 557, 161-175	2.6	12
50	The dominant detritus-feeding invertebrate in Arctic peat soils derives its essential amino acids from gut symbionts. <i>Journal of Animal Ecology</i> , 2016 , 85, 1275-85	4.7	30
49	Stable Isotope Ratios as Biomarkers of Diet for Health Research. <i>Annual Review of Nutrition</i> , 2015 , 35, 565-94	9.9	93
48	Linkage and association analysis of obesity traits reveals novel loci and interactions with dietary n-3 fatty acids in an Alaska Native (Yup'ik) population. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 689-97	2.7	17
47	Characterising the reproducibility and reliability of dietary patterns among Yup'ik Alaska Native people. <i>British Journal of Nutrition</i> , 2015 , 113, 634-43	3.6	13
46	Association between added sugar intake and dental caries in Yup'ik children using a novel hair biomarker. <i>BMC Oral Health</i> , 2015 , 15, 121	3.7	20
45	Sex, Adiposity, and Hypertension Status Modify the Inverse Effect of Marine Food Intake on Blood Pressure in Alaska Native (Yup'ik) People. <i>Journal of Nutrition</i> , 2015 , 145, 931-8	4.1	6
44	Validation of Stable Isotope Ratios as Biomarkers of Meat, Fish, and Soda Intake. <i>FASEB Journal</i> , 2015 , 29, 131.4	0.9	

43	DNA methylation patterns are associated with n-3 fatty acid intake in Yup'ik people. <i>Journal of Nutrition</i> , 2014 , 144, 425-30	4.1	43
42	Isotopic estimates of sugar intake are related to chronic disease risk factors but not obesity in an Alaska native (Yup'ik) study population. <i>European Journal of Clinical Nutrition</i> , 2014 , 68, 91-6	5.2	11
41	CDKAL1 and HHEX are associated with type 2 diabetes-related traits among Yup'ik people. <i>Journal of Diabetes</i> , 2014 , 6, 251-9	3.8	11
40	Anthropogenic food use and diet overlap between red foxes (<i>Vulpes vulpes</i>) and arctic foxes (<i>Vulpes lagopus</i>) in Prudhoe Bay, Alaska. <i>Canadian Journal of Zoology</i> , 2014 , 92, 657-663	1.5	15
39	Stable isotope models of sugar intake using hair, red blood cells, and plasma, but not fasting plasma glucose, predict sugar intake in a Yup'ik study population. <i>Journal of Nutrition</i> , 2014 , 144, 75-80	4.1	25
38	Using exploratory factor analysis of FFQ data to identify dietary patterns among Yup'ik people. <i>Public Health Nutrition</i> , 2014 , 17, 510-8	3.3	12
37	A stable isotope biomarker of marine food intake captures associations between n-3 fatty acid intake and chronic disease risk in a Yup'ik study population, and detects new associations with blood pressure and adiponectin. <i>Journal of Nutrition</i> , 2014 , 144, 706-13	4.1	22
36	Obesity polymorphisms identified in genome-wide association studies interact with n-3 polyunsaturated fatty acid intake and modify the genetic association with adiposity phenotypes in Yup'ik people. <i>Genes and Nutrition</i> , 2013 , 8, 495-505	4.3	16
35	The carbon isotope ratio of alanine in red blood cells is a new candidate biomarker of sugar-sweetened beverage intake. <i>Journal of Nutrition</i> , 2013 , 143, 878-84	4.1	39
34	Carbon and nitrogen stable isotope ratios predict intake of sweeteners in a Yup'ik study population. <i>Journal of Nutrition</i> , 2013 , 143, 161-5	4.1	37
33	Evidence for novel genetic loci associated with metabolic traits in Yup'ik people. <i>American Journal of Human Biology</i> , 2013 , 25, 673-80	2.7	9
32	Tracing carbon sources through aquatic and terrestrial food webs using amino acid stable isotope fingerprinting. <i>PLoS ONE</i> , 2013 , 8, e73441	3.7	145
31	Can amino acid carbon isotope ratios distinguish primary producers in a mangrove ecosystem?. <i>Rapid Communications in Mass Spectrometry</i> , 2012 , 26, 1541-8	2.2	30
30	Stable isotopes identify dietary changes associated with beak deformities in Black-capped Chickadees (<i>Poecile atricapillus</i>). <i>Auk</i> , 2012 , 129, 460-466	2.1	18
29	A test of alternative models for increased tissue nitrogen isotope ratios during fasting in hibernating arctic ground squirrels. <i>Journal of Experimental Biology</i> , 2012 , 215, 3354-61	3	40
28	Genetic polymorphisms in carnitine palmitoyltransferase 1A gene are associated with variation in body composition and fasting lipid traits in Yup'ik Eskimos. <i>Journal of Lipid Research</i> , 2012 , 53, 175-84	6.3	52
27	Stable nitrogen and carbon isotope ratios indicate traditional and market food intake in an indigenous circumpolar population. <i>Journal of Nutrition</i> , 2012 , 142, 84-90	4.1	54
26	Seasonal and annual variation in intake of traditional marine foods by a Yup'ik Eskimo population: a sequential dietary record from hair stable isotope analysis. <i>FASEB Journal</i> , 2012 , 26, lb347	0.9	

25	Associations between stable isotope ratios and animal-derived dietary protein intake in Yupik eskimos. <i>FASEB Journal</i> , 2012 , 26, lb342	0.9	
24	Associations of obesity with triglycerides and C-reactive protein are attenuated in adults with high red blood cell eicosapentaenoic and docosahexaenoic acids. <i>European Journal of Clinical Nutrition</i> , 2011 , 65, 808-17	5.2	51
23	Contrasting effects of nitrogen limitation and amino acid imbalance on carbon and nitrogen turnover in three species of Collembola. <i>Soil Biology and Biochemistry</i> , 2011 , 43, 749-759	7.5	24
22	Estimating lean mass over a wide range of body composition: a calibration of deuterium dilution in the arctic ground squirrel. <i>Rapid Communications in Mass Spectrometry</i> , 2011 , 25, 3491-6	2.2	6
21	Effects of Lipid Extraction on Stable Isotope Ratios in Avian Egg Yolk: Is Arithmetic Correction a Reliable Alternative?. <i>Auk</i> , 2010 , 127, 72-78	2.1	32
20	King eiders use an income strategy for egg production: a case study for incorporating individual dietary variation into nutrient allocation research. <i>Oecologia</i> , 2010 , 164, 1-12	2.9	23
19	Resource allocation to testes in walnut flies and implications for reproductive strategy. <i>Journal of Insect Physiology</i> , 2010 , 56, 1523-9	2.4	3
18	Relation between stable isotope ratios in human red blood cells and hair: implications for using the nitrogen isotope ratio of hair as a biomarker of eicosapentaenoic acid and docosahexaenoic acid. <i>American Journal of Clinical Nutrition</i> , 2009 , 90, 1642-7	7	39
17	Red blood cell delta15N: a novel biomarker of dietary eicosapentaenoic acid and docosahexaenoic acid intake. <i>American Journal of Clinical Nutrition</i> , 2009 , 89, 913-9	7	55
16	Disentangling effects of growth and nutritional status on seabird stable isotope ratios. <i>Oecologia</i> , 2009 , 159, 41-8	2.9	77
15	The influence of diet and water on the stable oxygen and hydrogen isotope composition of Chironomidae (Diptera) with paleoecological implications. <i>Oecologia</i> , 2009 , 160, 225-33	2.9	60
14	Stable isotope fingerprinting: a novel method for identifying plant, fungal, or bacterial origins of amino acids. <i>Ecology</i> , 2009 , 90, 3526-35	4.6	140
13	Use of stable isotopes to examine how dietary restriction extends <i>Drosophila</i> lifespan. <i>Current Biology</i> , 2008 , 18, R155-6	6.3	61
12	Using stable isotopes to assess carbon and nitrogen turnover in the Arctic sympagic amphipod <i>Onisimus litoralis</i> . <i>Oecologia</i> , 2008 , 158, 11-22	2.9	50
11	Age-related variation in red blood cell stable isotope ratios (delta13C and delta15N) from two Yupik villages in southwest Alaska: a pilot study. <i>International Journal of Circumpolar Health</i> , 2007 , 66, 31-41	1.7	29
10	Tracking human travel using stable oxygen and hydrogen isotope analyses of hair and urine. <i>Rapid Communications in Mass Spectrometry</i> , 2007 , 21, 2422-30	2.2	105
9	Resource allocation to reproduction and soma in <i>Drosophila</i> : a stable isotope analysis of carbon from dietary sugar. <i>Journal of Insect Physiology</i> , 2006 , 52, 763-70	2.4	39
8	An examination of the carbon isotope effects associated with amino acid biosynthesis. <i>Astrobiology</i> , 2006 , 6, 867-80	3.7	67

7	The amino acids used in reproduction by butterflies: a comparative study of dietary sources using compound-specific stable isotope analysis. <i>Physiological and Biochemical Zoology</i> , 2005 , 78, 819-27	2	66
6	Energy metabolism in orchid bee flight muscles: carbohydrate fuels all. <i>Journal of Experimental Biology</i> , 2005 , 208, 3573-9	3	91
5	Making eggs from nectar: the role of life history and dietary carbon turnover in butterfly reproductive resource allocation. <i>Oikos</i> , 2004 , 105, 279-291	4	109
4	Pollen feeding in the butterfly <i>Heliconius charitonia</i> : isotopic evidence for essential amino acid transfer from pollen to eggs. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2003 , 270, 2631-6	4.4	90
3	Renewable and nonrenewable resources: amino acid turnover and allocation to reproduction in Lepidoptera. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 4413-8	11.5	217
2	ALLOCATION TO REPRODUCTION IN A HAWKMOTH: A QUANTITATIVE ANALYSIS USING STABLE CARBON ISOTOPES. <i>Ecology</i> , 2000 , 81, 2822-2831	4.6	107
1	ALLOCATION TO REPRODUCTION IN A HAWKMOTH: A QUANTITATIVE ANALYSIS USING STABLE CARBON ISOTOPES 2000 , 81, 2822		5