

Sidney Strickland

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3866970/sidney-strickland-publications-by-year.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

65 papers	5,421 citations	37 h-index	67 g-index
67 ext. papers	6,059 ext. citations	9.8 avg, IF	5.75 L-index

#	Paper	IF	Citations
65	Plasmin-mediated cleavage of high-molecular-weight kininogen contributes to acetaminophen-induced acute liver failure. <i>Blood</i> , 2021 , 138, 259-272	2.2	4
64	The contact activation system and vascular factors as alternative targets for Alzheimer's disease therapy. <i>Research and Practice in Thrombosis and Haemostasis</i> , 2021 , 5, e12504	5.1	3
63	Factor XII plays a pathogenic role in organ failure and death in baboons challenged with <i>Staphylococcus aureus</i> . <i>Blood</i> , 2021 , 138, 178-189	2.2	5
62	Vascular endothelial growth factor associated dissimilar cerebrovascular phenotypes in two different mouse models of Alzheimer's Disease. <i>Neurobiology of Aging</i> , 2021 , 107, 96-108	5.6	0
61	Cerebral amyloid angiopathy-linked β amyloid mutations promote cerebral fibrin deposits via increased binding affinity for fibrinogen. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 14482-14492	11.5	10
60	Increased plasma bradykinin level is associated with cognitive impairment in Alzheimer's patients. <i>Neurobiology of Disease</i> , 2020 , 139, 104833	7.5	15
59	High molecular weight kininogen contributes to early mortality and kidney dysfunction in a mouse model of sickle cell disease. <i>Journal of Thrombosis and Haemostasis</i> , 2020 , 18, 2329-2340	15.4	3
58	Increased Contact System Activation in Mild Cognitive Impairment Patients with Impaired Short-Term Memory. <i>Journal of Alzheimer's Disease</i> , 2020 , 77, 59-65	4.3	6
57	A critical role for plasminogen in inflammation. <i>Journal of Experimental Medicine</i> , 2020 , 217,	16.6	20
56	Plasminogen mediates communication between the peripheral and central immune systems during systemic immune challenge with lipopolysaccharide. <i>Journal of Neuroinflammation</i> , 2019 , 16, 172	10.1	7
55	Long-Term Dabigatran Treatment Delays Alzheimer's Disease Pathogenesis in the TgCRND8 Mouse Model. <i>Journal of the American College of Cardiology</i> , 2019 , 74, 1910-1923	15.1	38
54	Knockdown of circulating C1 inhibitor induces neurovascular impairment, glial cell activation, neuroinflammation, and behavioral deficits. <i>Glia</i> , 2019 , 67, 1359-1373	9	16
53	Inflaming the Brain. <i>Neuron</i> , 2019 , 101, 991-993	13.9	4
52	Neutrophil adhesion in brain capillaries reduces cortical blood flow and impairs memory function in Alzheimer's disease mouse models. <i>Nature Neuroscience</i> , 2019 , 22, 413-420	25.5	152
51	An antibody against HK blocks Alzheimer's disease peptide β amyloid-induced bradykinin release in human plasma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 22921-22923	11.5	11
50	Aminopyrimidine Class Aggregation Inhibitor Effectively Blocks A β -Fibrinogen Interaction and A β -Induced Contact System Activation. <i>Biochemistry</i> , 2018 , 57, 1399-1409	3.2	9
49	A novel detection method of cleaved plasma high-molecular-weight kininogen reveals its correlation with Alzheimer's pathology and cognitive impairment. <i>Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring</i> , 2018 , 10, 480-489	5.2	16

48	Blood will out: vascular contributions to Alzheimer's disease. <i>Journal of Clinical Investigation</i> , 2018 , 128, 556-563	15.9	52
47	Analysis of β Amyloid-induced Abnormalities on Fibrin Clot Structure by Spectroscopy and Scanning Electron Microscopy. <i>Journal of Visualized Experiments</i> , 2018 ,	1.6	1
46	Blood-derived plasminogen drives brain inflammation and plaque deposition in a mouse model of Alzheimer's disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, E9687-E9696	11.5	39
45	Abnormal clotting of the intrinsic/contact pathway in Alzheimer disease patients is related to cognitive ability. <i>Blood Advances</i> , 2018 , 2, 954-963	7.8	21
44	Depletion of coagulation factor XII ameliorates brain pathology and cognitive impairment in Alzheimer disease mice. <i>Blood</i> , 2017 , 129, 2547-2556	2.2	54
43	Fibrinogen in the Nervous System: Glia Beware. <i>Neuron</i> , 2017 , 96, 951-953	13.9	5
42	Interactions of β Amyloid peptide with fibrinogen and coagulation factor XII may contribute to Alzheimer's disease. <i>Current Opinion in Hematology</i> , 2017 , 24, 427-431	3.3	24
41	Impact of the Coagulation System on the Pathogenesis of Alzheimer's Disease. <i>Blood</i> , 2017 , 130, SCI-3-SC123	1.3	1
40	Biochemical and structural analysis of the interaction between β Amyloid and fibrinogen. <i>Blood</i> , 2016 , 128, 1144-51	2.2	40
39	Laminin regulates PDGFR(+) cell stemness and muscle development. <i>Nature Communications</i> , 2016 , 7, 11415	17.4	29
38	A possible new role for A β in vascular and inflammatory dysfunction in Alzheimer's disease. <i>Thrombosis Research</i> , 2016 , 141 Suppl 2, S59-61	8.2	22
37	The cellular origin of laminin determines its role in blood pressure regulation. <i>Cellular and Molecular Life Sciences</i> , 2015 , 72, 999-1008	10.3	11
36	Activation of the factor XII-driven contact system in Alzheimer's disease patient and mouse model plasma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 4068-73	11.5	66
35	Fibrin deposited in the Alzheimer's disease brain promotes neuronal degeneration. <i>Neurobiology of Aging</i> , 2015 , 36, 608-17	5.6	106
34	Chronic Hypertension Leads to Neurodegeneration in the TgSwDI Mouse Model of Alzheimer's Disease. <i>Hypertension</i> , 2015 , 66, 175-82	8.5	63
33	Astrocytic laminin regulates pericyte differentiation and maintains blood brain barrier integrity. <i>Nature Communications</i> , 2014 , 5, 3413	17.4	195
32	A novel A β fibrinogen interaction inhibitor rescues altered thrombosis and cognitive decline in Alzheimer's disease mice. <i>Journal of Experimental Medicine</i> , 2014 , 211, 1049-62	16.6	76
31	The APOE e4/e4 genotype potentiates vascular fibrin(ogen) deposition in amyloid-laden vessels in the brains of Alzheimer's disease patients. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2013 , 33, 1251-8	7.3	100

30	Ablation of astrocytic laminin impairs vascular smooth muscle cell function and leads to hemorrhagic stroke. <i>Journal of Cell Biology</i> , 2013 , 202, 381-95	7.3	72
29	Allylamine delays fibrin clot lysis by altering fibrin structure and attenuating plasminogen binding to fibrin. <i>Blood</i> , 2012 , 119, 3342-51	2.2	80
28	Fibrinogen and altered hemostasis in Alzheimer's disease. <i>Journal of Alzheimer's Disease</i> , 2012 , 32, 599-608	4.9	106
27	Mesenchymal stem cells facilitate axon sorting, myelination, and functional recovery in paralyzed mice deficient in Schwann cell-derived laminin. <i>Glia</i> , 2011 , 59, 267-77	9	40
26	Laminin Expression Is Necessary for Maintenance of the Vascular Hematopoietic Niche. <i>Blood</i> , 2011 , 118, 217-217	2.2	
25	Alzheimer's disease peptide beta-amyloid interacts with fibrinogen and induces its oligomerization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 21812-7	11.5	145
24	Fibrinogen and beta-amyloid association alters thrombosis and fibrinolysis: a possible contributing factor to Alzheimer's disease. <i>Neuron</i> , 2010 , 66, 695-709	13.9	226
23	Laminin is required for Schwann cell morphogenesis. <i>Journal of Cell Science</i> , 2009 , 122, 929-36	5.3	53
22	Maximum-entropy network analysis reveals a role for tumor necrosis factor in peripheral nerve development and function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 12494-9	11.5	12
21	Disruption of laminin in the peripheral nervous system impedes nonmyelinating Schwann cell development and impairs nociceptive sensory function. <i>Glia</i> , 2009 , 57, 850-9	9	34
20	Cortical deficiency of laminin gamma1 impairs the AKT/GSK-3beta signaling pathway and leads to defects in neurite outgrowth and neuronal migration. <i>Developmental Biology</i> , 2009 , 327, 158-68	3.1	68
19	Regulation of Schwann cell function by the extracellular matrix. <i>Glia</i> , 2008 , 56, 1498-507	9	131
18	Fibrin deposition accelerates neurovascular damage and neuroinflammation in mouse models of Alzheimer's disease. <i>Journal of Experimental Medicine</i> , 2007 , 204, 1999-2008	16.6	227
17	Schwann cell-specific ablation of laminin gamma1 causes apoptosis and prevents proliferation. <i>Journal of Neuroscience</i> , 2005 , 25, 4463-72	6.6	123
16	Fibrin depletion decreases inflammation and delays the onset of demyelination in a tumor necrosis factor transgenic mouse model for multiple sclerosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 6698-703	11.5	109
15	The hippocampal laminin matrix is dynamic and critical for neuronal survival. <i>Molecular Biology of the Cell</i> , 2003 , 14, 2665-76	3.5	54
14	The tissue plasminogen activator-plasminogen proteolytic cascade accelerates amyloid-beta (Abeta) degradation and inhibits Abeta-induced neurodegeneration. <i>Journal of Neuroscience</i> , 2003 , 23, 8867-71	6.6	188
13	The possible role of tissue-type plasminogen activator (tPA) and tPA blockers in the pathogenesis and treatment of Alzheimer's disease. <i>Journal of Molecular Neuroscience</i> , 2003 , 20, 287-9	3.3	15

12	Laminin gamma1 is critical for Schwann cell differentiation, axon myelination, and regeneration in the peripheral nerve. <i>Journal of Cell Biology</i> , 2003 , 163, 889-99	7.3	221
11	Nervous system pathology: the fibrin perspective. <i>Biological Chemistry</i> , 2002 , 383, 37-45	4.5	44
10	Fibrin inhibits peripheral nerve remyelination by regulating Schwann cell differentiation. <i>Neuron</i> , 2002 , 33, 861-75	13.9	150
9	Cortex, a Drosophila gene required to complete oocyte meiosis, is a member of the Cdc20/fizzy protein family. <i>Genesis</i> , 2001 , 29, 141-52	1.9	56
8	Tissue plasminogen activator-mediated fibrinolysis protects against axonal degeneration and demyelination after sciatic nerve injury. <i>Journal of Cell Biology</i> , 2000 , 149, 1157-66	7.3	152
7	The only function of Grauzone required for Drosophila oocyte meiosis is transcriptional activation of the cortex gene. <i>Genetics</i> , 2000 , 155, 1831-9	4	17
6	Mapping of Drosophila mutations using site-specific male recombination. <i>Genetics</i> , 1998 , 149, 157-63	4	69
5	Neuronal death in the hippocampus is promoted by plasmin-catalyzed degradation of laminin. <i>Cell</i> , 1997 , 91, 917-25	56.2	562
4	An extracellular proteolytic cascade promotes neuronal degeneration in the mouse hippocampus. <i>Journal of Neuroscience</i> , 1997 , 17, 543-52	6.6	382
3	Neuronal cell death and tPA. <i>Nature</i> , 1996 , 384, 123-4	50.4	210
2	Excitotoxin-induced neuronal degeneration and seizure are mediated by tissue plasminogen activator. <i>Nature</i> , 1995 , 377, 340-4	50.4	609
1	Tissue plasminogen activator mRNA in murine tissues. <i>FEBS Letters</i> , 1988 , 229, 100-6	3.8	40