

Margaret Heslin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3877446/publications.pdf>

Version: 2024-02-01

58
papers

1,955
citations

361296

20
h-index

265120

42
g-index

59
all docs

59
docs citations

59
times ranked

3455
citing authors

#	ARTICLE	IF	CITATIONS
1	Antidepressant and antipsychotic treatment of Psychotic Major Depression in a British mental healthcare setting. <i>Journal of Mental Health</i> , 2023, 32, 71-77.	1.0	1
2	Assessing the feasibility of a web-based outcome measurement system in child and adolescent mental health services – myHealthE a randomised controlled feasibility pilot study. <i>Child and Adolescent Mental Health</i> , 2023, 28, 128-147.	1.8	4
3	Rethinking the course of psychotic disorders: modelling long-term symptom trajectories. <i>Psychological Medicine</i> , 2022, 52, 2641-2650.	2.7	11
4	Effectiveness and cost-effectiveness of psychiatric mother and baby units: quasi-experimental study. <i>British Journal of Psychiatry</i> , 2022, 221, 628-636.	1.7	9
5	Perinatal mental health services in pregnancy and the year after birth: the ESMI research programme including RCT. <i>Programme Grants for Applied Research</i> , 2022, 10, 1-142.	0.4	9
6	Neurological Signs at the First Psychotic Episode as Correlates of Long-Term Outcome: Results From the AESOP-10 Study. <i>Schizophrenia Bulletin</i> , 2021, 47, 118-127.	2.3	20
7	Diagnostic stability and outcome after first episode psychosis. <i>Journal of Mental Health</i> , 2021, 30, 104-112.	1.0	9
8	Association between air pollution exposure and mental health service use among individuals with first presentations of psychotic and mood disorders: retrospective cohort study. <i>British Journal of Psychiatry</i> , 2021, 219, 678-685.	1.7	40
9	Recovery From Severe Mental Health Problems: A Systematic Review of Service User and Informal Caregiver Perspectives. <i>Frontiers in Psychiatry</i> , 2021, 12, 712026.	1.3	18
10	Cost-Effectiveness of Combination Disease-Modifying Antirheumatic Drugs Versus Tumor Necrosis Factor Inhibitors in Active Rheumatoid Arthritis: A Pragmatic, Randomized, Multicenter Trial. <i>Arthritis Care and Research</i> , 2020, 72, 334-342.	1.5	6
11	Symptom remission at 12-weeks strongly predicts long-term recovery from the first episode of psychosis. <i>Psychological Medicine</i> , 2020, 50, 1452-1462.	2.7	16
12	An exploratory parallel-group randomised controlled trial of antenatal Guided Self-Help (plus usual) Disorders, 2020, 261, 187-197.	2.0	14
13	Reasons behind the rising rate of involuntary admissions under the Mental Health Act (1983): Service use and cost impact. <i>International Journal of Law and Psychiatry</i> , 2020, 68, 101506.	0.5	16
14	Out of pocket expenses in obsessive compulsive disorder. <i>Journal of Mental Health</i> , 2020, , 1-6.	1.0	0
15	Psychological interventions for people with psychotic experiences: A systematic review and meta-analysis of controlled and uncontrolled effectiveness and economic studies. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 673-695.	1.3	12
16	Low-Intensity Guided Help Through Mindfulness (LIGHTMIND): study protocol for a randomised controlled trial comparing supported mindfulness-based cognitive therapy self-help to supported cognitive behavioural therapy self-help for adults experiencing depression. <i>Trials</i> , 2020, 21, 374.	0.7	7
17	A health promotion intervention to improve lifestyle choices and health outcomes in people with psychosis: a research programme including the IMPaCT RCT. <i>Programme Grants for Applied Research</i> , 2020, 8, 1-124.	0.4	3
18	Guidance for researchers wanting to link NHS data using non-consent approaches: a thematic analysis of feedback from the Health Research Authority Confidentiality Advisory Group. <i>International Journal of Population Data Science</i> , 2020, 5, 1355.	0.1	2

#	ARTICLE	IF	CITATIONS
19	Protocol for a quasi-experimental study of the effectiveness and cost-effectiveness of mother and baby units compared with general psychiatric inpatient wards and crisis resolution team services (The Tj ETQq1 1 0784314 ngBT /Ov	0.784314	19
20	A qualitative study of the acceptability of cognitive bias modification for paranoia (CBM-pa) in patients with psychosis. <i>BMC Psychiatry</i> , 2019, 19, 225.	1.1	10
21	Effect of lifestyle, medication and ethnicity on cardiometabolic risk in the year following the first episode of psychosis: prospective cohort study. <i>British Journal of Psychiatry</i> , 2019, 215, 712-719.	1.7	25
22	Psychometric properties of the five-level EuroQoL-5 dimension and Short Form-6 dimension measures of health-related quality of life in a population of pregnant women with depression. <i>BJPsych Open</i> , 2019, 5, e88.	0.3	13
23	Psychological interventions for people with psychotic experiences: protocol for a systematic review and meta-analysis. <i>Systematic Reviews</i> , 2019, 8, 124.	2.5	2
24	For Baby's Sake: Intervention Development and Evaluation Design of a Whole-Family Perinatal Intervention to Break the Cycle of Domestic Abuse. <i>Journal of Family Violence</i> , 2019, 34, 539-551.	2.1	20
25	A Comparison of Different Approaches for Costing Medication Use in an Economic Evaluation. <i>Value in Health</i> , 2018, 21, 185-192.	0.1	6
26	Psychotic major depression: challenges in clinical practice and research. <i>British Journal of Psychiatry</i> , 2018, 212, 131-133.	1.7	18
27	Comparison of effectiveness and cost-effectiveness of an intensive community supported discharge service versus treatment as usual for adolescents with psychiatric emergencies: a randomised controlled trial. <i>Lancet Psychiatry</i> , 2018, 5, 477-485.	3.7	70
28	Early sustained recovery following first episode psychosis: Evidence from the AESOP10 follow-up study. <i>Schizophrenia Research</i> , 2018, 199, 341-345.	1.1	12
29	Inpatient use and area-level socio-environmental factors in people with psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 1133-1140.	1.6	8
30	Nurse-led psychological intervention for type 2 diabetes: a cluster randomised controlled trial (Diabetes-6 study) in primary care. <i>British Journal of General Practice</i> , 2018, 68, e531-e540.	0.7	26
31	5.2 RETHINKING THE COURSE OF PSYCHOTIC DISORDERS: IDENTIFYING LATENT TRAJECTORIES. <i>Schizophrenia Bulletin</i> , 2018, 44, S6-S6.	2.3	0
32	Costs of the police service and mental healthcare pathways experienced by individuals with enduring mental health needs. <i>British Journal of Psychiatry</i> , 2017, 210, 157-164.	1.7	19
33	Antipsychotic treatment resistance in first-episode psychosis: prevalence, subtypes and predictors. <i>Psychological Medicine</i> , 2017, 47, 1981-1989.	2.7	200
34	Scaling-up services for psychosis, depression and epilepsy in sub-Saharan Africa and South Asia: development and application of a mental health systems planning tool (OneHealth). <i>Epidemiology and Psychiatric Sciences</i> , 2017, 26, 234-244.	1.8	47
35	Cognitive Bias Modification for paranoia (CBM-pa): study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 298.	0.7	15
36	Ethnicity and long-term course and outcome of psychotic disorders in a UK sample: The AESOP-10 study. <i>British Journal of Psychiatry</i> , 2017, 211, 88-94.	1.7	71

#	ARTICLE	IF	CITATIONS
37	Low-intensity cognitive-behaviour therapy interventions for obsessive-compulsive disorder compared to waiting list for therapist-led cognitive-behaviour therapy: 3-arm randomised controlled trial of clinical effectiveness. <i>PLoS Medicine</i> , 2017, 14, e1002337.	3.9	25
38	Randomised controlled trial to improve health and reduce substance use in established psychosis (IMPACT): cost-effectiveness of integrated psychosocial health promotion. <i>BMC Psychiatry</i> , 2017, 17, 407.	1.1	10
39	Clinical effectiveness, cost-effectiveness and acceptability of low-intensity interventions in the management of obsessive-compulsive disorder: the Obsessive-Compulsive Treatment Efficacy randomised controlled Trial (OCTET). <i>Health Technology Assessment</i> , 2017, 21, 1-132.	1.3	41
40	Ten-year outcomes in first episode psychotic major depression patients compared with schizophrenia and bipolar patients. <i>Schizophrenia Research</i> , 2016, 176, 417-422.	1.1	23
41	Depression: an exploratory parallel-group randomised controlled trial of Antenatal guided self help for WomeN (DAWN): study protocol for a randomised controlled trial. <i>Trials</i> , 2016, 17, 503.	0.7	13
42	Outcomes following first-episode psychosis – Why we should intervene early in all ages, not only in youth. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 1055-1063.	1.3	25
43	Decision analytic model exploring the cost and cost-offset implications of street triage. <i>BMJ Open</i> , 2016, 6, e009670.	0.8	18
44	A systematic review of the economic evidence for interventions for family carers of stroke patients. <i>Clinical Rehabilitation</i> , 2016, 30, 119-133.	1.0	13
45	Diagnostic Stability of ICD/DSM First Episode Psychosis Diagnoses: Meta-analysis. <i>Schizophrenia Bulletin</i> , 2016, 42, 1395-1406.	2.3	151
46	Biological and psychosocial risk factors for psychotic major depression. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 233-245.	1.6	19
47	Diagnostic change 10 years after a first episode of psychosis. <i>Psychological Medicine</i> , 2015, 45, 2757-2769.	2.7	74
48	Ten-Year Outcomes of First-Episode Psychoses in the MRC A†SOP-10 Study. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 379-386.	0.5	90
49	Tumour necrosis factor inhibitors versus combination intensive therapy with conventional disease modifying anti-rheumatic drugs in established rheumatoid arthritis: TACIT non-inferiority randomised controlled trial. <i>BMJ</i> , The, 2015, 350, h1046-h1046.	3.0	59
50	Mortality in Schizophrenia and Other Psychoses: A 10-Year Follow-up of the Ó”SOP First-Episode Cohort. <i>Schizophrenia Bulletin</i> , 2015, 41, 664-673.	2.3	128
51	Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study – CORRIGENDUM. <i>Psychological Medicine</i> , 2014, 44, 2727-2727.	2.7	2
52	Randomised controlled trial of Tumour necrosis factor inhibitors Against Combination Intensive Therapy with conventional disease-modifying antirheumatic drugs in established rheumatoid arthritis: the TACIT trial and associated systematic reviews. <i>Health Technology Assessment</i> , 2014, 18, 1-164.	1.3	17
53	Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study. <i>Psychological Medicine</i> , 2014, 44, 2713-2726.	2.7	192
54	Bilateral hippocampal increase following first-episode psychosis is associated with good clinical, functional and cognitive outcomes. <i>Psychological Medicine</i> , 2014, 44, 1279-1291.	2.7	40

#	ARTICLE	IF	CITATIONS
55	Randomized controlled trial of supported employment in England: 2 year follow-up of the Supported Work and Needs (SWAN) study. <i>World Psychiatry</i> , 2011, 10, 132-137.	4.8	59
56	The economic cost of pathways to care in first episode psychosis. <i>International Review of Psychiatry</i> , 2011, 23, 55-60.	1.4	3
57	Supported employment: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2010, 196, 404-411.	1.7	82
58	Multiple Sclerosis in the UK. <i>Pharmacoeconomics</i> , 2008, 26, 847-860.	1.7	96