

Zoltan Pozsonyi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3879060/publications.pdf>

Version: 2024-02-01

19
papers

855
citations

840776

11
h-index

839539

18
g-index

20
all docs

20
docs citations

20
times ranked

1548
citing authors

#	ARTICLE	IF	CITATIONS
1	Red cell distribution width in heart failure: Prediction of clinical events and relationship with markers of ineffective erythropoiesis, inflammation, renal function, and nutritional state. <i>American Heart Journal</i> , 2009, 158, 659-666.	2.7	525
2	Interaction of serum 70-kDa heat shock protein levels and HspA1B (+1267) gene polymorphism with disease severity in patients with chronic heart failure. <i>Cell Stress and Chaperones</i> , 2008, 13, 199-206.	2.9	42
3	Levels of von Willebrand factor antigen and von Willebrand factor cleaving protease (ADAMTS13) activity predict clinical events in chronic heart failure. <i>Thrombosis and Haemostasis</i> , 2009, 102, 573-580.	3.4	40
4	Copeptin (C-terminal pro Arginine-Vasopressin) is an Independent Long-Term Prognostic Marker in Heart Failure with Reduced Ejection Fraction. <i>Heart Lung and Circulation</i> , 2015, 24, 359-367.	0.4	39
5	Elevated extracellular HSP70 (HSPA1A) level as an independent prognostic marker of mortality in patients with heart failure. <i>Cell Stress and Chaperones</i> , 2013, 18, 809-813.	2.9	34
6	Association of Ficolin-3 with Severity and Outcome of Chronic Heart Failure. <i>PLoS ONE</i> , 2013, 8, e60976.	2.5	34
7	Complement anaphylatoxin C3a as a novel independent prognostic marker in heart failure. <i>Clinical Research in Cardiology</i> , 2012, 101, 607-615.	3.3	26
8	Long-Term Survival and Apolipoprotein A1 Level in Chronic Heart Failure: Interaction With Tumor Necrosis Factor \pm 308 G/A Polymorphism. <i>Journal of Cardiac Failure</i> , 2017, 23, 113-120.	1.7	19
9	Adrenomedullin and endothelin-1 are related to inflammation in chronic heart failure. <i>Inflammation Research</i> , 2009, 58, 298-305.	4.0	18
10	Red cell distribution width: a powerful prognostic marker in heart failure. <i>European Journal of Heart Failure</i> , 2010, 12, 415-415.	7.1	17
11	Serum soluble E-selectin and NT-proBNP levels additively predict mortality in diabetic patients with chronic heart failure. <i>Clinical Research in Cardiology</i> , 2011, 100, 587-594.	3.3	14
12	Coexistence of aortic valve stenosis and cardiac amyloidosis: echocardiographic and clinical significance. <i>Cardiovascular Ultrasound</i> , 2019, 17, 32.	1.6	12
13	Severe Mitral Regurgitation and Heart Failure due to Caseous Calcification of the Mitral Annulus. <i>Cardiology</i> , 2011, 118, 79-82.	1.4	10
14	Hemochromatosis and Hemojuvelin G320V Homozygosity in a Hungarian Woman. <i>Acta Haematologica</i> , 2010, 123, 191-193.	1.4	7
15	Red cell distribution width as predictive marker in CHF: Testing of model performance by reclassification methods. <i>International Journal of Cardiology</i> , 2014, 174, 783-785.	1.7	7
16	Successful thrombolysis of late, non-obstructive mitral bioprosthetic valve thrombosis: case report and review of the literature. <i>Journal of Heart Valve Disease</i> , 2011, 20, 526-30.	0.5	5
17	Variant Transthyretin Amyloidosis (ATTRv) in Hungary: First Data on Epidemiology and Clinical Features. <i>Genes</i> , 2021, 12, 1152.	2.4	4
18	994 Correlations of spontaneous echocontrast and inflammatory, haemostaseologic and echocardiographic parameters in atrial fibrillation. <i>European Journal of Echocardiography</i> , 2005, 6, S159-S159.	2.3	0

#	ARTICLE	IF	CITATIONS
19	Multiple Valvular Complications of Hypereosinophilic Syndrome. Journal of Heart Valve Disease, 2016, 25, 752-755.	0.5	0