

Ruediger Klein

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3884209/ruediger-klein-publications-by-year.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

176 papers	31,173 citations	84 h-index	176 g-index
190 ext. papers	33,586 ext. citations	15.9 avg, IF	6.92 L-index

#	Paper	IF	Citations
176	Amyloid-like aggregating proteins cause lysosomal defects in neurons via gain-of-function toxicity.. <i>Life Science Alliance</i> , 2022 , 5,	5.8	4
175	Natural loss of function of ephrin-B3 shapes spinal flight circuitry in birds. <i>Science Advances</i> , 2021 , 7,	14.3	1
174	FLRT2 and FLRT3 Cooperate in Maintaining the Tangential Migratory Streams of Cortical Interneurons during Development. <i>Journal of Neuroscience</i> , 2021 , 41, 7350-7362	6.6	1
173	Recent advances in inter-cellular interactions during neural circuit assembly. <i>Current Opinion in Neurobiology</i> , 2021 , 69, 25-32	7.6	2
172	Fluc-EGFP reporter mice reveal differential alterations of neuronal proteostasis in aging and disease. <i>EMBO Journal</i> , 2021 , 40, e107260	13	5
171	FLRT3 Marks Direction-Selective Retinal Ganglion Cells That Project to the Medial Terminal Nucleus.. <i>Frontiers in Molecular Neuroscience</i> , 2021 , 14, 790466	6.1	1
170	Structural Basis of Teneurin-Latrophilin Interaction in Repulsive Guidance of Migrating Neurons. <i>Cell</i> , 2020 , 180, 323-339.e19	56.2	37
169	The Insula Cortex Contacts Distinct Output Streams of the Central Amygdala. <i>Journal of Neuroscience</i> , 2020 , 40, 8870-8882	6.6	4
168	Spinal Inhibitory Ptf1a-Derived Neurons Prevent Self-Generated Itch. <i>Cell Reports</i> , 2020 , 33, 108422	10.6	4
167	Identification of Spinal Neurons Contributing to the Dorsal Column Projection Mediating Fine Touch and Corrective Motor Movements. <i>Neuron</i> , 2019 , 104, 749-764.e6	13.9	13
166	Cortical circuit alterations precede motor impairments in Huntington's disease mice. <i>Scientific Reports</i> , 2019 , 9, 6634	4.9	22
165	Gulp1 controls Eph/ephrin trogocytosis and is important for cell rearrangements during development. <i>Journal of Cell Biology</i> , 2019 , 218, 3455-3471	7.3	10
164	Regulation of Cerebral Cortex Folding by Controlling Neuronal Migration via FLRT Adhesion Molecules. <i>Cell</i> , 2017 , 169, 621-635.e16	56.2	72
163	Placental labyrinth formation in mice requires endothelial FLRT2/UNC5B signaling. <i>Development (Cambridge)</i> , 2017 , 144, 2392-2401	6.6	14
162	Central amygdala circuits modulate food consumption through a positive-valence mechanism. <i>Nature Neuroscience</i> , 2017 , 20, 1384-1394	25.5	112
161	In Situ Architecture and Cellular Interactions of PolyQ Inclusions. <i>Cell</i> , 2017 , 171, 179-187.e10	56.2	177
160	Spatiotemporal Proteomic Profiling of Huntington's Disease Inclusions Reveals Widespread Loss of Protein Function. <i>Cell Reports</i> , 2017 , 21, 2291-2303	10.6	71

159	Intrinsic Circuits in the Lateral Central Amygdala. <i>ENeuro</i> , 2017 , 4,	3.9	29
158	Exosomes mediate cell contact-independent ephrin-Eph signaling during axon guidance. <i>Journal of Cell Biology</i> , 2016 , 214, 35-44	7.3	73
157	Mechanisms of ephrin-Eph signalling in development, physiology and disease. <i>Nature Reviews Molecular Cell Biology</i> , 2016 , 17, 240-56	48.7	317
156	Multimodal Eph/Ephrin signaling controls several phases of urogenital development. <i>Kidney International</i> , 2016 , 90, 373-388	9.9	6
155	Super-complexes of adhesion GPCRs and neural guidance receptors. <i>Nature Communications</i> , 2016 , 7, 11184	17.4	53
154	Structural Perspectives on Axon Guidance. <i>Annual Review of Cell and Developmental Biology</i> , 2016 , 32, 577-608	12.6	62
153	Tiam-Rac signaling mediates trans-endocytosis of ephrin receptor EphB2 and is important for cell repulsion. <i>Journal of Cell Biology</i> , 2016 , 214, 735-52	7.3	19
152	Structural basis of latrophilin-FLRT interaction. <i>Structure</i> , 2015 , 23, 774-81	5.2	45
151	The Eph Receptor Family 2015 , 165-264		2
150	Cis and trans RET signaling control the survival and central projection growth of rapidly adapting mechanoreceptors. <i>ELife</i> , 2015 , 4, e06828	8.9	20
149	Ret rescues mitochondrial morphology and muscle degeneration of Drosophila Pink1 mutants. <i>EMBO Journal</i> , 2014 , 33, 341-55	13	52
148	Ephrin signalling in the developing nervous system. <i>Current Opinion in Neurobiology</i> , 2014 , 27, 16-24	7.6	62
147	The composition of EphB2 clusters determines the strength in the cellular repulsion response. <i>Journal of Cell Biology</i> , 2014 , 204, 409-22	7.3	59
146	Genetic evidence for the adhesion protein IgSF9/Dasm1 to regulate inhibitory synapse development independent of its intracellular domain. <i>Journal of Neuroscience</i> , 2014 , 34, 4187-99	6.6	20
145	EphA4-mediated ipsilateral corticospinal tract misprojections are necessary for bilateral voluntary movements but not bilateral stereotypic locomotion. <i>Journal of Neuroscience</i> , 2014 , 34, 5211-21	6.6	29
144	FLRT structure: balancing repulsion and cell adhesion in cortical and vascular development. <i>Neuron</i> , 2014 , 84, 370-85	13.9	70
143	EphA4 receptor shedding regulates spinal motor axon guidance. <i>Current Biology</i> , 2014 , 24, 2355-65	6.3	24
142	FLRT3 is a Robo1-interacting protein that determines Netrin-1 attraction in developing axons. <i>Current Biology</i> , 2014 , 24, 494-508	6.3	59

141	Structurally encoded intraclass differences in EphA clusters drive distinct cell responses. <i>Nature Structural and Molecular Biology</i> , 2013 , 20, 958-64	17.6	69
140	Ephrins and Eph Receptors [Synaptogenesis and Synaptic Function 2013 , 659-670		
139	Ephrin-B1 controls the columnar distribution of cortical pyramidal neurons by restricting their tangential migration. <i>Neuron</i> , 2013 , 79, 1123-35	13.9	46
138	EphrinB3/EphA4-mediated guidance of ascending and descending spinal tracts. <i>Neuron</i> , 2013 , 80, 1407-20	13.9	45
137	Protein tyrosine phosphatase receptor type O inhibits trigeminal axon growth and branching by repressing TrkB and Ret signaling. <i>Journal of Neuroscience</i> , 2013 , 33, 5399-410	6.6	17
136	Integration of guidance cues: parallel signaling and crosstalk. <i>Trends in Neurosciences</i> , 2013 , 36, 295-304	13.3	75
135	c-Jun in Schwann cells promotes axonal regeneration and motoneuron survival via paracrine signaling. <i>Journal of Cell Biology</i> , 2012 , 198, 127-41	7.3	166
134	Eph/ephrin signalling during development. <i>Development (Cambridge)</i> , 2012 , 139, 4105-9	6.6	144
133	Genetic evidence for a contribution of EphA:ephrinA reverse signaling to motor axon guidance. <i>Journal of Neuroscience</i> , 2012 , 32, 5209-15	6.6	36
132	Repairing the parkinsonian brain with neurotrophic factors. <i>Trends in Neurosciences</i> , 2011 , 34, 88-100	13.3	90
131	The axon's balancing act: cis- and trans-interactions between Ephs and ephrins. <i>Neuron</i> , 2011 , 71, 1-3	13.9	16
130	Anatomical coupling of sensory and motor nerve trajectory via axon tracking. <i>Neuron</i> , 2011 , 71, 263-77	13.9	45
129	Pitx3 is a critical mediator of GDNF-induced BDNF expression in nigrostriatal dopaminergic neurons. <i>Journal of Neuroscience</i> , 2011 , 31, 12802-15	6.6	74
128	Rüdiger Klein: reading the guideposts for axon guidance. Interview by Caitlin Sedwick. <i>Journal of Cell Biology</i> , 2011 , 194, 162-3	7.3	
127	The in vivo contribution of motor neuron TrkB receptors to mutant SOD1 motor neuron disease. <i>Human Molecular Genetics</i> , 2011 , 20, 4116-31	5.6	17
126	FLRT2 and FLRT3 act as repulsive guidance cues for Unc5-positive neurons. <i>EMBO Journal</i> , 2011 , 30, 2920-33	10.3	101
125	Signaling from axon guidance receptors. <i>Cold Spring Harbor Perspectives in Biology</i> , 2010 , 2, a001941	10.2	175
124	Inactivation of VCP/ter94 suppresses retinal pathology caused by misfolded rhodopsin in <i>Drosophila</i> . <i>PLoS Genetics</i> , 2010 , 6, e1001075	6	58

123	Pro-survival role for Parkinson's associated gene DJ-1 revealed in trophically impaired dopaminergic neurons. <i>PLoS Biology</i> , 2010 , 8, e1000349	9.7	46
122	Cell sorting during regenerative tissue formation. <i>Cell</i> , 2010 , 143, 32-4	56.2	8
121	Topography in hippocampal mossy fiber plasticity. <i>Neuron</i> , 2010 , 65, 580-2	13.9	1
120	Neuron-astrocyte communication and synaptic plasticity. <i>Current Opinion in Neurobiology</i> , 2010 , 20, 466-73	7.3	68
119	GDNF acts as a chemoattractant to support ephrinA-induced repulsion of limb motor axons. <i>Current Biology</i> , 2010 , 20, 2150-6	6.3	54
118	Progressive postnatal motoneuron loss in mice lacking GDF-15. <i>Journal of Neuroscience</i> , 2009 , 29, 13640-8	6.6	76
117	Bidirectional modulation of synaptic functions by Eph/ephrin signaling. <i>Nature Neuroscience</i> , 2009 , 12, 15-20	25.5	331
116	Neuron-glia communication via EphA4/ephrin-A3 modulates LTP through glial glutamate transport. <i>Nature Neuroscience</i> , 2009 , 12, 1285-92	25.5	206
115	Serine phosphorylation of ephrinB2 regulates trafficking of synaptic AMPA receptors. <i>Nature Neuroscience</i> , 2008 , 11, 1035-43	25.5	62
114	The Rab5 guanylate exchange factor Rin1 regulates endocytosis of the EphA4 receptor in mature excitatory neurons. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 12539-44	11.5	54
113	The protein dendrite arborization and synapse maturation 1 (Dasm-1) is dispensable for dendrite arborization. <i>Molecular and Cellular Biology</i> , 2008 , 28, 2782-91	4.8	18
112	Absence of functional peroxisomes from mouse CNS causes dysmyelination and axon degeneration. <i>Journal of Neuroscience</i> , 2008 , 28, 4015-27	6.6	86
111	Genetic ablation of FLRT3 reveals a novel morphogenetic function for the anterior visceral endoderm in suppressing mesoderm differentiation. <i>Genes and Development</i> , 2008 , 22, 3349-62	12.6	51
110	Role for ephrinB2 in postnatal lung alveolar development and elastic matrix integrity. <i>Developmental Dynamics</i> , 2008 , 237, 2220-34	2.9	30
109	Brain IGF-1 receptors control mammalian growth and lifespan through a neuroendocrine mechanism. <i>PLoS Biology</i> , 2008 , 6, e254	9.7	204
108	Neocortical and cerebellar developmental abnormalities in conditions of selective elimination of peroxisomes from brain or from liver. <i>Journal of Neuroscience Research</i> , 2007 , 85, 58-72	4.4	67
107	Bidirectional Eph-ephrin signaling during axon guidance. <i>Trends in Cell Biology</i> , 2007 , 17, 230-8	18.3	295
106	Tyrosine phosphorylation sites in ephrinB2 are required for hippocampal long-term potentiation but not long-term depression. <i>Journal of Neuroscience</i> , 2007 , 27, 11279-88	6.6	41

105	Deletion of Shp2 in the brain leads to defective proliferation and differentiation in neural stem cells and early postnatal lethality. <i>Molecular and Cellular Biology</i> , 2007 , 27, 6706-17	4.8	109
104	RET signaling does not modulate MPTP toxicity but is required for regeneration of dopaminergic axon terminals. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 20049-54	11.5	49
103	EphA4-dependent axon guidance is mediated by the RacGAP alpha2-chimaerin. <i>Neuron</i> , 2007 , 55, 756-67	13.9	124
102	Absence of Ret signaling in mice causes progressive and late degeneration of the nigrostriatal system. <i>PLoS Biology</i> , 2007 , 5, e39	9.7	144
101	Transgenic mouse proteomics identifies new 14-3-3-associated proteins involved in cytoskeletal rearrangements and cell signaling. <i>Molecular and Cellular Proteomics</i> , 2006 , 5, 2211-27	7.6	110
100	Release of full-length EphB2 receptors from hippocampal neurons to cocultured glial cells. <i>Journal of Neuroscience</i> , 2006 , 26, 11575-81	6.6	32
99	EphB receptors and ephrin-B3 regulate axon guidance at the ventral midline of the embryonic mouse spinal cord. <i>Journal of Neuroscience</i> , 2006 , 26, 8909-14	6.6	43
98	Genetic analysis of EphA-dependent signaling mechanisms controlling topographic mapping in vivo. <i>Development (Cambridge)</i> , 2006 , 133, 4415-20	6.6	25
97	Cooperation between GDNF/Ret and ephrinA/EphA4 signals for motor-axon pathway selection in the limb. <i>Neuron</i> , 2006 , 50, 35-47	13.9	162
96	TEF-1 and C/EBPbeta are major p38alpha MAPK-regulated transcription factors in proliferating cardiomyocytes. <i>Biochemical Journal</i> , 2006 , 396, 163-72	3.8	34
95	Mig6 is a negative regulator of EGF receptor-mediated skin morphogenesis and tumor formation. <i>Nature Medicine</i> , 2006 , 12, 568-73	50.5	203
94	PDZ interaction site in ephrinB2 is required for the remodeling of lymphatic vasculature. <i>Genes and Development</i> , 2005 , 19, 397-410	12.6	357
93	Axon guidance: opposing EPHefts in the growth cone. <i>Cell</i> , 2005 , 121, 4-6	56.2	3
92	Regulation of EphA 4 kinase activity is required for a subset of axon guidance decisions suggesting a key role for receptor clustering in Eph function. <i>Neuron</i> , 2005 , 47, 515-28	13.9	100
91	Altered expression patterns of EphrinB2 and EphB2 in human umbilical vessels and congenital venous malformations. <i>Pediatric Research</i> , 2005 , 57, 537-44	3.2	31
90	Mitogen-inducible gene 6 is an endogenous inhibitor of HGF/Met-induced cell migration and neurite growth. <i>Journal of Cell Biology</i> , 2005 , 171, 337-48	7.3	62
89	The neuregulin receptor, ErbB4, is not required for normal development and adult maintenance of the substantia nigra pars compacta. <i>Journal of Neurochemistry</i> , 2004 , 91, 1302-11	6	38
88	Hippocampal plasticity requires postsynaptic ephrinBs. <i>Nature Neuroscience</i> , 2004 , 7, 33-40	25.5	215

87	Receptor tyrosine kinase ErbB4 modulates neuroblast migration and placement in the adult forebrain. <i>Nature Neuroscience</i> , 2004 , 7, 1319-28	25.5	215
86	Eph/ephrin signaling in morphogenesis, neural development and plasticity. <i>Current Opinion in Cell Biology</i> , 2004 , 16, 580-9	9	261
85	Notch activation induces apoptosis in neural progenitor cells through a p53-dependent pathway. <i>Developmental Biology</i> , 2004 , 269, 81-94	3.1	232
84	EphB-ephrinB bi-directional endocytosis terminates adhesion allowing contact mediated repulsion. <i>Nature Cell Biology</i> , 2003 , 5, 869-78	23.4	281
83	Cortical and retinal defects caused by dosage-dependent reductions in VEGF-A paracrine signaling. <i>Developmental Biology</i> , 2003 , 262, 225-41	3.1	218
82	Forebrain-specific trkB-receptor knockout mice: behaviorally more hyperactive than "depressive". <i>Biological Psychiatry</i> , 2003 , 54, 972-82	7.9	124
81	Met signaling is required for recruitment of motor neurons to PEA3-positive motor pools. <i>Neuron</i> , 2003 , 39, 767-77	13.9	59
80	Control of skeletal patterning by ephrinB1-EphB interactions. <i>Developmental Cell</i> , 2003 , 5, 217-30	10.2	199
79	Role of EphA4 and EphrinB3 in local neuronal circuits that control walking. <i>Science</i> , 2003 , 299, 1889-92	33.3	276
78	Multiple roles of ephrins in morphogenesis, neuronal networking, and brain function. <i>Genes and Development</i> , 2003 , 17, 1429-50	12.6	213
77	Eph Receptors 2003 , 421-425		
76	Immunohistochemical evidence of seizure-induced activation of trkB receptors in the mossy fiber pathway of adult mouse hippocampus. <i>Journal of Neuroscience</i> , 2002 , 22, 7502-8	6.6	79
75	Axon guidance: receptor complexes and signaling mechanisms. <i>Current Opinion in Neurobiology</i> , 2002 , 12, 250-9	7.6	69
74	Long-term monitoring of hippocampus-dependent behavior in naturalistic settings: mutant mice lacking neurotrophin receptor TrkB in the forebrain show spatial learning but impaired behavioral flexibility. <i>Hippocampus</i> , 2002 , 12, 27-38	3.5	55
73	Mechanisms and functions of Eph and ephrin signalling. <i>Nature Reviews Molecular Cell Biology</i> , 2002 , 3, 475-86	48.7	934
72	Discoidin domain receptor 2 interacts with Src and Shc following its activation by type I collagen. <i>Journal of Biological Chemistry</i> , 2002 , 277, 19206-12	5.4	99
71	Discoidin domain receptor 2 regulates fibroblast proliferation and migration through the extracellular matrix in association with transcriptional activation of matrix metalloproteinase-2. <i>Journal of Biological Chemistry</i> , 2002 , 277, 3606-13	5.4	175
70	Distinct requirements for TrkB and TrkC signaling in target innervation by sensory neurons. <i>Genes and Development</i> , 2002 , 16, 633-45	12.6	69

69	Receptor-specific regulation of phosphatidylinositol 3Kinase activation by the protein tyrosine phosphatase Shp2. <i>Molecular and Cellular Biology</i> , 2002 , 22, 4062-72	4.8	210
68	EphrinB phosphorylation and reverse signaling: regulation by Src kinases and PTP-BL phosphatase. <i>Molecular Cell</i> , 2002 , 9, 725-37	17.6	254
67	Mechanism of TrkB-mediated hippocampal long-term potentiation. <i>Neuron</i> , 2002 , 36, 121-37	13.9	353
66	Long-term monitoring of hippocampus-dependent behavior in naturalistic settings: Mutant mice lacking neurotrophin receptor TrkB in the forebrain show spatial learning but impaired behavioral flexibility 2002 , 12, 27		3
65	Excitatory Eph receptors and adhesive ephrin ligands. <i>Current Opinion in Cell Biology</i> , 2001 , 13, 196-203	9	141
64	Ephrin-B3 is the midline barrier that prevents corticospinal tract axons from recrossing, allowing for unilateral motor control. <i>Genes and Development</i> , 2001 , 15, 877-88	12.6	197
63	The collagen receptor DDR2 regulates proliferation and its elimination leads to dwarfism. <i>EMBO Reports</i> , 2001 , 2, 446-52	6.5	209
62	Kinase-dependent and kinase-independent functions of EphA4 receptors in major axon tract formation in vivo. <i>Neuron</i> , 2001 , 29, 73-84	13.9	210
61	Beta1-class integrins regulate the development of laminae and folia in the cerebral and cerebellar cortex. <i>Neuron</i> , 2001 , 31, 367-79	13.9	490
60	Kinase-independent requirement of EphB2 receptors in hippocampal synaptic plasticity. <i>Neuron</i> , 2001 , 32, 1027-40	13.9	259
59	Coupling Met to specific pathways results in distinct developmental outcomes. <i>Molecular Cell</i> , 2001 , 7, 1293-306	17.6	125
58	The cytoplasmic domain of the ligand ephrinB2 is required for vascular morphogenesis but not cranial neural crest migration. <i>Cell</i> , 2001 , 104, 57-69	56.2	239
57	Knocking the NT4 gene into the BDNF locus rescues BDNF deficient mice and reveals distinct NT4 and BDNF activities. <i>Nature Neuroscience</i> , 2000 , 3, 350-7	25.5	85
56	Eph receptors and ephrin ligands. essential mediators of vascular development. <i>Trends in Cardiovascular Medicine</i> , 2000 , 10, 183-8	6.9	120
55	Role of brain insulin receptor in control of body weight and reproduction. <i>Science</i> , 2000 , 289, 2122-5	33.3	1729
54	Essential Role of p38MAP Kinase in Placental but Not Embryonic Cardiovascular Development. <i>Molecular Cell</i> , 2000 , 6, 109-116	17.6	432
53	Shc-binding site in the TrkB receptor is not required for hippocampal long-term potentiation. <i>Neuropharmacology</i> , 2000 , 39, 717-24	5.5	30
52	Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. <i>Nature Genetics</i> , 1999 , 23, 99-103	36.3	1430

51	Tyro-3 family receptors are essential regulators of mammalian spermatogenesis. <i>Nature</i> , 1999 , 398, 723-8	50.4	379
50	Hepatocyte growth factor, a versatile signal for developing neurons. <i>Nature Neuroscience</i> , 1999 , 2, 213-7	25.5	203
49	Bidirectional signals establish boundaries. <i>Current Biology</i> , 1999 , 9, R691-4	6.3	32
48	EphrinB ligands recruit GRIP family PDZ adaptor proteins into raft membrane microdomains. <i>Neuron</i> , 1999 , 22, 511-24	13.9	308
47	Essential role for TrkB receptors in hippocampus-mediated learning. <i>Neuron</i> , 1999 , 24, 401-14	13.9	666
46	Roles of ephrinB ligands and EphB receptors in cardiovascular development: demarcation of arterial/venous domains, vascular morphogenesis, and sprouting angiogenesis. <i>Genes and Development</i> , 1999 , 13, 295-306	12.6	799
45	Ephrin-B3, a ligand for the receptor EphB3, expressed at the midline of the developing neural tube. <i>Oncogene</i> , 1998 , 16, 471-80	9.2	64
44	Signaling by Eph receptors and their ephrin ligands. <i>Current Opinion in Neurobiology</i> , 1998 , 8, 375-82	7.6	127
43	Multiple roles for hepatocyte growth factor in sympathetic neuron development. <i>Neuron</i> , 1998 , 20, 835-46	15.9	130
42	Point mutation in trkB causes loss of NT4-dependent neurons without major effects on diverse BDNF responses. <i>Neuron</i> , 1998 , 21, 335-45	13.9	169
41	TrkB and neurotrophin-4 are important for development and maintenance of sympathetic preganglionic neurons innervating the adrenal medulla. <i>Journal of Neuroscience</i> , 1998 , 18, 7272-84	6.6	41
40	Met receptor signaling is required for sensory nerve development and HGF promotes axonal growth and survival of sensory neurons. <i>Genes and Development</i> , 1997 , 11, 3341-50	12.6	196
39	Tyrosine phosphorylation of transmembrane ligands for Eph receptors. <i>Science</i> , 1997 , 275, 1640-3	33.3	351
38	Survival of inner ear sensory neurons in trk mutant mice. <i>Mechanisms of Development</i> , 1997 , 64, 77-85	1.7	25
37	A role for the Ras signalling pathway in synaptic transmission and long-term memory. <i>Nature</i> , 1997 , 390, 281-6	50.4	419
36	The Eph receptor family: axonal guidance by contact repulsion. <i>Trends in Genetics</i> , 1997 , 13, 354-9	8.5	162
35	The N-terminal globular domain of Eph receptors is sufficient for ligand binding and receptor signaling. <i>EMBO Journal</i> , 1997 , 16, 3889-97	13	64
34	Reduced acetylcholinesterase (AChE) activity in adrenal medulla and loss of sympathetic preganglionic neurons in TrkA-deficient, but not TrkB-deficient, mice. <i>Journal of Neuroscience</i> , 1997 , 17, 891-903	6.6	27

33	Similarities and differences in the way transmembrane-type ligands interact with the Elk subclass of Eph receptors. <i>Molecular and Cellular Neurosciences</i> , 1996 , 8, 199-209	4.8	59
32	Nuk controls pathfinding of commissural axons in the mammalian central nervous system. <i>Cell</i> , 1996 , 86, 35-46	56.2	468
31	Uncoupling of Grb2 from the Met receptor in vivo reveals complex roles in muscle development. <i>Cell</i> , 1996 , 87, 531-42	56.2	275
30	Sek4 and Nuk receptors cooperate in guidance of commissural axons and in palate formation.. <i>EMBO Journal</i> , 1996 , 15, 6035-6049	13	260
29	TrkB and TrkC neurotrophin receptors cooperate in promoting survival of hippocampal and cerebellar granule neurons. <i>Genes and Development</i> , 1996 , 10, 2849-58	12.6	196
28	Aberrant neural and cardiac development in mice lacking the ErbB4 neuregulin receptor. <i>Nature</i> , 1995 , 378, 390-4	50.4	928
27	The neurotrophin receptors TrkA and TrkB are inhibitory for neurite outgrowth. <i>European Journal of Neuroscience</i> , 1995 , 7, 1424-8	3.5	15
26	Membrane-bound LERK2 ligand can signal through three different Eph-related receptor tyrosine kinases.. <i>EMBO Journal</i> , 1995 , 14, 3116-3126	13	107
25	Developmental changes in NT3 signalling via TrkA and TrkB in embryonic neurons.. <i>EMBO Journal</i> , 1995 , 14, 4482-4489	13	122
24	Telling axons where to grow: a role for Eph receptor tyrosine kinases in guidance. <i>Molecular and Cellular Neurosciences</i> , 1995 , 6, 487-95	4.8	74
23	ELF-2, a new member of the Eph ligand family, is segmentally expressed in mouse embryos in the region of the hindbrain and newly forming somites. <i>Molecular and Cellular Biology</i> , 1995 , 15, 4921-9	4.8	149
22	Role of neurotrophins in mouse neuronal development. <i>FASEB Journal</i> , 1994 , 8, 738-44	0.9	221
21	Severe sensory and sympathetic neuropathies in mice carrying a disrupted Trk/NGF receptor gene. <i>Nature</i> , 1994 , 368, 246-9	50.4	859
20	Disruption of the neurotrophin-3 receptor gene trkC eliminates la muscle afferents and results in abnormal movements. <i>Nature</i> , 1994 , 368, 249-51	50.4	560
19	High-affinity nerve growth factor receptor (Trk) immunoreactivity is localized in cholinergic neurons of the basal forebrain and striatum in the adult rat brain. <i>Brain Research</i> , 1993 , 612, 330-5	3.7	135
18	Targeted disruption of the trkB neurotrophin receptor gene results in nervous system lesions and neonatal death. <i>Cell</i> , 1993 , 75, 113-122	56.2	538
17	Similarities and differences in the way neurotrophins interact with the Trk receptors in neuronal and nonneuronal cells. <i>Neuron</i> , 1993 , 10, 137-49	13.9	497
16	Induction of noncatalytic TrkB neurotrophin receptors during axonal sprouting in the adult hippocampus. <i>Journal of Neuroscience</i> , 1993 , 13, 4001-14	6.6	122

15	The trkB tyrosine protein kinase is a receptor for neurotrophin-4. <i>Neuron</i> , 1992 , 8, 947-56	13.9	288
14	The trk family of tyrosine protein kinase receptors. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 1991 , 1072, 115-27	11.2	58
13	The trk tyrosine protein kinase mediates the mitogenic properties of nerve growth factor and neurotrophin-3. <i>Cell</i> , 1991 , 66, 173-83	56.2	495
12	The trk proto-oncogene encodes a receptor for nerve growth factor. <i>Cell</i> , 1991 , 65, 189-97	56.2	1266
11	trkC, a new member of the trk family of tyrosine protein kinases, is a receptor for neurotrophin-3. <i>Cell</i> , 1991 , 66, 967-79	56.2	959
10	The trkB tyrosine protein kinase is a receptor for brain-derived neurotrophic factor and neurotrophin-3. <i>Cell</i> , 1991 , 66, 395-403	56.2	813
9	Human trk oncogenes activated by point mutation, in-frame deletion, and duplication of the tyrosine kinase domain. <i>Molecular and Cellular Biology</i> , 1990 , 10, 4202-10	4.8	76
8	The trkB tyrosine protein kinase gene codes for a second neurogenic receptor that lacks the catalytic kinase domain. <i>Cell</i> , 1990 , 61, 647-56	56.2	666
7	trkB, a novel tyrosine protein kinase receptor expressed during mouse neural development.. <i>EMBO Journal</i> , 1989 , 8, 3701-3709	13	433
6	Highly glycosylated PDGF-like molecule secreted by simian sarcoma virus-transformed cells. <i>Virology</i> , 1988 , 164, 403-10	3.6	2
5	The 89,000-Mr murine cytomegalovirus immediate-early protein stimulates c-fos expression and cellular DNA synthesis. <i>Journal of Virology</i> , 1988 , 62, 3341-7	6.6	16
4	Diurnal variation of several blood parameters in the owl monkey, <i>Aotus trivirgatus griseimembra</i> . <i>Folia Primatologica</i> , 1985 , 45, 195-203	1.2	9
3	Amyloid-like aggregates cause lysosomal defects in neurons via gain-of-function toxicity		4
2	Central amygdala circuits modulate food consumption through a positive valence mechanism		1
1	Natural loss of function of ephrin-B3 shapes spinal flight circuitry in birds		1