

Grozdana Bogdanic

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3884692/publications.pdf>

Version: 2024-02-01

48
papers

487
citations

840119

11
h-index

752256

20
g-index

50
all docs

50
docs citations

50
times ranked

272
citing authors

#	ARTICLE	IF	CITATIONS
1	Vapor-liquid and liquid-liquid equilibria in the water+poly(propylene glycol) system. Journal of Molecular Liquids, 2021, 337, 116336.	2.3	4
2	Vapour-liquid equilibria in water+poly(ethylene glycol) systems: New experiments and cumulative thermodynamic processing of all data. Journal of Chemical Thermodynamics, 2020, 140, 105901.	1.0	4
3	Simple Apparatus for the Measurement of Total Pressure of Polymer+Solvent Mixtures. Chemical Engineering and Technology, 2017, 40, 991-996.	0.9	3
4	An improved apparatus for vapour+liquid equilibria measurement in polymer+ solvent systems at higher temperatures: A study of the water+ poly(ethylene glycol) system. Fluid Phase Equilibria, 2017, 454, 111-115.	1.4	3
5	Ebulliometric measurement of total pressure in the binary polystyrene+butan-2-one system. Fluid Phase Equilibria, 2016, 424, 41-43.	1.4	3
6	Vapour+Liquid Equilibria in the Polystyrene + Toluene System at Higher Concentrations of Solvent. Chemical and Biochemical Engineering Quarterly, 2015, 29, 1-4.	0.5	6
7	Vapour+liquid equilibria in the polymer+solvent system containing lower concentrations of solute at normal or reduced pressures. Fluid Phase Equilibria, 2013, 358, 301-303.	1.4	3
8	Vapour+liquid equilibria in binary and ternary systems composed of 2,3-dimethylbutane, diisopropyl ether, and 3-methyl-2-butanone at 313.15, 323.15 and 313.15K. Fluid Phase Equilibria, 2013, 344, 59-64.	1.4	13
9	Experiments and Modelling of Liquid+liquid Equilibria in the Mineral Oil + N,N-dimethylformamide System. Procedia Engineering, 2012, 42, 721-725.	1.2	0
10	Vapour+liquid and chemical equilibria in the ethyl ethanoate+ethanol+propyl ethanoate+propanol system accompanied with transesterification reaction. Fluid Phase Equilibria, 2012, 328, 61-68.	1.4	19
11	Vapor+Liquid Equilibrium in Diluted Polymer + Solvent Systems. Journal of Chemical & Engineering Data, 2011, 56, 1080-1083.	1.0	9
12	Circulation micro-ebulliometer for determination of pressure above mixtures containing solvent and non-volatile component. Fluid Phase Equilibria, 2010, 297, 142-148.	1.4	14
13	Studies on the influence of long chain acrylic esters polymers with polar monomers as crude oil flow improver additives. Fuel, 2008, 87, 2943-2950.	3.4	102
14	Copolymerization of N+tert+Butylacrylamide with Ethylene Glycol Dimethacrylate. Journal of Macromolecular Science - Pure and Applied Chemistry, 2007, 44, 721-725.	1.2	0
15	Flow improver additives for gas condensate. Fuel, 2007, 86, 1409-1416.	3.4	9
16	Synthesis of N+Acryl+EN,N+di+tert+Butylurea and Copolymerization with Ethylene Glycol Dimethacrylate. Journal of Macromolecular Science - Pure and Applied Chemistry, 2006, 43, 879-887.	1.2	2
17	Group contribution methods for estimating the properties of polymer systems. Hemijska Industrija, 2006, 60, 287-305.	0.3	8
18	Structural Differences Between Copolymers of Acryl+and Methacryl+dicyclohexylurea with Ethylene Glycol Dimethacrylate and their Thermal Degradation Products. Journal of Macromolecular Science - Pure and Applied Chemistry, 2005, 42, 1621-1626.	1.2	2

#	ARTICLE	IF	CITATIONS
19	Free-Radical Initiated Polymerization of N-methacryl-N- ϵ -diisopropylurea with Styrene. Journal of Macromolecular Science - Pure and Applied Chemistry, 2005, 42, 535-542.	1.2	0
20	Nanoporous Crosslinked Copolymers Prepared by Thermal Degradation of Poly(Methacryl-N- ϵ -diisopropylurea- ϵ -ethylene Glycol Dimethacrylate). Journal of Macromolecular Science - Pure and Applied Chemistry, 2004, 41, 1087-1094.	1.2	3
21	Preparation of Nanoporous Crosslinked Poly(Methacryl-N- ϵ -cyclohexylamide- ϵ -ethylene Glycol) Tj ETQq1 1 0.784314 rgBT /Overl	1.2	4
22	Synthesis of Nanoporous Crosslinked Poly(Acryl-N- ϵ -cyclohexyl Amide- ϵ -ethylene Glycol Dimethacrylate) by Thermal Degradation of Poly(Acryl-N- ϵ -dicyclohexylurea- ϵ -ethylene Glycol Dimethacrylate). Journal of Macromolecular Science - Pure and Applied Chemistry, 2003, 40, 747-754.	1.2	6
23	Pearson-type I Distribution Function for Polydisperse Polymer Systems. Molar Mass Distribution. Journal of Chemical Information and Computer Sciences, 2003, 43, 880-884.	2.8	1
24	Separation of Cyclohexylisocyanate from the Crosslinked Copolymers of N-Acryl-dicyclohexylurea with Ethylene Glycol Dimethacrylate or Divinyl Benzene. Journal of Macromolecular Science - Pure and Applied Chemistry, 2003, 40, 81-85.	1.2	6
25	POLYMERIZATION OF N(p-PHENOXY-PHENYL)ACRYLAMIDE AND COPOLYMERS WITH STYRENE. Journal of Macromolecular Science - Pure and Applied Chemistry, 2001, 38, 1075-1086.	1.2	2
26	COPOLYMERIZATION OF N-ACRYL-N- ϵ -DICYCLOHEXYLUREA AND N-METHACRYL-N- ϵ -DICYCLOHEXYLUREA WITH STYRENE. Journal of Macromolecular Science - Pure and Applied Chemistry, 2001, 38, 839-850.	1.2	3
27	The FV-UNIQUAC segmental interaction model for liquid-liquid equilibrium calculations for polymer solutions. Fluid Phase Equilibria, 2001, 191, 49-57.	1.4	5
28	FREE-RADICAL-INITIATED COPOLYMERIZATION OF 2-CHLOROSTYRENE, 4-CHLOROSTYRENE, AND 2,6-DICHLOROSTYRENE WITH MALEIC ANHYDRIDE. Journal of Macromolecular Science - Pure and Applied Chemistry, 2001, 38, 253-261.	1.2	2
29	FREE RADICAL-INITIATED COPOLYMERIZATION OF 2,6-DICHLOROSTYRENE WITH MALEIMIDE, N-METHYLMALEIMIDE, AND N-PHENYLMALEIMIDE. Journal of Macromolecular Science - Pure and Applied Chemistry, 2000, 37, 513-524.	1.2	12
30	A segmental interaction model for liquid-liquid equilibrium calculations for polymer solutions. Fluid Phase Equilibria, 2000, 173, 241-252.	1.4	35
31	Phase Behavior and Miscibility in Binary Blends Containing Polymers and Copolymers of Styrene, of 2,6-Dimethyl-1,4-Phenylene Oxide, and of Their Derivatives. Journal of Physical and Chemical Reference Data, 1999, 28, 851-868.	1.9	17
32	A segmental interaction model for liquid-liquid equilibria correlation and prediction. Polymer Bulletin, 1998, 40, 117-123.	1.7	7
33	Phase behaviour in blends of poly[styrene-co-ortho(para)-bromostyrene] and phenylsulfonylated poly(2,6-dimethyl-1,4-phenylene oxide) copolymers. Polymer, 1998, 39, 2847-2850.	1.8	3
34	Estimation of the segmental interaction parameters of polymer blends based on styrene and 2,6-dimethyl-1,4-phenylene oxide derivatives. Fluid Phase Equilibria, 1997, 139, 277-294.	1.4	3
35	Miscibility-immiscibility behaviour in blends of phenylsulfonylated poly(2,6-dimethyl-1,4-phenylene) Tj ETQq1 1 0.784314 rgBT /Overl	1.2	3
36	Copolymer blends of phenylsulfonylated poly(2,6-dimethyl-1,4-phenylene oxide) and poly(p-fluorostyrene-co-p(o)-chlorostyrene). Thermochimica Acta, 1996, 275, 259-268.	1.2	4

#	ARTICLE	IF	CITATIONS
37	Miscibility and phase separation study of blends of random copolymers of poly[ortho(para)-fluorostyrene-co-ortho(para)-bromostyrene] with phenylsulphonylated poly(2,6-dimethyl-1,4-phenylene oxide) copolymers by thermal methods. <i>Thermochimica Acta</i> , 1996, 285, 141-154.	1.2	4
38	Phase behaviour in copolymer blends of phenylsulphonylated poly(2,6-dimethyl-1,4-phenylene oxide) and poly(o-fluorostyrene-co-p(o)-chlorostyrene). <i>Thermochimica Acta</i> , 1995, 264, 125-135.	1.2	6
39	Revision of the Group-Contribution-Flory Equation of State for Phase Equilibria Calculations in Mixtures with Polymers. 2. Prediction of Liquid-Liquid Equilibria for Polymer Solutions. <i>Industrial & Engineering Chemistry Research</i> , 1995, 34, 1835-1841.	1.8	17
40	Prediction of Vapor-Liquid Equilibria for Mixtures with Copolymers. <i>Industrial & Engineering Chemistry Research</i> , 1995, 34, 324-331.	1.8	15
41	DSC study of the miscibility of PPO and sulphonylated PPO in blends with alternating copolymers of β -substituted styrene derivatives with maleic anhydride and with N-substituted maleimides. <i>Thermochimica Acta</i> , 1994, 233, 75-86.	1.2	3
42	Phase behavior in copolymer blends of poly (p-chlorostyrene-co-o-chlorostyrene) and phenylsulfonylated poly (2,6-dimethyl-1,4-phenylene oxide). <i>Journal of Polymer Science, Part B: Polymer Physics</i> , 1994, 32, 1079-1087.	2.4	10
43	Miscibility in blends of sulfonylated poly(2,6-dimethyl-1,4-phenylene oxide) and poly(p-bromostyrene-co-o-bromostyrene). <i>Journal of Applied Polymer Science</i> , 1994, 52, 1499-1503.	1.3	8
44	Miscibility in blends of phenylsulfonylated poly(2,6-dimethyl-1,4-phenylene oxide) and poly(p-fluorostyrene-co-o-fluorostyrene). <i>Polymer</i> , 1994, 35, 3055-3059.	1.8	6
45	Revision of the Group-Contribution Flory Equation of State for Phase Equilibria Calculations in Mixtures with Polymers. 1. Prediction of Vapor-Liquid Equilibria for Polymer Solutions. <i>Industrial & Engineering Chemistry Research</i> , 1994, 33, 1331-1340.	1.8	70
46	Miscibility in blends of sulfonylated poly(2,6-dimethyl-1,4-phenylene oxide) (SPPO) with homopolymers of halogen-substituted styrene derivatives. <i>Polymer</i> , 1993, 34, 1449-1453.	1.8	15
47	Miscibility behaviour of sulfonylated poly(2,6-dimethyl-1,4-phenylene oxide) copolymers in the blends with poly(styrene-co-maleic anhydride) and with poly(β -methylstyrene-co-maleic anhydride). <i>Polymer Bulletin</i> , 1992, 28, 473-479.	1.7	6
48	Investigation of the decomposition of copolymers of styrene and maleic anhydride using thermogravimetric analysis. <i>Thermochimica Acta</i> , 1990, 171, 39-47.	1.2	7