

Dariusz Kotlega

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3888035/publications.pdf>

Version: 2024-02-01

36
papers

359
citations

840585

11
h-index

887953

17
g-index

38
all docs

38
docs citations

38
times ranked

535
citing authors

#	ARTICLE	IF	CITATIONS
1	High Levels of Thromboxane (TX) Are Associated with the Sex-Dependent Non-Dipping Phenomenon in Ischemic Stroke Patients. <i>Journal of Clinical Medicine</i> , 2022, 11, 2652.	1.0	1
2	Could Lipoxins Represent a New Standard in Ischemic Stroke Treatment?. <i>International Journal of Molecular Sciences</i> , 2021, 22, 4207.	1.8	13
3	Prostaglandin E2, 9S-, 13S-HODE and resolvin D1 are strongly associated with the post-stroke cognitive impairment. <i>Prostaglandins and Other Lipid Mediators</i> , 2021, 156, 106576.	1.0	10
4	Free Fatty Acids Are Associated with the Cognitive Functions in Stroke Survivors. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 6500.	1.2	13
5	The Role of Thromboxane in the Course and Treatment of Ischemic Stroke: Review. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11644.	1.8	26
6	The Role of Resolvins: EPA and DHA Derivatives Can Be Useful in the Prevention and Treatment of Ischemic Stroke. <i>International Journal of Molecular Sciences</i> , 2020, 21, 7628.	1.8	24
7	The Association of Free Fatty Acids and Eicosanoids with the Severity of Depressive Symptoms in Stroke Patients. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5220.	1.8	9
8	Lipoxins, RevD1 and 9, 13 HODE as the most important derivatives after an early incident of ischemic stroke. <i>Scientific Reports</i> , 2020, 10, 12849.	1.6	31
9	Free Fatty Acids and Their Inflammatory Derivatives Affect BDNF in Stroke Patients. <i>Mediators of Inflammation</i> , 2020, 2020, 1-12.	1.4	11
10	Multiple Cryotherapy Attenuates Oxi-Inflammatory Response Following Skeletal Muscle Injury. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7855.	1.2	11
11	NEUROPSYCHOLOGICAL CHARACTERISTIC OF COGNITIVE DISORDERS IN PATIENT WITH CLIPPERS SYNDROME. <i>Acta Neuropsychologica</i> , 2020, 18, 393-401.	0.3	0
12	Detection of stroke risk factors over the decade in the polish population of ischemic stroke patients. <i>Postepy Psychiatrii i Neurologii</i> , 2019, 28, 83-87.	0.2	3
13	The early outcome of thrombolytic treatment in ischaemic stroke patients: a retrospective analysis. <i>Studia Medyczne</i> , 2019, 35, 16-22.	0.0	0
14	Association between the Urinary Bladder Volume and the Incidence of "De Novo" Overactive Bladder in Patients with Stress Urinary Incontinence Subjected to Sling Surgeries or Burch Procedure. <i>BioMed Research International</i> , 2019, 2019, 1-6.	0.9	2
15	Beneficial effects of pre-stroke statins use in cardioembolic stroke patients with atrial fibrillation: a hospital-based retrospective analysis. <i>Archives of Medical Science</i> , 2019, 15, 385-392.	0.4	11
16	Elevated Inflammatory Parameter Levels Negatively Impact Populations of Circulating Stem Cells (CD133+), Early Endothelial Progenitor Cells (CD133+/VEGFR2+), and Fibroblast Growth Factor in Stroke Patients. <i>Current Neurovascular Research</i> , 2019, 16, 19-26.	0.4	21
17	STROKE MIMICS: A PSYCHOGENIC STROKE PATIENT TREATED WITH ALTEPLASE. <i>Acta Neuropsychologica</i> , 2019, 17, 97-102.	0.3	0
18	Plastic surgery of nail folds combined with Arkada™'s method in patient with ingrown nail " case report. <i>Pomeranian Journal of Life Sciences</i> , 2019, 65, 42-44.	0.1	1

#	ARTICLE	IF	CITATIONS
19	The use of botulinum toxin for the treatment of patients with overactive bladder. Pomeranian Journal of Life Sciences, 2019, 64, 9-13.	0.1	0
20	Differences between Subjective and Objective Assessment of Speech Deficiency in Parkinson Disease. Journal of Voice, 2018, 32, 715-722.	0.6	11
21	Effects of Angiotensin-Converting Enzyme Inhibition on Circulating Endothelial Progenitor Cells in Patients with Acute Ischemic Stroke. Stem Cells International, 2018, 2018, 1-10.	1.2	5
22	Statins Therapy is Associated with Increased Populations of Early Endothelial Progenitor (CD133+/VEGFR2+) and Endothelial (CD34-/CD133- /VEGFR2+) Cells in Patients with Acute Ischemic Stroke. Current Neurovascular Research, 2018, 15, 120-128.	0.4	11
23	The role of brain-derived neurotrophic factor and its single nucleotide polymorphisms in stroke patients. Neurologia I Neurochirurgia Polska, 2017, 51, 240-246.	0.6	39
24	NEUROPSYCHOLOGICAL SYMPTOMS AND THERAPY IN PATIENT WITH A BILATERAL THALAMIC ISCHEMIC STROKE DUE TO THE ARTERY OF PERCHERON TERRITORY INFARCT. Acta Neuropsychologica, 2017, 15, 0-0.	0.3	1
25	Delayed neurological sequelae in carbon monoxide poisoning – a case report and present state of knowledge. Pomeranian Journal of Life Sciences, 2017, 63, .	0.1	0
26	Risk Factors of Fatigue in Idiopathic Parkinson’s Disease in a Polish Population. Parkinson's Disease, 2016, 2016, 1-8.	0.6	6
27	Early Emergency Medical Service Calls for Stroke: Was the Long-Term Education Program Based on the Experience of West Pomerania Successful?. Journal of Stroke and Cerebrovascular Diseases, 2016, 25, 254-258.	0.7	3
28	The emotional stress and risk of ischemic stroke. Neurologia I Neurochirurgia Polska, 2016, 50, 265-270.	0.6	34
29	Simultaneous acute shoulder arthritis and multiple mononeuropathy in a newly diagnosed type 2 diabetes patient – First case report. Neurologia I Neurochirurgia Polska, 2016, 50, 474-480.	0.6	0
30	Association between selected gene polymorphisms and statin metabolism, risk of ischemic stroke and cardiovascular disorders. Postepy Higieny I Medycyny Doswiadczalnej, 2016, 70, 435-447.	0.1	12
31	Prolonged effect of temporary rituximab therapy in neuromyelitis optica – A case study. Postepy Psychiatrii I Neurologii, 2015, 24, 239-243.	0.2	1
32	Potential role of statins in the intracerebral hemorrhage and subarachnoid hemorrhage. Neurologia I Neurochirurgia Polska, 2015, 49, 322-328.	0.6	12
33	Atrial fibrillation and stroke – Coexistence and attitude to preventive therapy on the basis of Szczecin and Szczecin region patients. Neurologia I Neurochirurgia Polska, 2014, 48, 410-415.	0.6	3
34	Anti-oxLDL antibodies are clinically insignificant for stroke patients. Neurological Research, 2014, 36, 86-91.	0.6	10
35	Association between CRP gene polymorphism 717A/G, C-reactive protein and neurological deficit in ischemic stroke. Journal of Clinical Neuroscience, 2014, 21, 574-577.	0.8	7
36	Risk factors of stroke and rs2794521 CRP gene polymorphism among stroke patients in West Pomerania province of Poland. Neurologia I Neurochirurgia Polska, 2014, 48, 30-34.	0.6	12