

Yanjin Hu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3891727/publications.pdf>

Version: 2024-02-01

19
papers

322
citations

1051969

10
h-index

1051228

16
g-index

20
all docs

20
docs citations

20
times ranked

634
citing authors

#	ARTICLE	IF	CITATIONS
1	The Changes of Lipidomic Profiles Reveal Therapeutic Effects of Exenatide in Patients With Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2022, 13, 677202.	1.5	11
2	The Prevalence of Euthyroid Hypertriiodothyroninemia in Newly Diagnosed Multiple Myeloma and its Clinical Characteristics. <i>Endocrine Practice</i> , 2021, 27, 236-240.	1.1	5
3	Primary Hyperparathyroidism in Pregnancy: Insights From a Case of a 28-Year-Old Woman With Miscarriages and Hyperemesis Gravidarum. <i>Annals of Laboratory Medicine</i> , 2021, 41, 336-338.	1.2	0
4	Relation of kidney function and homocysteine in patients with hypothyroidism. <i>Endocrine Connections</i> , 2021, 10, 502-510.	0.8	1
5	The association between albuminuria and thyroid antibodies in newly diagnosed type 2 diabetes mellitus patients with Hashimoto's thyroiditis and euthyroidism. <i>BMC Endocrine Disorders</i> , 2020, 20, 172.	0.9	2
6	The effect of exenatide on fasting bile acids in newly diagnosed type 2 diabetes mellitus patients, a pilot study. <i>BMC Pharmacology & Toxicology</i> , 2020, 21, 44.	1.0	0
7	The Relationship Between the Impairment of Endothelial Function and Thyroid Antibodies in Hashimoto's Thyroiditis Patients with Euthyroidism. <i>Hormone and Metabolic Research</i> , 2020, 52, 642-646.	0.7	3
8	Homocysteine Levels are Associated with Endothelial Function in Newly Diagnosed Type 2 Diabetes Mellitus Patients. <i>Metabolic Syndrome and Related Disorders</i> , 2019, 17, 323-327.	0.5	15
9	The Effects of Exenatide and Metformin on Endothelial Function in Newly Diagnosed Type 2 Diabetes Mellitus Patients: A Case-Control Study. <i>Diabetes Therapy</i> , 2018, 9, 1295-1305.	1.2	14
10	Comparison of Exenatide and Metformin Monotherapy in Overweight/Obese Patients with Newly Diagnosed Type 2 Diabetes. <i>International Journal of Endocrinology</i> , 2017, 2017, 1-6.	0.6	8
11	PPAR α Agonist Fenofibrate Reduced the Secreting Load of β -Cells in Hypertriglyceridemia Patients with Normal Glucose Tolerance. <i>PPAR Research</i> , 2016, 2016, 1-7.	1.1	5
12	Levothyroxine treatment restored the decreased circulating fibroblast growth factor 21 levels in patients with hypothyroidism. <i>European Journal of Internal Medicine</i> , 2016, 31, 94-98.	1.0	11
13	Exenatide treatment increases serum irisin levels in patients with obesity and newly diagnosed type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1555-1559.	1.2	34
14	Potential harmful correlation between homocysteine and low-density lipoprotein cholesterol in patients with hypothyroidism. <i>Medicine (United States)</i> , 2016, 95, e4291.	0.4	17
15	Comparison of β -cell dysfunction and insulin resistance correlating obesity with type 2 diabetes: A cross-sectional study. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 898-902.	1.2	5
16	High prevalence of vitamin D deficiency in urban health checkup population. <i>Clinical Nutrition</i> , 2016, 35, 859-863.	2.3	41
17	Clinical Study of Serum Homocysteine and Non-Alcoholic Fatty Liver Disease in Euglycemic Patients. <i>Medical Science Monitor</i> , 2016, 22, 4146-4151.	0.5	10
18	The role of fibroblast growth factor 21 in the pathogenesis of non-alcoholic fatty liver disease and implications for therapy. <i>Metabolism: Clinical and Experimental</i> , 2015, 64, 380-390.	1.5	96

#	ARTICLE	IF	CITATIONS
19	Novel Clinical Evidence of an Association between Homocysteine and Insulin Resistance in Patients with Hypothyroidism or Subclinical Hypothyroidism. PLoS ONE, 2015, 10, e0125922.	1.1	44