Hyo Jung Seo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3895808/publications.pdf

Version: 2024-02-01

| 19 papers | 377 citations | 933264 10 h-index | 17 g-index |
|--------------|------------------|-------------------------|----------------|
| 19 | 19 | 19 | 758 |
| all docs | docs citations | times ranked | citing authors |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Gadoxetate Disodium-Enhanced Magnetic Resonance Imaging Versus Contrast-Enhanced 18F-Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography for the Detection of Colorectal Liver Metastases. Investigative Radiology, 2011, 46, 548-555. | 3.5 | 83 |
| 2 | Prognostic value of 18Fâ€FDG PET for hepatocellular carcinoma patients treated with sorafenib. Liver International, 2011, 31, 1144-1149. | 1.9 | 56 |
| 3 | Usefulness of Integrated PET/MRI in Head and Neck Cancer: A Preliminary Study. Nuclear Medicine and Molecular Imaging, 2014, 48, 98-105. | 0.6 | 34 |
| 4 | Rapid Hepatobiliary Excretion of Micelle-Encapsulated/Radiolabeled Upconverting Nanoparticles as an Integrated Form. Scientific Reports, 2015, 5, 15685. | 1.6 | 34 |
| 5 | Positron Emission Tomography/Magnetic Resonance Imaging Evaluation of Lung Cancer. Journal of Thoracic Imaging, 2014, 29, 4-16. | 0.8 | 33 |
| 6 | 18F-FDG PET/CT in hepatocellular carcinoma. Nuclear Medicine Communications, 2015, 36, 226-233. | 0.5 | 23 |
| 7 | Correlation Between 18F-Fluorodeoxyglucose Uptake and Epidermal Growth Factor Receptor Mutations in Advanced Lung Cancer. Nuclear Medicine and Molecular Imaging, 2012, 46, 169-175. | 0.6 | 21 |
| 8 | Ratio of Mediastinal Lymph Node SUV to Primary Tumor SUV in 18F-FDG PET/CT for Nodal Staging in Non-Small-Cell Lung Cancer. Nuclear Medicine and Molecular Imaging, 2017, 51, 140-146. | 0.6 | 18 |
| 9 | Clinical Performance of Whole-Body 18F-FDG PET/Dixon-VIBE, T1-Weighted, and T2-Weighted MRI Protocol in Colorectal Cancer. Clinical Nuclear Medicine, 2015, 40, e392-e398. | 0.7 | 17 |
| 10 | Evaluation of Bone Metastasis from Hepatocellular Carcinoma Using 18F-FDG PET/CT and 99mTc-HDP Bone Scintigraphy: Characteristics of Soft Tissue Formation. Nuclear Medicine and Molecular Imaging, 2011, 45, 203-211. | 0.6 | 13 |
| 11 | Using 18F-FDG PET/CT to Detect an Occult Mesenchymal Tumor Causing Oncogenic Osteomalacia. Nuclear Medicine and Molecular Imaging, 2011, 45, 233-237. | 0.6 | 10 |
| 12 | Usefulness of Additional SPECT/CT Identifying Lymphatico-renal Shunt in a Patient with Chyluria. Nuclear Medicine and Molecular Imaging, 2015, 49, 61-64. | 0.6 | 8 |
| 13 | A Hepatoid Adenocarcinoma of the Stomach Evaluated With 18F-FDG PET/CT. Clinical Nuclear Medicine, 2014, 39, 442-445. | 0.7 | 7 |
| 14 | Usefulness of 131I-SPECT/CT and 18F-FDG PET/CT in Evaluating Successful 131I and Retinoic Acid Combined Therapy in a Patient with Metastatic Struma Ovarii. Nuclear Medicine and Molecular Imaging, 2015, 49, 52-56. | 0.6 | 7 |
| 15 | Hemodynamic Significance of Internal Carotid or Middle Cerebral Artery Stenosis Detected on Magnetic Resonance Angiography. Yonsei Medical Journal, 2015, 56, 1686. | 0.9 | 5 |
| 16 | Usefulness of 18F-FDG PET/CT to Detect Metastatic Mucinous Adenocarcinoma Within an Inguinal Hernia. Nuclear Medicine and Molecular Imaging, 2016, 50, 85-89. | 0.6 | 4 |
| 17 | Prognostic significance of interim ¹⁸ Fâ€fluorodeoxyglucose positron emission tomography–computed tomography volumetric parameters in metastatic or recurrent gastric cancer. Asia-Pacific Journal of Clinical Oncology, 2018, 14, e302-e309. | 0.7 | 4 |
| 18 | High Serum Levels of Thyroid-Stimulating Hormone and Sustained Weight Gain in Patients with Thyroid Cancer Undergoing Radioiodine Therapy. International Journal of Thyroidology, 2016, 9, 19. | 0.1 | 0 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Clinical Use of Radiopharmaceuticals in Boron Neutron Capture Therapy. Nuclear Medicine and Molecular Imaging, $0, 1$. | 0.6 | O |