
Hyo Jung Seo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3895808/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Prognostic significance of interim ¹⁸ Fâ€fluorodeoxyglucose positron emission tomography–computed tomography volumetric parameters in metastatic or recurrent gastric cancer. Asia-Pacific Journal of Clinical Oncology, 2018, 14, e302-e309.	0.7	4
2	Ratio of Mediastinal Lymph Node SUV to Primary Tumor SUV in 18F-FDG PET/CT for Nodal Staging in Non-Small-Cell Lung Cancer. Nuclear Medicine and Molecular Imaging, 2017, 51, 140-146.	0.6	18
3	High Serum Levels of Thyroid-Stimulating Hormone and Sustained Weight Gain in Patients with Thyroid Cancer Undergoing Radioiodine Therapy. International Journal of Thyroidology, 2016, 9, 19.	0.1	0
4	Usefulness of 18F-FDG PET/CT to Detect Metastatic Mucinous Adenocarcinoma Within an Inguinal Hernia. Nuclear Medicine and Molecular Imaging, 2016, 50, 85-89.	0.6	4
5	Rapid Hepatobiliary Excretion of Micelle-Encapsulated/Radiolabeled Upconverting Nanoparticles as an Integrated Form. Scientific Reports, 2015, 5, 15685.	1.6	34
6	Clinical Performance of Whole-Body 18F-FDG PET/Dixon-VIBE, T1-Weighted, and T2-Weighted MRI Protocol in Colorectal Cancer. Clinical Nuclear Medicine, 2015, 40, e392-e398.	0.7	17
7	Hemodynamic Significance of Internal Carotid or Middle Cerebral Artery Stenosis Detected on Magnetic Resonance Angiography. Yonsei Medical Journal, 2015, 56, 1686.	0.9	5
8	18F-FDG PET/CT in hepatocellular carcinoma. Nuclear Medicine Communications, 2015, 36, 226-233.	0.5	23
9	Usefulness of 131I-SPECT/CT and 18F-FDG PET/CT in Evaluating Successful 131I and Retinoic Acid Combined Therapy in a Patient with Metastatic Struma Ovarii. Nuclear Medicine and Molecular Imaging, 2015, 49, 52-56.	0.6	7
10	Usefulness of Additional SPECT/CT Identifying Lymphatico-renal Shunt in a Patient with Chyluria. Nuclear Medicine and Molecular Imaging, 2015, 49, 61-64.	0.6	8
11	Positron Emission Tomography/Magnetic Resonance Imaging Evaluation of Lung Cancer. Journal of Thoracic Imaging, 2014, 29, 4-16.	0.8	33
12	A Hepatoid Adenocarcinoma of the Stomach Evaluated With 18F-FDG PET/CT. Clinical Nuclear Medicine, 2014, 39, 442-445.	0.7	7
13	Usefulness of Integrated PET/MRI in Head and Neck Cancer: A Preliminary Study. Nuclear Medicine and Molecular Imaging, 2014, 48, 98-105.	0.6	34
14	Correlation Between 18F-Fluorodeoxyglucose Uptake and Epidermal Growth Factor Receptor Mutations in Advanced Lung Cancer. Nuclear Medicine and Molecular Imaging, 2012, 46, 169-175.	0.6	21
15	Gadoxetate Disodium-Enhanced Magnetic Resonance Imaging Versus Contrast-Enhanced 18F-Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography for the Detection of Colorectal Liver Metastases. Investigative Radiology, 2011, 46, 548-555.	3.5	83
16	Prognostic value of 18Fâ€FDG PET for hepatocellular carcinoma patients treated with sorafenib. Liver International, 2011, 31, 1144-1149.	1.9	56
17	Using 18F-FDG PET/CT to Detect an Occult Mesenchymal Tumor Causing Oncogenic Osteomalacia. Nuclear Medicine and Molecular Imaging, 2011, 45, 233-237.	0.6	10
18	Evaluation of Bone Metastasis from Hepatocellular Carcinoma Using 18F-FDG PET/CT and 99mTc-HDP Bone Scintigraphy: Characteristics of Soft Tissue Formation. Nuclear Medicine and Molecular Imaging, 2011, 45, 203-211.	0.6	13

#	Article	IF	CITATIONS
19	Clinical Use of Radiopharmaceuticals in Boron Neutron Capture Therapy. Nuclear Medicine and Molecular Imaging, O, , 1.	0.6	Ο