
Michal Hoskovec

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3896142/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Cholesteryl esters of ω-(O-acyl)-hydroxy fatty acids in vernix caseosa. Journal of Lipid Research, 2017, 58, 1579-1590.	4.2	22
2	Novel, male-produced aggregation pheromone of the cerambycid beetle Rosalia alpina, a priority species of European conservation concern. PLoS ONE, 2017, 12, e0183279.	2.5	19
3	Analysis of 1,2-diol diesters in vernix caseosa by high-performance liquid chromatography – atmospheric pressure chemical ionization mass spectrometry. Journal of Chromatography A, 2015, 1378, 8-18.	3.7	10
4	Analyses of volatiles produced by the African fruit fly species complex (Diptera, Tephritidae). ZooKeys, 2015, 540, 385-404.	1.1	12
5	Resolution of three cryptic agricultural pests (<i>Ceratitis fasciventris, C. anonae, C. rosa</i> ,) Tj ETQq1 1 0.78 2014, 104, 631-638.	4314 rgBT 1.0	/Overlock 10 39
6	Volatiles from spruce trap-trees detected by Ips typographus bark beetles: chemical and electrophysiological analyses. Arthropod-Plant Interactions, 2014, 8, 305-316.	1.1	25
7	Pheromone Analyses of the <i>Anastrepha fraterculus</i> (Diptera: Tephritidae) Cryptic Species Complex. Florida Entomologist, 2013, 96, 1107-1115.	0.5	37
8	Cuticular Hydrocarbons of the South American Fruit Fly Anastrepha fraterculus: Variability with Sex and Age. Journal of Chemical Ecology, 2012, 38, 1133-1142.	1.8	40
9	Are the Wild and Laboratory Insect Populations Different in Semiochemical Emission? The Case of the Medfly Sex Pheromone. Journal of Agricultural and Food Chemistry, 2012, 60, 7168-7176.	5.2	33
10	Chiral and Nonchiral GC×GC/TOFMS Analysis of Natural Compounds: The Case of Possible Aggregation Pheromones of Chinese Bark Beetles Ips shangrila and Ips nitidus. , 2012, , .		1
11	Indirect Determination of Vapor Pressures by Capillary Gas–Liquid Chromatography: Analysis of the Reference Vapor-Pressure Data and Their Treatment. Journal of Chemical & Engineering Data, 2012, 57, 1349-1368.	1.9	27
12	Identification of two components of the female sex pheromone of the sugarcaneâ€borer <i>Diatraea flavipennella</i> (Lepidoptera: Crambidae). Journal of Applied Entomology, 2012, 136, 203-211.	1.8	2
13	Chemical communication in termites: syn-4,6-dimethylundecan-1-ol as trail-following pheromone, syn-4,6-dimethylundecanal and (5E)-2,6,10-trimethylundeca-5,9-dienal as the respective male and female sex pheromones in Hodotermopsis sjoestedti (Isoptera, Archotermopsidae). Journal of Insect Physiology. 2011, 57, 1585-1591.	2.0	12
14	Irresistible bouquet of death—how are burying beetles (Coleoptera: Silphidae: Nicrophorus) attracted by carcasses. Die Naturwissenschaften, 2009, 96, 889-899.	1.6	122
15	Use of infochemicals to attract carrion beetles into pitfall traps. Entomologia Experimentalis Et Applicata, 2009, 132, 59-64.	1.4	32
16	GC×GC/TOF MS technique—A new tool in identification of insect pheromones: Analysis of the persimmon bark borer sex pheromone gland. Talanta, 2006, 69, 542-547.	5.5	25
17	Determining the vapour pressures of plant volatiles from gas chromatographic retention data. Journal of Chromatography A, 2005, 1083, 161-172.	3.7	73
18	(11Z)-hexadec-11-enal enhances the attractiveness of Diatraea saccharalis main pheromone component in wind tunnel experiments. Journal of Applied Entomology, 2005, 129, 70-74.	1.8	8

MICHAL HOSKOVEC

#	Article	IF	CITATIONS
19	Sex pheromone of horse-chestnut leafminer Camneraria ohridella and its use in a pheromone-based monitoring system. Journal of Chemical Ecology, 2003, 29, 387-404.	1.8	22
20	Efficient syntheses of (10E,12Z,15Z)-9-oxo- and (9Z,11E,15E)-13-oxo-octadecatrienoic acids; two stress metabolites of wounded plants. Tetrahedron, 2002, 58, 3271-3274.	1.9	17
21	Biosynthesis of sex pheromones in moths: stereochemistry of fatty alcohol oxidation in Manduca sexta. Tetrahedron, 2002, 58, 9193-9201.	1.9	14
22	Detection of Sex Pheromone Components in Manduca sexta (L.). Chemical Senses, 2001, 26, 1175-1186.	2.0	52
23	Analysis of the Silkworm Moth Pheromone Binding Protein–Pheromone Complex by Electrospray-Ionization Mass Spectrometry. Angewandte Chemie - International Edition, 2000, 39, 4341-4343.	13.8	28
24	Synthesis of (8E,10Z)-Tetradeca-8,10-dienal, Sex Pheromone of Horse Chestnut Leafminer (Cameraria) Tj ETQq0 0 2000, 65, 511-523.	0 rgBT /C 1.0	overlock 10 7 11
25	Identification of a new lepidopteran sex pheromone in picogram quantities using an antennal biodetector: (8E,10Z)-tetradeca-8,10-dienal from Cameraria ohridella. Tetrahedron Letters, 1999, 40, 7011-7014.	1.4	42
26	Additivity of vaporization properties in pheromone-like homologous series. Journal of the Chemical Society Perkin Transactions II, 1998, , 1351-1356.	0.9	7
27	New Potential Inhibitors of Pheromonal Attractionin the Oriental Fruit Moth, Cydia molesta. Collection of Czechoslovak Chemical Communications, 1998, 63, 1031-1044.	1.0	2
28	Initial water content and lipase-mediated ester formation in hexane. Biotechnology Letters, 1997, 19, 745-750.	2.2	8
29	Gas chromatographic determination of vapour pressures of pheromone-like compounds IV. Acetates, a reinvestigation. Journal of Chromatography A, 1997, 759, 93-109.	3.7	14
30	Gas chromatographic determination of vapour pressures of pheromone-like compounds III. Aldehydes. Journal of Chromatography A, 1996, 719, 391-400.	3.7	10
31	Biotransformations of Î ³ -methyl-Î ² -ketosulfones: stereoselectivity of 3-methyl-1-(phenylsulfonyl)hexan-2-one reductions by various yeasts. Tetrahedron: Asymmetry, 1996, 7, 1285-1294.	1.8	24
32	New mimics of the acetate function in pheromone-based attraction. Bioorganic and Medicinal Chemistry, 1996, 4, 479-488.	3.0	12
33	Synthesis of (Z)-14-Heptadecen-4-olide and (Z)-11-Pentadecen-4-olide, Sex Pheromone Analogues of Ostrinia nubilalis and Cydia molesta. Collection of Czechoslovak Chemical Communications, 1994, 59, 1211-1218.	1.0	2
34	Gas chromatographic determination of vapour pressures of pheromone-like compounds II. Alcohols. Journal of Chromatography A, 1994, 679, 307-317.	3.7	11
35	α,α-Disubstituted Allyl Sulfones: An approach to the synthesis of vinyl-branched pheromone analogues. Helvetica Chimica Acta, 1994, 77, 1281-1287.	1.6	5
36	Estimation of Critical Properties from the Number of Carbon Atoms in Homologous Series. Collection of Czechoslovak Chemical Communications, 1994, 59, 1483-1494.	1.0	5

#	Article	IF	CITATIONS
37	Structure-activity correlations among analogs of the currant clearwing moth pheromone. Journal of Chemical Ecology, 1993, 19, 735-750.	1.8	10
38	Gas chromatographic determination of vapour pressures of pheromone-like acetates. Journal of Chromatography A, 1992, 626, 215-221.	3.7	11
39	A simple method for estimating vaporization properties of pheromone-like acetates from their molecular structures. Collection of Czechoslovak Chemical Communications, 1991, 56, 727-735.	1.0	2
40	A convenient synthesis of 2,13- and 3,13-octadecadienyl acetates, sex pheromone components of the Synanthedon species. Collection of Czechoslovak Chemical Communications, 1990, 55, 2270-2281.	1.0	14
41	Mannich and Grignard reaction of some N-(2-propynyl)azaheterocycles. Collection of Czechoslovak Chemical Communications, 1989, 54, 1067-1081.	1.0	1