
## Marcel Spaargaren

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3896883/publications.pdf Version: 2024-02-01



| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Design and standardization of PCR primers and protocols for detection of clonal immunoglobulin<br>and T-cell receptor gene recombinations in suspect lymphoproliferations: Report of the BIOMED-2<br>Concerted Action BMH4-CT98-3936. Leukemia, 2003, 17, 2257-2317. | 3.3 | 2,788     |
| 2  | The clinically active BTK inhibitor PCI-32765 targets B-cell receptor– and chemokine-controlled adhesion and migration in chronic lymphocytic leukemia. Blood, 2012, 119, 2590-2594.   | 0.6 | 493       |
| 3  | EuroClonality/BIOMED-2 guidelines for interpretation and reporting of Ig/TCR clonality testing in suspected lymphoproliferations. Leukemia, 2012, 26, 2159-2171.   | 3.3 | 409       |
| 4  | Differential Interaction of the Ras Family GTP-binding Proteins H-Ras, Rap1A, and R-Ras with the Putative<br>Effector Molecules Raf Kinase and Ral-Guanine Nucleotide Exchange Factor. Journal of Biological<br>Chemistry, 1996, 271, 6794-6800.                     | 1.6 | 298       |
| 5  | Illegitimate WNT signaling promotes proliferation of multiple myeloma cells. Proceedings of the National Academy of Sciences of the United States of America, 2004, 101, 6122-6127.  | 3.3 | 293       |
| 6  | Significantly improved PCR-based clonality testing in B-cell malignancies by use of multiple<br>immunoglobulin gene targets. Report of the BIOMED-2 Concerted Action BHM4-CT98-3936. Leukemia,<br>2007, 21, 207-214.   | 3.3 | 292       |
| 7  | Bruton's Tyrosine Kinase and Phospholipase Cγ2 Mediate Chemokine-Controlled B Cell Migration and<br>Homing. Immunity, 2007, 26, 93-104.  | 6.6 | 262       |
| 8  | Identification of the guanine nucleotide dissociation stimulator for Ral as a putative effector<br>molecule of R-ras, H-ras, K-ras, and Rap Proceedings of the National Academy of Sciences of the<br>United States of America, 1994, 91, 12609-12613.               | 3.3 | 251       |
| 9  | Cell surface proteoglycan syndecan-1 mediates hepatocyte growth factor binding and promotes Met signaling in multiple myeloma. Blood, 2002, 99, 1405-1410.   | 0.6 | 235       |
| 10 | Powerful strategy for polymerase chain reaction-based clonality assessment in T-cell malignancies<br>Report of the BIOMED-2 Concerted Action BHM4 CT98-3936. Leukemia, 2007, 21, 215-221.  | 3.3 | 222       |
| 11 | The B Cell Antigen Receptor Controls Integrin Activity through Btk and PLCγ2. Journal of Experimental Medicine, 2003, 198, 1539-1550.  | 4.2 | 211       |
| 12 | Heparan sulfate proteoglycan binding promotes APRIL-induced tumor cell proliferation. Cell Death and Differentiation, 2005, 12, 637-648.   | 5.0 | 204       |
| 13 | Heparan Sulfate-modified CD44 Promotes Hepatocyte Growth Factor/Scatter Factor-induced Signal<br>Transduction through the Receptor Tyrosine Kinase c-Met. Journal of Biological Chemistry, 1999, 274,<br>6499-6506.  | 1.6 | 198       |
| 14 | Egress of CD19+CD5+ cells into peripheral blood following treatment with the Bruton tyrosine kinase inhibitor ibrutinib in mantle cell lymphoma patients. Blood, 2013, 122, 2412-2424.   | 0.6 | 185       |
| 15 | High prevalence of oncogenic MYD88 and CD79B mutations in diffuse large B-cell lymphomas presenting at immune-privileged sites. Blood Cancer Journal, 2013, 3, e139-e139.  | 2.8 | 164       |
| 16 | Deletion of the WNT Target and Cancer Stem Cell Marker CD44 in Apc(Min/+) Mice Attenuates<br>Intestinal Tumorigenesis. Cancer Research, 2008, 68, 3655-3661.   | 0.4 | 163       |
| 17 | Lymphoma dissemination: the other face of lymphocyte homing. Blood, 2007, 110, 3102-3111.  | 0.6 | 157       |
| 18 | The hepatocyte growth factor/Met pathway controls proliferation and apoptosis in multiple myeloma.<br>Leukemia, 2003, 17, 764-774.   | 3.3 | 145       |

2

MARCEL SPAARGAREN

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Aberrant Wnt signaling in multiple myeloma: molecular mechanisms and targeting options. Leukemia, 2019, 33, 1063-1075.  | 3.3 | 119       |
| 20 | Stem cell CD44v isoforms promote intestinal cancer formation in Apc(min) mice downstream of Wnt signaling. Oncogene, 2014, 33, 665-670.   | 2.6 | 116       |
| 21 | Primary Follicular Lymphoma of the Small Intestine. American Journal of Pathology, 2003, 162, 105-113.  | 1.9 | 96        |
| 22 | The hepatocyte growth factor/ met pathway in development, tumorigenesis, and B-cell differentiation.<br>Advances in Cancer Research, 2000, 79, 39-90.                               | 1.9 | 95        |
| 23 | High prevalence of oncogenic MYD88 and CD79B mutations in primary testicular diffuse large B-cell<br>lymphoma. Leukemia, 2014, 28, 719-720.   | 3.3 | 91        |
| 24 | FOXP1 directly represses transcription of proapoptotic genes and cooperates with NF-κB to promote survival of human B cells. Blood, 2014, 124, 3431-3440.                           | 0.6 | 86        |
| 25 | Factor VIIa/Tissue Factor-induced Signaling via Activation of Src-like Kinases, Phosphatidylinositol<br>3-Kinase, and Rac. Journal of Biological Chemistry, 2000, 275, 28750-28756. | 1.6 | 85        |
| 26 | Association of RACK1 and PKCl <sup>2</sup> with the common l <sup>2</sup> -chain of the IL-5/IL-3/GM-CSF receptor. Oncogene, 1999, 18, 5126-5130.                                   | 2.6 | 81        |
| 27 | Functional analysis of HGF/MET signaling and aberrant HGF-activator expression in diffuse large B-cell<br>lymphoma. Blood, 2006, 107, 760-768.                                      | 0.6 | 80        |
| 28 | Ibrutinib and idelalisib synergistically target BCR-controlled adhesion in MCL and CLL: a rationale for combination therapy. Blood, 2015, 125, 2306-2309.                           | 0.6 | 79        |
| 29 | Rab5 Induces Rac-independent Lamellipodia Formation and Cell Migration. Molecular Biology of the Cell, 1999, 10, 3239-3250.   | 0.9 | 77        |
| 30 | Human sprouty 4, a new ras antagonist on 5q31, interacts with the dual specificity kinase TESK1. FEBS<br>Journal, 2002, 269, 2546-2556.   | 0.2 | 76        |
| 31 | Expression of c-Met and Heparan-Sulfate Proteoglycan Forms of CD44 in Colorectal Cancer. American<br>Journal of Pathology, 2000, 157, 1563-1573.                                    | 1.9 | 75        |
| 32 | Syndecan-1 promotes Wnt/β-catenin signaling in multiple myeloma by presenting Wnts and R-spondins.<br>Blood, 2018, 131, 982-994.  | 0.6 | 68        |
| 33 | The Ras-related protein R-ras interacts directly with Raf-1 in a GTP-dependent manner. Biochemical<br>Journal, 1994, 300, 303-307.  | 1.7 | 62        |
| 34 | c-Cbl Is Involved in Met Signaling in B Cells and Mediates Hepatocyte Growth Factor-Induced Receptor<br>Ubiquitination. Journal of Immunology, 2002, 169, 3793-3800.                | 0.4 | 57        |
| 35 | Characterization and identification of an epidermal-growth-factor-activated phospholipase A2.<br>Biochemical Journal, 1992, 287, 37-43.   | 1.7 | 56        |
| 36 | <i>MYD88</i> mutations identify a molecular subgroup of diffuse large B-cell lymphoma with an unfavorable prognosis. Haematologica, 2020, 105, 424-434.                             | 1.7 | 55        |

MARCEL SPAARGAREN

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | Multiple myeloma cells catalyze hepatocyte growth factor (HGF) activation by secreting the serine protease HGF-activator. Blood, 2004, 104, 2172-2175.  | 0.6 | 54        |
| 38 | Tubular epithelial syndecan-1 maintains renal function in murine ischemia/reperfusion and human<br>transplantation. Kidney International, 2012, 81, 651-661.  | 2.6 | 54        |
| 39 | Epidermal Growth Factor (EGF) Induces Serine Phosphorylation-Dependent Activation and<br>Calcium-Dependent Translocation of the Cytosolic Phospholipase A2. FEBS Journal, 1995, 231, 593-601.                               | 0.2 | 53        |
| 40 | Transcriptional Silencing of the Wnt-Antagonist Dickkopf-1 (DKK1) by Promoter Methylation Unleashes<br>Aberrant Wnt Signaling In Advanced Multiple Myeloma. Blood, 2010, 116, 1919-1919.                                    | 0.6 | 53        |
| 41 | Targeting EXT1 reveals a crucial role for heparan sulfate in the growth of multiple myeloma. Blood, 2010, 115, 601-604.   | 0.6 | 50        |
| 42 | Infection and transmission of SARS oVâ€2 depend on heparan sulfate proteoglycans. EMBO Journal,<br>2021, 40, e106765.   | 3.5 | 50        |
| 43 | Disruption of heparan sulfate proteoglycan conformation perturbs B-cell maturation and APRIL-mediated plasma cell survival. Blood, 2011, 117, 6162-6171.  | 0.6 | 48        |
| 44 | MET Signaling Mediates Intestinal Crypt-Villus Development, Regeneration, and Adenoma Formation and Is Promoted by Stem Cell CD44 Isoforms. Gastroenterology, 2017, 153, 1040-1053.e4.                                      | 0.6 | 48        |
| 45 | Heparan sulfate proteoglycans in the control of <scp>B</scp> cell development and the pathogenesis of multiple myeloma. FEBS Journal, 2013, 280, 2180-2193.   | 2.2 | 47        |
| 46 | Regulation of Cytokine Signaling by B Cell Antigen Receptor and Cd40-Controlled Expression of<br>Heparan Sulfate Proteoglycans. Journal of Experimental Medicine, 2000, 192, 1115-1124.                                     | 4.2 | 46        |
| 47 | The small GTPase Ral mediates SDF-1–induced migration of B cells and multiple myeloma cells. Blood, 2008, 111, 3364-3372.   | 0.6 | 43        |
| 48 | The forkhead transcription factor FOXP1 represses human plasma cell differentiation. Blood, 2015, 126, 2098-2109.   | 0.6 | 42        |
| 49 | Illegitimate WNT Pathway Activation by $\hat{l}^2$ -Catenin Mutation or Autocrine Stimulation in T-Cell Malignancies. Cancer Research, 2008, 68, 6969-6977.   | 0.4 | 41        |
| 50 | Transcriptional Silencing of the Wnt-Antagonist DKK1 by Promoter Methylation Is Associated with<br>Enhanced Wnt Signaling in Advanced Multiple Myeloma. PLoS ONE, 2012, 7, e30359.  | 1.1 | 41        |
| 51 | The hypoxia target adrenomedullin is aberrantly expressed in multiple myeloma and promotes angiogenesis. Leukemia, 2013, 27, 1729-1737.   | 3.3 | 41        |
| 52 | Tubulointerstitial heparan sulfate proteoglycan changes in human renal diseases correlate with<br>leukocyte influx and proteinuria. American Journal of Physiology - Renal Physiology, 2008, 294,<br>F253-F263.             | 1.3 | 39        |
| 53 | Hepatocyte growth factor triggers signaling cascades mediating vascular smooth muscle cell<br>migration. Biochemical and Biophysical Research Communications, 2002, 298, 80-86.   | 1.0 | 37        |
| 54 | Aberrantly expressed LGR4 empowers Wnt signaling in multiple myeloma by hijacking<br>osteoblast-derived R-spondins. Proceedings of the National Academy of Sciences of the United States<br>of America, 2017, 114, 376-381. | 3.3 | 37        |

MARCEL SPAARGAREN

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Interaction of Epidermal growth factor receptors with the cytoskeleton is related to receptor clustering. Journal of Cellular Physiology, 1990, 145, 365-375.  | 2.0 | 36        |
| 56 | N-cadherin-mediated interaction with multiple myeloma cells inhibits osteoblast differentiation.<br>Haematologica, 2011, 96, 1653-1661.  | 1.7 | 36        |
| 57 | Loss of CYLD expression unleashes Wnt signaling in multiple myeloma and is associated with aggressive disease. Oncogene, 2017, 36, 2105-2115.  | 2.6 | 34        |
| 58 | The B Cell Antigen Receptor Controls AP-1 and NFAT Activity through Ras-Mediated Activation of Ral.<br>Journal of Immunology, 2007, 178, 1405-1414.  | 0.4 | 31        |
| 59 | Ibrutinib and idelalisib target B cell receptor- but not CXCL12/CXCR4-controlled integrin-mediated adhesion in Waldenstrom macroglobulinemia. Haematologica, 2016, 101, e111-e115.   | 1.7 | 30        |
| 60 | BTK inhibitors in chronic lymphocytic leukemia: a glimpse to the future. Oncogene, 2015, 34, 2426-2436.  | 2.6 | 29        |
| 61 | The HGF/MET pathway as target for the treatment of multiple myeloma and B-cell lymphomas.<br>Biochimica Et Biophysica Acta: Reviews on Cancer, 2010, 1806, 208-219.  | 3.3 | 28        |
| 62 | Cell lines generated from a chronic lymphocytic leukemia mouse model exhibit constitutive Btk and<br>Akt signaling. Oncotarget, 2017, 8, 71981-71995.  | 0.8 | 27        |
| 63 | WNT signaling controls expression of pro-apoptotic BOK and BAX in intestinal cancer. Biochemical and Biophysical Research Communications, 2011, 406, 1-6.  | 1.0 | 26        |
| 64 | Hepatocyte growth factor/MET and CD44 in colorectal cancer: partners in tumorigenesis and therapy resistance. Biochimica Et Biophysica Acta: Reviews on Cancer, 2020, 1874, 188437.  | 3.3 | 26        |
| 65 | Impaired Lymphoid Organ Development in Mice Lacking the Heparan Sulfate Modifying Enzyme<br>Glucuronyl C5-Epimerase. Journal of Immunology, 2010, 184, 3656-3664.  | 0.4 | 25        |
| 66 | Follicular Dendritic Cells Catalyze Hepatocyte Growth Factor (HGF) Activation in the Germinal Center<br>Microenvironment by Secreting the Serine Protease HGF Activator. Journal of Immunology, 2005, 175,<br>2807-2813.                     | 0.4 | 24        |
| 67 | Stimulated plasmacytoid dendritic cells impair human T-cell development. Blood, 2006, 108, 3792-3800.  | 0.6 | 24        |
| 68 | CD44 Expression in Intestinal Epithelium and Colorectal Cancer Is Independent of p53 Status. PLoS<br>ONE, 2013, 8, e72849.   | 1.1 | 23        |
| 69 | AKT signaling restrains tumor suppressive functions of FOXO transcription factors and GSK3 kinase in multiple myeloma. Blood Advances, 2020, 4, 4151-4164.   | 2.5 | 20        |
| 70 | R-Ras Alters Ca2+ Homeostasis by Increasing the Ca2+ Leak across the Endoplasmic Reticular<br>Membrane. Journal of Biological Chemistry, 2003, 278, 13672-13679.   | 1.6 | 18        |
| 71 | The small FOXP1 isoform predominantly expressed in activated B cell-like diffuse large B-cell lymphoma<br>and full-length FOXP1 exert similar oncogenic and transcriptional activity in human B cells.<br>Haematologica, 2017, 102, 573-583. | 1.7 | 18        |
| 72 | The pan phosphoinositide 3-kinase/mammalian target of rapamycin inhibitor SAR245409<br>(voxtalisib/XL765) blocks survival, adhesion and proliferation of primary chronic lymphocytic<br>leukemia cells. Leukemia, 2016, 30, 337-345.         | 3.3 | 17        |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | The CXCL12gamma chemokine immobilized by heparan sulfate on stromal niche cells controls adhesion and mediates drug resistance in multiple myeloma. Journal of Hematology and Oncology, 2021, 14, 11.                              | 6.9 | 15        |
| 74 | MET Signaling Overcomes Epidermal Growth Factor Receptor Inhibition in Normal and Colorectal Cancer Stem Cells Causing Drug Resistance. Gastroenterology, 2019, 157, 1153-1155.e1.   | 0.6 | 14        |
| 75 | Identification of the SRC-family tyrosine kinase HCK as a therapeutic target in mantle cell lymphoma.<br>Leukemia, 2021, 35, 881-886.  | 3.3 | 14        |
| 76 | Syndecan-1 and stromal heparan sulfate proteoglycans: key moderators of plasma cell biology and myeloma pathogenesis. Blood, 2021, 137, 1713-1718.   | 0.6 | 14        |
| 77 | The anaphase-promoting complex/cyclosome: a new promising target in diffuse large B-cell lymphoma and mantle cell lymphoma. British Journal of Cancer, 2019, 120, 1137-1146.   | 2.9 | 12        |
| 78 | Diffuse large B cell lymphomas relapsing in the CNS lack oncogenic MYD88 and CD79B mutations.<br>Blood Cancer Journal, 2014, 4, e266-e266.   | 2.8 | 11        |
| 79 | Antibody-induced activation of the epidermal growth factor receptor tyrosine kinase requires the presence of detergent. Biochemical and Biophysical Research Communications, 1990, 171, 882-889.                                   | 1.0 | 10        |
| 80 | Tipping the balance: toward rational combination therapies to overcome venetoclax resistance in mantle cell lymphoma. Leukemia, 2022, 36, 2165-2176.   | 3.3 | 8         |
| 81 | Possible Mechanisms Of Resistance To The Novel BH3-Mimetic ABT-199 In In Vitro Lymph Node Models Of<br>CLL – The Role Of Abl and Btk. Blood, 2013, 122, 4188-4188.   | 0.6 | 6         |
| 82 | Redirecting T-cell Activity with Anti-BCMA/Anti-CD3 Bispecific Antibodies in Chronic Lymphocytic Leukemia and Other B-cell Lymphomas. Cancer Research Communications, 2022, 2, 330-341.  | 0.7 | 6         |
| 83 | Targeting cell adhesion and homing as strategy to cure Waldenström's macroglobulinemia. Best<br>Practice and Research in Clinical Haematology, 2016, 29, 161-168.  | 0.7 | 5         |
| 84 | Immune evasion in primary testicular and central nervous system lymphomas: HLA loss rather than<br>9p24.1/ <i>PD-L1/PD-L2</i> alterations. Blood, 2021, 138, 1194-1197.  | 0.6 | 5         |
| 85 | Lymphoma spread? Target CD47-SIRPα!. Blood, 2011, 118, 4762-4764.  | 0.6 | 4         |
| 86 | A loss-of-adhesion CRISPR-Cas9 screening platform to identify cell adhesion-regulatory proteins and signaling pathways. Nature Communications, 2022, 13, 2136.   | 5.8 | 4         |
| 87 | Instant conditional transgenesis in the mouse hematopoietic compartment. Journal of Immunological<br>Methods, 2008, 339, 259-263.  | 0.6 | 3         |
| 88 | Combined Inhibition of mTOR and DNA-PK Blocks Survival, Adhesion, Proliferation and<br>Chemoresistance in Primary Chronic Lymphocytic Leukemia (CLL) Cells. Blood, 2014, 124, 1981-1981.   | 0.6 | 3         |
| 89 | Mapping the Targets of Dasatinib in Chronic Lymphocytic Leukemia Reveals Distinct Roles for Abl and<br>Btk in Drug Resistance and Adhesion, and Explains Clinical Effects On Lymph Node Reduction. Blood,<br>2012, 120, 3900-3900. | 0.6 | 2         |
| 90 | lbrutinib for AML? Check CD117 (KIT)!. Lancet Haematology,the, 2015, 2, e180-e181.   | 2.2 | 1         |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 91 | Targeting EXT-1 Reveals a Crucial Role of Heparan Sulfate in the Growth of Multiple Myeloma Blood, 2009, 114, 1830-1830.  | 0.6 | 1         |
| 92 | Egress of CD19+CD5+ Cells Into Peripheral Blood Following Treatment with the Bruton Tyrosine<br>Kinase Inhibitor, PCI-32765, in Mantle Cell Lymphoma Patients. Blood, 2011, 118, 954-954.   | 0.6 | 1         |
| 93 | Combined Inhibition of Phosphatidylinositol 3-Kinase (PI3K) Isoform α and δ By the Pan-Class I PI3K<br>Inhibitor SAR245409 (XL765) in Primary Chronic Lymphocytic Leukemia Cells Blocks Survival, Adhesion<br>and Proliferation. Blood, 2014, 124, 4691-4691. | 0.6 | 1         |
| 94 | Aberrant Wnt signaling in multiple myeloma: molecular mechanisms and targeting options. Clinical<br>Lymphoma, Myeloma and Leukemia, 2019, 19, e108.   | 0.2 | 0         |
| 95 | Syndecan-1 and stromal HSPGs: key moderators of communication between myeloma plasma cells and the bone marrow niche. Clinical Lymphoma, Myeloma and Leukemia, 2019, 19, e96-e97.   | 0.2 | 0         |
| 96 | Dimerization Activates the Epidermal Growth Factor Receptor Tyrosine Kinase. , 1991, , 45-58.   |     | 0         |
| 97 | General Mechanistic Patterns of Signal Transduction Across Membranes. , 0, , 1-59.  |     | 0         |