

Mi-Suk Park

List of Publications by Citations

Source: <https://exaly.com/author-pdf/391132/mi-suk-park-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

196
papers

4,161
citations

34
h-index

54
g-index

202
ext. papers

4,910
ext. citations

6.3
avg, IF

5.18
L-index

#	Paper	IF	Citations
196	Differentiation of extrahepatic bile duct cholangiocarcinoma from benign stricture: findings at MRCP versus ERCP. <i>Radiology</i> , 2004 , 233, 234-40	20.5	203
195	CT and PET in stomach cancer: preoperative staging and monitoring of response to therapy. <i>Radiographics</i> , 2006 , 26, 143-56	5.4	133
194	Perfusion CT: noninvasive surrogate marker for stratification of pancreatic cancer response to concurrent chemo- and radiation therapy. <i>Radiology</i> , 2009 , 250, 110-7	20.5	123
193	Can microvessel invasion of hepatocellular carcinoma be predicted by pre-operative MRI?. <i>European Radiology</i> , 2009 , 19, 1744-51	8	118
192	Complete response at first chemoembolization is still the most robust predictor for favorable outcome in hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2015 , 62, 1304-10	13.4	108
191	Diffusion-weighted MR imaging of liver on 3.0-Tesla system: effect of intravenous administration of gadoteric acid disodium. <i>European Radiology</i> , 2010 , 20, 1052-60	8	90
190	Hepatocellular carcinoma: detection with diffusion-weighted versus contrast-enhanced magnetic resonance imaging in pretransplant patients. <i>Hepatology</i> , 2012 , 56, 140-8	11.2	87
189	Capecitabine combined with gemcitabine (CapGem) as first-line treatment in patients with advanced/metastatic biliary tract carcinoma. <i>Cancer</i> , 2005 , 104, 2753-8	6.4	86
188	Accuracy of gadoteric acid-enhanced magnetic resonance imaging for the diagnosis of sinusoidal obstruction syndrome in patients with chemotherapy-treated colorectal liver metastases. <i>European Radiology</i> , 2012 , 22, 864-71	8	85
187	Comparison of gadoteric acid-enhanced dynamic imaging and diffusion-weighted imaging for the preoperative evaluation of colorectal liver metastases. <i>Journal of Magnetic Resonance Imaging</i> , 2011 , 34, 345-53	5.6	77
186	Comparison of CT and 18F-FDG pet for detecting peritoneal metastasis on the preoperative evaluation for gastric carcinoma. <i>Korean Journal of Radiology</i> , 2006 , 7, 249-56	6.9	72
185	Preoperative prediction of the microvascular invasion of hepatocellular carcinoma with diffusion-weighted imaging. <i>Liver Transplantation</i> , 2012 , 18, 1171-8	4.5	71
184	Autologous bone marrow infusion activates the progenitor cell compartment in patients with advanced liver cirrhosis. <i>Cell Transplantation</i> , 2010 , 19, 1237-46	4	71
183	Comparison of MRI and endoscopic ultrasound in the characterization of pancreatic cystic lesions. <i>American Journal of Roentgenology</i> , 2010 , 195, 947-52	5.4	69
182	Abdominal cocoon: preoperative diagnostic clues from radiologic imaging with pathologic correlation. <i>American Journal of Roentgenology</i> , 2004 , 182, 639-41	5.4	66
181	Differentiation of hepatic hyperintense lesions seen on gadoteric acid-enhanced hepatobiliary phase MRI. <i>American Journal of Roentgenology</i> , 2011 , 197, W44-52	5.4	64
180	Peritoneal mesotheliomas: clinicopathologic features, CT findings, and differential diagnosis. <i>American Journal of Roentgenology</i> , 2008 , 191, 814-25	5.4	64

179	Effects of neoadjuvant combined chemotherapy and radiation therapy on the CT evaluation of resectability and staging in patients with pancreatic head cancer. <i>Radiology</i> , 2009 , 250, 758-65	20.5	62
178	Response evaluation in patients with colorectal liver metastases: RECIST version 1.1 versus modified CT criteria. <i>American Journal of Roentgenology</i> , 2012 , 199, 809-15	5.4	60
177	Predisposing factors of bile duct injury after transcatheter arterial chemoembolization (TACE) for hepatic malignancy. <i>CardioVascular and Interventional Radiology</i> , 2002 , 25, 270-4	2.7	59
176	Growth rate of early-stage hepatocellular carcinoma in patients with chronic liver disease. <i>Clinical and Molecular Hepatology</i> , 2015 , 21, 279-86	6.9	53
175	Comparison of breathhold, navigator-triggered, and free-breathing diffusion-weighted MRI for focal hepatic lesions. <i>Journal of Magnetic Resonance Imaging</i> , 2013 , 38, 109-18	5.6	52
174	Recurrent pyogenic cholangitis: comparison between MR cholangiography and direct cholangiography. <i>Radiology</i> , 2001 , 220, 677-82	20.5	50
173	Prediction of the histopathological grade of hepatocellular carcinoma using qualitative diffusion-weighted, dynamic, and hepatobiliary phase MRI. <i>European Radiology</i> , 2012 , 22, 1701-8	8	45
172	Acute cholecystitis at T2-weighted and manganese-enhanced T1-weighted MR cholangiography: preliminary study. <i>Radiology</i> , 2003 , 227, 580-4	20.5	45
171	Biliary ductal involvement of hilar cholangiocarcinoma: multidetector computed tomography versus magnetic resonance cholangiography. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 72-8	2.2	41
170	Quantitative evaluation of liver cirrhosis using T1 relaxation time with 3 tesla MRI before and after oxygen inhalation. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 36, 405-10	5.6	39
169	Added value of subtraction imaging in detecting arterial enhancement in small (. <i>European Radiology</i> , 2013 , 23, 924-30	8	38
168	Early biliary complications of laparoscopic cholecystectomy: evaluation on T2-weighted MR cholangiography in conjunction with mangafodipir trisodium-enhanced 3D T1-weighted MR cholangiography. <i>American Journal of Roentgenology</i> , 2004 , 183, 1559-66	5.4	38
167	Small (. <i>American Journal of Roentgenology</i> , 2004 , 182, 1233-40	5.4	38
166	Primary malignant melanoma of the rectum: CT findings in eight patients. <i>Radiology</i> , 2004 , 232, 181-6	20.5	35
165	Hepatocellular carcinoma variants: radiologic-pathologic correlation. <i>American Journal of Roentgenology</i> , 2009 , 193, W7-13	5.4	34
164	Gastric true leiomyoma: computed tomographic findings and pathological correlation. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 204-8	2.2	34
163	Radiologic findings of gastrointestinal tract involvement in hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 95-101	2.2	34
162	Evaluation of treatment response in hepatocellular carcinoma in the explanted liver with Liver Imaging Reporting and Data System version 2017. <i>European Radiology</i> , 2020 , 30, 261-271	8	34

161	Feasibility of 3D navigator-triggered magnetic resonance cholangiopancreatography with combined parallel imaging and compressed sensing reconstruction at 3T. <i>Journal of Magnetic Resonance Imaging</i> , 2017 , 46, 1289-1297	5.6	33
160	Added value of smooth hypointense rim in the hepatobiliary phase of gadoxetic acid-enhanced MRI in identifying tumour capsule and diagnosing hepatocellular carcinoma. <i>European Radiology</i> , 2017 , 27, 2610-2618	8	33
159	Radiological and clinical features of sarcomatoid hepatocellular carcinoma in 11 cases. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 745-9	2.2	33
158	Bile duct injuries leading to portal vein obliteration after transcatheter arterial chemoembolization in the liver: CT findings and initial observations. <i>Radiology</i> , 2001 , 221, 429-36	20.5	33
157	Hepatic hemangiomas with arteriportal shunt: sonographic appearances with CT and MRI correlation. <i>American Journal of Roentgenology</i> , 2006 , 187, W406-14	5.4	32
156	Detection of liver metastases using gadoxetic-enhanced dynamic and 10- and 20-minute delayed phase MR imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 35, 635-43	5.6	31
155	Radiographic findings of primary B-cell lymphoma of the stomach: low-grade versus high-grade malignancy in relation to the mucosa-associated lymphoid tissue concept. <i>American Journal of Roentgenology</i> , 2002 , 179, 1297-304	5.4	31
154	Preoperative MRI of potential living-donor-related liver transplantation using a single dose of gadobenate dimeglumine. <i>American Journal of Roentgenology</i> , 2005 , 185, 424-31	5.4	30
153	Mucinous versus nonmucinous gastric carcinoma: differentiation with helical CT. <i>Radiology</i> , 2002 , 223, 540-6	20.5	30
152	Value of "patent track" sign on Doppler sonography after percutaneous liver biopsy in detection of postbiopsy bleeding: a prospective study in 352 patients. <i>American Journal of Roentgenology</i> , 2007 , 189, 109-16	5.4	29
151	Pain palliation in patients with bone metastases using magnetic resonance-guided focused ultrasound with conformal bone system: a preliminary report. <i>Yonsei Medical Journal</i> , 2015 , 56, 503-9	3	28
150	Pancreatic tumors: emphasis on CT findings and pathologic classification. <i>Korean Journal of Radiology</i> , 2011 , 12, 731-9	6.9	28
149	MR cholangiography for evaluation of hilar branching anatomy in transplantation of the right hepatic lobe from a living donor. <i>American Journal of Roentgenology</i> , 2008 , 191, 537-45	5.4	28
148	Characterization of incidental liver lesions: comparison of multidetector CT versus Gd-EOB-DTPA-enhanced MR imaging. <i>PLoS ONE</i> , 2013 , 8, e66141	3.7	27
147	Differentiation of benign and malignant ampullary obstructions on MR imaging. <i>European Journal of Radiology</i> , 2011 , 80, 198-203	4.7	27
146	Prognostic factors and 10-year survival in patients with hepatocellular carcinoma after curative hepatectomy. <i>Journal of Gastrointestinal Surgery</i> , 2011 , 15, 598-607	3.3	26
145	Hepatic splenosis preoperatively diagnosed as hepatocellular carcinoma in a patient with chronic hepatitis B: a case report. <i>Journal of Korean Medical Science</i> , 2008 , 23, 336-41	4.7	26
144	Preoperative staging accuracy of multidetector row computed tomography for extrahepatic bile duct carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 362-7	2.2	26

143	Scirrhous gastric carcinoma: endoscopy versus upper gastrointestinal radiography. <i>Radiology</i> , 2004 , 231, 421-6	20.5	26
142	Intraindividual comparison of diagnostic performance in patients with hepatic metastasis of full-dose standard and half-dose iterative reconstructions with dual-source abdominal computed tomography. <i>Investigative Radiology</i> , 2014 , 49, 195-200	10.1	25
141	Correlations of dynamic contrast-enhanced magnetic resonance imaging with morphologic, angiogenic, and molecular prognostic factors in rectal cancer. <i>Yonsei Medical Journal</i> , 2013 , 54, 123-30	3	25
140	Use of Imaging to Predict Complete Response of Colorectal Liver Metastases after Chemotherapy: MR Imaging versus CT Imaging. <i>Radiology</i> , 2017 , 284, 423-431	20.5	24
139	Radiation dose reduction via sinogram affirmed iterative reconstruction and automatic tube voltage modulation (CARE kV) in abdominal CT. <i>Korean Journal of Radiology</i> , 2013 , 14, 886-93	6.9	24
138	Role of EUS and MDCT in the diagnosis of gastric submucosal tumors according to the revised pathologic concept of gastrointestinal stromal tumors. <i>European Radiology</i> , 2009 , 19, 924-34	8	24
137	FOLFIRINOX gemcitabine/nab-paclitaxel for treatment of metastatic pancreatic cancer: Single-center cohort study. <i>World Journal of Gastrointestinal Oncology</i> , 2020 , 12, 182-194	3.4	23
136	Correlation between 18F-fluorodeoxyglucose positron emission tomography and pathologic differentiation in pancreatic cancer. <i>Annals of Nuclear Medicine</i> , 2014 , 28, 430-5	2.5	22
135	Intraductal ultrasonography combined with percutaneous transhepatic cholangioscopy for the preoperative evaluation of longitudinal tumor extent in hilar cholangiocarcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2010 , 25, 286-92	4	22
134	Preoperative radiologic and postoperative pathologic risk factors for early intra-hepatic recurrence in hepatocellular carcinoma patients who underwent curative resection. <i>Yonsei Medical Journal</i> , 2009 , 50, 789-95	3	22
133	Advanced gastric carcinoma with signet ring cell carcinoma versus non-signet ring cell carcinoma: differentiation with multidetector CT. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 880-4	2.2	22
132	Retrospective comparison of EASL 2018 and LI-RADS 2018 for the noninvasive diagnosis of hepatocellular carcinoma using magnetic resonance imaging. <i>Hepatology International</i> , 2020 , 14, 70-79	8.8	22
131	Diagnostic Performance of CT/MRI Liver Imaging Reporting and Data System v2017 for Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis. <i>Liver International</i> , 2020 , 40, 1488-1497	7.9	21
130	Differential features of pancreatobiliary- and intestinal-type ampullary carcinomas at MR imaging. <i>Radiology</i> , 2010 , 257, 384-93	20.5	21
129	Choledochal cyst and anomalous pancreaticobiliary ductal union in adults: radiological spectrum and complications. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 17-22	2.2	21
128	Gadoxetic acid-enhanced magnetic resonance imaging: Hepatocellular carcinoma and mimickers. <i>Clinical and Molecular Hepatology</i> , 2019 , 25, 223-233	6.9	21
127	Rectal Mucinous Adenocarcinoma: MR Imaging Assessment of Response to Concurrent Chemotherapy and Radiation Therapy-A Hypothesis-generating Study. <i>Radiology</i> , 2017 , 285, 124-133	20.5	20
126	Dynamic enhancement pattern of HCC smaller than 3 cm in diameter on gadoxetic acid-enhanced MRI: comparison with multiphasic MDCT. <i>Liver International</i> , 2014 , 34, 1593-602	7.9	20

125	Prediction of postoperative pancreatic fistulas after pancreatectomy: assessment with acoustic radiation force impulse elastography. <i>Journal of Ultrasound in Medicine</i> , 2014 , 33, 781-6	2.9	20
124	Intestinal parasitic infection. <i>Abdominal Imaging</i> , 2008 , 33, 166-71		20
123	Quantitative analysis of the effect of iterative reconstruction using a phantom: determining the appropriate blending percentage. <i>Yonsei Medical Journal</i> , 2015 , 56, 253-61	3	19
122	Detection of recurrent hepatocellular carcinoma on post-operative surveillance: comparison of MDCT and gadoxetic acid-enhanced MRI. <i>Abdominal Imaging</i> , 2014 , 39, 291-9		19
121	MRI findings of rectal submucosal tumors. <i>Korean Journal of Radiology</i> , 2011 , 12, 487-98	6.9	19
120	Enhancement patterns of hepatocellular carcinoma after transarterial chemoembolization using drug-eluting beads on arterial phase CT images: a pilot retrospective study. <i>American Journal of Roentgenology</i> , 2012 , 199, 349-59	5.4	19
119	Transient peritumoral enhancement during dynamic MRI of the liver: cavernous hemangioma versus hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 411-7	2.2	19
118	Optimal T2-weighted MR cholangiopancreatographic images can be obtained after administration of gadoxetic acid. <i>Radiology</i> , 2010 , 256, 475-84	20.5	18
117	Unusual cystic neoplasms in the pancreas: radiologic-pathologic correlation. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 610-6	2.2	18
116	Diffusion and perfusion MRI prediction of progression-free survival in patients with hepatocellular carcinoma treated with concurrent chemoradiotherapy. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 39, 286-92	5.6	17
115	Para-aortic lymph node metastasis in patients with intra-abdominal malignancies: CT vs PET. <i>World Journal of Gastroenterology</i> , 2009 , 15, 4434-8	5.6	17
114	Liver stiffness assessment with tagged MRI of cardiac-induced liver motion in cirrhosis patients. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 39, 1301-7	5.6	16
113	Eosinophilic hepatic necrosis: magnetic resonance imaging and computed tomography comparison. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 765-71	2.2	16
112	Metastasis versus focal eosinophilic infiltration of the liver in patients with extrahepatic abdominal cancer: an evaluation with gadobenate dimeglumine-enhanced magnetic resonance imaging. <i>Journal of Computer Assisted Tomography</i> , 2009 , 33, 119-24	2.2	15
111	Imaging findings of leukemic involvement of the pancreaticobiliary system in adults. <i>American Journal of Roentgenology</i> , 2007 , 188, 1589-95	5.4	15
110	The AFSUMB Consensus Statements and Recommendations for the Clinical Practice of Contrast-Enhanced Ultrasound using Sonazoid. <i>Ultrasonography</i> , 2020 , 39, 191-220	4.3	15
109	Gadoxetic acid-enhanced MRI of macrotrabecular-massive hepatocellular carcinoma and its prognostic implications. <i>Journal of Hepatology</i> , 2021 , 74, 109-121	13.4	15
108	Comparison of MR imaging features of solid pseudopapillary neoplasm of pancreas between male and female patients. <i>European Journal of Radiology</i> , 2015 , 84, 2065-70	4.7	14

107	18F -FDG PET metabolic parameters and MRI perfusion and diffusion parameters in hepatocellular carcinoma: a preliminary study. <i>PLoS ONE</i> , 2013 , 8, e71571	3.7	14
106	A Phase II study of capecitabine combined with gemcitabine in patients with advanced gallbladder carcinoma. <i>Yonsei Medical Journal</i> , 2005 , 46, 526-31	3	14
105	Contrast-enhanced US with Perfluorobutane(Sonazoid) used as a surveillance test for Hepatocellular Carcinoma (HCC) in Cirrhosis (SCAN): an exploratory cross-sectional study for a diagnostic trial. <i>BMC Cancer</i> , 2017 , 17, 279	4.8	13
104	Hyperintense nodules on non-enhanced T1-weighted gradient-echo magnetic resonance imaging of cirrhotic liver: fate and clinical implications. <i>Journal of Magnetic Resonance Imaging</i> , 2006 , 24, 630-6	5.6	13
103	Abdominal sparganosis presenting as an abscess with fistulous communication to the bowel. <i>American Journal of Roentgenology</i> , 2005 , 185, 1084-5	5.4	13
102	Contrast-enhanced US with Perfluorobutane for Hepatocellular Carcinoma Surveillance: A Multicenter Diagnostic Trial (SCAN). <i>Radiology</i> , 2019 , 292, 638-646	20.5	12
101	Efficacy and treatment-related adverse events of gemcitabine plus nab-paclitaxel for treatment of metastatic pancreatic cancer "in a Korean" population: A single-center cohort study. <i>Seminars in Oncology</i> , 2017 , 44, 420-427	5.5	12
100	Focal eosinophilic necrosis versus metastasis in the liver: the usefulness of two-phase dynamic CT. <i>American Journal of Roentgenology</i> , 2005 , 184, 1085-90	5.4	12
99	Comparison of CT and MRI for presurgical characterization of paraaortic lymph nodes in patients with pancreatico-biliary carcinoma. <i>World Journal of Gastroenterology</i> , 2008 , 14, 2208-12	5.6	12
98	The Role of Imaging in Current Treatment Strategies for Pancreatic Adenocarcinoma. <i>Korean Journal of Radiology</i> , 2021 , 22, 23-40	6.9	12
97	Imaging-guided minimally invasive laparoscopic resection of intraluminal small-bowel tumor: report of two cases. <i>American Journal of Roentgenology</i> , 2007 , 189, 56-60	5.4	11
96	Hepatic cavernous hemangioma in cirrhotic liver: imaging findings. <i>Korean Journal of Radiology</i> , 2000 , 1, 185-90	6.9	11
95	Lack of anti-tumor activity by anti-VEGF treatments in hepatic hemangiomas. <i>Angiogenesis</i> , 2016 , 19, 147-53	10.6	10
94	MRI Findings and Prediction of Time to Progression of Patients with Hepatocellular Carcinoma Treated with Drug-eluting Bead Transcatheter Arterial Chemoembolization. <i>Journal of Korean Medical Science</i> , 2015 , 30, 965-73	4.7	10
93	Using multi-detector-row CT to diagnose ampullary adenoma or adenocarcinoma in situ. <i>European Journal of Radiology</i> , 2011 , 80, e340-5	4.7	10
92	Primary hepatic schwannoma mimicking malignancy on fluorine-18 2-fluoro-2-deoxy-D-glucose positron emission tomography-computed tomography. <i>Hepatology</i> , 2010 , 51, 1080-1	11.2	10
91	Spontaneous regression of a cystic tumor in a postpartum woman; is it a cystic lymphangioma?. <i>Yonsei Medical Journal</i> , 2007 , 48, 715-8	3	10
90	Solitary fibrous tumor arising from stomach: CT findings. <i>Yonsei Medical Journal</i> , 2007 , 48, 1056-60	3	10

89	Is Non-Contrast CT Adequate for the Evaluation of Hepatic Metastasis in Patients Who Cannot Receive Iodinated Contrast Media?. <i>PLoS ONE</i> , 2015 , 10, e0134133	3.7	10
88	Incremental Role of Pancreatic Magnetic Resonance Imaging after Staging Computed Tomography to Evaluate Patients with Pancreatic Ductal Adenocarcinoma. <i>Cancer Research and Treatment</i> , 2019 , 51, 24-33	5.2	10
87	Optimal lexicon of gadoxetic acid-enhanced magnetic resonance imaging for the diagnosis of hepatocellular carcinoma modified from LI-RADS. <i>Abdominal Radiology</i> , 2019 , 44, 3078-3088	3	9
86	Characterization of focal liver lesions using the stretched exponential model: comparison with monoexponential and biexponential diffusion-weighted magnetic resonance imaging. <i>European Radiology</i> , 2019 , 29, 5111-5120	8	9
85	Long-Term Follow-Up of Patients After Autologous Bone Marrow Cell Infusion for Decompensated Liver Cirrhosis. <i>Cell Transplantation</i> , 2017 , 26, 1059-1066	4	9
84	Gadobenate dimeglumine-enhanced MRI of intraductal papillary mucinous tumor of the bile ducts. <i>Journal of Magnetic Resonance Imaging</i> , 2007 , 25, 625-7	5.6	9
83	MR cholangiography of accessory bile duct connected to the stomach. <i>American Journal of Roentgenology</i> , 2007 , 189, W344-7	5.4	9
82	Hepatic MRI using the double-echo chemical shift phase-selective gradient-echo technique. <i>American Journal of Roentgenology</i> , 2007 , 188, W49-56	5.4	9
81	Dynamic Contrast-Enhanced Magnetic Resonance Imaging as a Surrogate Biomarker for Bevacizumab in Colorectal Cancer Liver Metastasis: A Single-Arm, Exploratory Trial. <i>Cancer Research and Treatment</i> , 2016 , 48, 1210-1221	5.2	9
80	Imaging Features of Hepatocellular Carcinoma: Quantitative and Qualitative Comparison Between MRI-Enhanced With Gd-EOB-DTPA and Gd-DTPA. <i>Investigative Radiology</i> , 2019 , 54, 494-499	10.1	9
79	Clinical Feasibility of MR Elastography in Patients With Biliary Obstruction. <i>American Journal of Roentgenology</i> , 2018 , 210, 1273-1278	5.4	8
78	Hepatocellular carcinoma: clinical and radiological findings in patients with chronic B viral hepatitis and chronic C viral hepatitis. <i>Abdominal Imaging</i> , 2012 , 37, 591-4		8
77	The Modified Response Evaluation Criteria in Solid Tumors (RECIST) Yield a More Accurate Prognoses Than the RECIST 1.1 in Hepatocellular Carcinoma Treated with Transarterial Radioembolization. <i>Gut and Liver</i> , 2020 , 14, 765-774	4.8	8
76	Contrast-enhanced ultrasound liver imaging reporting and data system for diagnosing hepatocellular carcinoma: A meta-analysis. <i>Liver International</i> , 2020 , 40, 2345-2352	7.9	8
75	CT features of hepatic metastases from hepatoid adenocarcinoma. <i>Abdominal Radiology</i> , 2017 , 42, 2402-2409	3	7
74	Diagnostic performance of the LR-M criteria and spectrum of LI-RADS imaging features among primary hepatic carcinomas. <i>Abdominal Radiology</i> , 2020 , 45, 3743-3754	3	7
73	Potential conditions causing impairment of selective hepatobiliary enhancement of gadobenate dimeglumine-enhanced delayed magnetic resonance imaging. <i>Journal of Computer Assisted Tomography</i> , 2010 , 34, 113-20	2.2	7
72	Value of manganese-enhanced T1- and T2-weighted MR cholangiography for differentiating cystic parenchymal lesions from cystic abnormalities which communicate with bile ducts. <i>Yonsei Medical Journal</i> , 2007 , 48, 1072-4	3	7

71	Double common bile duct: curved-planar reformatted computed tomography (CT) and gadobenate dimeglumine-enhanced MR cholangiography. <i>Journal of Magnetic Resonance Imaging</i> , 2008 , 27, 209-11	5.6	7
70	Lymphoma presenting as an echogenic periportal mass: sonographic findings. <i>Journal of Clinical Ultrasound</i> , 2008 , 36, 437-9	1	7
69	FDG uptake in PET by bladder hernia simulating inguinal metastasis. <i>Yonsei Medical Journal</i> , 2007 , 48, 886-90	3	7
68	Magnetic Resonance Imaging for Colorectal Cancer Metastasis to the Liver: Comparative Effectiveness Research for the Choice of Contrast Agents. <i>Cancer Research and Treatment</i> , 2018 , 50, 60-70	5.2	7
67	Liver trauma diagnosis with contrast-enhanced ultrasound: interobserver variability between radiologist and emergency physician in an animal study. <i>American Journal of Emergency Medicine</i> , 2012 , 30, 1229-34	2.9	6
66	Quantification of superparamagnetic iron oxide-mediated signal intensity change in patients with liver cirrhosis using T2 and T2* mapping: a preliminary report. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 31, 1379-86	5.6	6
65	Dynamic contrast-enhanced MRI coupled with a subtraction technique is useful for treatment response evaluation of malignant melanoma hepatic metastasis. <i>Oncotarget</i> , 2016 , 7, 38513-38522	3.3	6
64	Preoperative prediction of postsurgical outcomes in mass-forming intrahepatic cholangiocarcinoma based on clinical, radiologic, and radiomics features. <i>European Radiology</i> , 2021 , 31, 8638-8648	8	6
63	Feasibility of radiation dose reduction with iterative reconstruction in abdominopelvic CT for patients with inappropriate arm positioning. <i>PLoS ONE</i> , 2018 , 13, e0209754	3.7	6
62	Optimal criteria for hepatocellular carcinoma diagnosis using CT in patients undergoing liver transplantation. <i>European Radiology</i> , 2019 , 29, 1022-1031	8	5
61	Hepatobiliary versus Extracellular MRI Contrast Agents in Hepatocellular Carcinoma Detection: Hepatobiliary Phase Features in Relation to Disease-free Survival. <i>Radiology</i> , 2019 , 293, 594-604	20.5	5
60	Preoperative Clinical and Computed Tomography (CT)-Based Nomogram to Predict Oncologic Outcomes in Patients with Pancreatic Head Cancer Resected with Curative Intent: A Retrospective Study. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	5
59	Inter-observer variability of response evaluation criteria for hepatocellular carcinoma treated with chemoembolization. <i>Digestive and Liver Disease</i> , 2015 , 47, 682-8	3.3	5
58	Diagnostic accuracy of multidetector-row computed tomography for common bile duct calculi: is it necessary to add non-contrast-enhanced images to contrast-enhanced images?. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 508-12	2.2	5
57	Gadoxetic acid-enhanced MRI of hepatocellular carcinoma: Diagnostic performance of category-adjusted LR-5 using modified criteria. <i>PLoS ONE</i> , 2020 , 15, e0242344	3.7	5
56	The AFSUMB Consensus Statements and Recommendations for the Clinical Practice of Contrast-Enhanced Ultrasound using Sonazoid. <i>Journal of Medical Ultrasound</i> , 2020 , 28, 59-82	0.8	5
55	Distal common bile duct metastasis from hepatocellular carcinoma. <i>Hepatology</i> , 2012 , 55, 1638-9	11.2	4
54	Imaging findings of biliary and nonbiliary complications following laparoscopic surgery. <i>European Radiology</i> , 2006 , 16, 1906-14	8	4

53	CT/MRI and CEUS LI-RADS Major Features Association with Hepatocellular Carcinoma: Individual Patient Data Meta-Analysis. <i>Radiology</i> , 2021 , 211244	20.5	4
52	MRI Ancillary Features for LI-RADS Category 3 and 4 Observations: Improved Categorization to Indicate the Risk of Hepatic Malignancy. <i>American Journal of Roentgenology</i> , 2020 , 215, 1354-1362	5.4	4
51	Comparison of the current guidelines for diagnosing hepatocellular carcinoma using gadoteric acid-enhanced magnetic resonance imaging. <i>European Radiology</i> , 2021 , 31, 4492-4503	8	4
50	LI-RADS Major Features on MRI for Diagnosing Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 518-525	5.6	4
49	Diagnostic performance of Liver Imaging Reporting and Data System in patients at risk of both hepatocellular carcinoma and metastasis. <i>Abdominal Radiology</i> , 2020 , 45, 3789-3799	3	3
48	Is there association between statin usage and contrast-associated acute kidney injury after intravenous administration of iodine-based contrast media in enhanced computed tomography?. <i>European Radiology</i> , 2020 , 30, 5261-5271	8	3
47	The impact of CT follow-up interval on stages of hepatocellular carcinomas detected during the surveillance of patients with liver cirrhosis. <i>American Journal of Roentgenology</i> , 2012 , 199, 816-21	5.4	3
46	Mouse Hepatic Tumor Vascular Imaging by Experimental Selective Angiography. <i>PLoS ONE</i> , 2015 , 10, e0131687	3.7	3
45	LI-RADS treatment response algorithm for detecting incomplete necrosis in hepatocellular carcinoma after locoregional treatment: a systematic review and meta-analysis using individual patient data. <i>Abdominal Radiology</i> , 2021 , 46, 3717-3728	3	3
44	Initial experience of irreversible electroporation for locally advanced pancreatic cancer in a Korean population. <i>Acta Radiologica</i> , 2021 , 62, 164-171	2	3
43	Clinical Staging of Mass-Forming Intrahepatic Cholangiocarcinoma: Computed Tomography Versus Magnetic Resonance Imaging. <i>Hepatology Communications</i> , 2021 , 5, 2009-2018	6	3
42	Multimodality imaging studies of intraductal tubulopapillary neoplasms of the pancreas. <i>Diagnostic and Interventional Radiology</i> , 2019 , 25, 251-256	3.2	2
41	Characterisation of small hypoattenuating hepatic lesions in multi-detector CT (MDCT) in patients with underlying extrahepatic malignancy: added value of contrast-enhanced MR images. <i>European Radiology</i> , 2010 , 20, 2853-61	8	2
40	Malignant Mixed M¶erian Tumor with Small Bowel Metastasis: A Case Report. <i>Journal of the Korean Society of Magnetic Resonance in Medicine</i> , 2012 , 16, 257		2
39	Noninvasive Biomarker for Predicting Treatment Response to Concurrent Chemoradiotherapy in Patients with Hepatocellular Carcinoma 2019 , 23, 351		2
38	Diagnostic Performance of Liver Imaging Reporting and Data System Version 2017 Versus Version 2018 for Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis of Comparative Studies. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 1912-1919	5.6	2
37	Characteristics and Early Recurrence of Hepatocellular Carcinomas Categorized as LR-M: Comparison with Those Categorized as LR-4 or 5. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 1446-1454	5.6	2
36	Prognostic factors in patients with locally advanced or borderline resectable pancreatic ductal adenocarcinoma: chemotherapy vs. chemoradiotherapy. <i>Abdominal Radiology</i> , 2021 , 46, 655-666	3	2

35	Extended application of subtraction arterial phase imaging in LI-RADS version 2018: a strategy to improve the diagnostic performance for hepatocellular carcinoma on gadoxetate disodium-enhanced MRI. <i>European Radiology</i> , 2021 , 31, 1620-1629	8	2
34	Noninvasive evaluation of liver fibrosis: comparison of the stretched exponential diffusion-weighted model to other diffusion-weighted MRI models and transient elastography. <i>European Radiology</i> , 2021 , 31, 4813-4823	8	2
33	Systematic review and meta-analysis of diagnostic performance of CT imaging for assessing resectability of pancreatic ductal adenocarcinoma after neoadjuvant therapy: importance of CT criteria. <i>Abdominal Radiology</i> , 2021 , 46, 5201-5217	3	2
32	Retrospective Evaluation of Treatment Response in Patients with Nonmetastatic Pancreatic Cancer Using CT and CA 19-9.. <i>Radiology</i> , 2022 , 212236	20.5	2
31	Metal implants influence CT scan parameters leading to increased local radiation exposure: A proposal for correction techniques. <i>PLoS ONE</i> , 2019 , 14, e0221692	3.7	1
30	Electronic clinical challenges and images in GI. Hepatic venoocclusive disease. <i>Gastroenterology</i> , 2008 , 135, e3-4	13.3	1
29	Necrotic lymphoma in a patient with post-transplantation lymphoproliferative disorder: ultrasonography and CT findings with pathologic correlation. <i>Ultrasonography</i> , 2015 , 34, 148-52	4.3	1
28	A lexicon for hepatocellular carcinoma surveillance ultrasonography: benign versus malignant lesions. <i>Clinical and Molecular Hepatology</i> , 2017 , 23, 57-65	6.9	1
27	Magnetization-tagged MRI is a simple method for predicting liver fibrosis. <i>Clinical and Molecular Hepatology</i> , 2016 , 22, 140-5	6.9	1
26	Non-contiguous Multi-organ Involvement of an Inflammatory Myofibroblastic Tumor: A Case Report. <i>Journal of the Korean Radiological Society</i> , 2007 , 57, 265		1
25	Primary Malignant Melanoma of the Esophagus: A Case Report. <i>Journal of the Korean Radiological Society</i> , 2007 , 57, 37		1
24	Radiographic portal or superior mesenteric vein invasion is an independent prognostic factor in non-metastatic pancreatic ductal adenocarcinoma: A missing block of clinical T staging?. <i>Pancreatology</i> , 2020 , 20, 952-959	3.8	1
23	Contrast-enhanced ultrasound Liver Imaging Reporting and Data System category M: a systematic review and meta-analysis. <i>Ultrasonography</i> , 2021 ,	4.3	1
22	Antiangiogenic Therapy Induces Hepatic Tumor Vascular Network Rearrangement to Receive Perfusion via the Portal Vein and Hepatic Artery. <i>Journal of Vascular Research</i> , 2016 , 53, 72-82	1.9	1
21	Radiological patterns of secondary sclerosing cholangitis in patients after lung transplantation. <i>Abdominal Radiology</i> , 2019 , 44, 1361-1366	3	1
20	Liver MRI with amide proton transfer imaging: feasibility and accuracy for the characterization of focal liver lesions. <i>European Radiology</i> , 2021 , 31, 222-231	8	1
19	Should Threshold Growth Be Considered a Major Feature in the Diagnosis of Hepatocellular Carcinoma Using LI-RADS?. <i>Korean Journal of Radiology</i> , 2021 , 22, 1628-1639	6.9	1
18	Effectiveness of Hepatocellular Carcinoma Surveillance and an Optimal Surveillance Interval: Nationwide Cohort of Korea. <i>Yonsei Medical Journal</i> , 2021 , 62, 758-766	3	1

17	MRI-diagnosis of category LR-M observations in the Liver Imaging Reporting and Data System v2018: a systematic review and meta-analysis.. <i>European Radiology</i> , 2022 , 1	8	o
16	Portal venous perfusion steal causing graft dysfunction after orthotopic liver transplantation: serial imaging findings in a successfully treated patient. <i>Ultrasonography</i> , 2016 , 35, 78-82	4.3	o
15	Subcentimeter hepatocellular carcinoma in treatment-naïve patients: noninvasive diagnostic criteria and tumor staging on gadoteric acid-enhanced MRI. <i>European Radiology</i> , 2021 , 31, 2321-2331	8	o
14	Hepatobiliary phase signal intensity: A potential method of diagnosing HCC with atypical imaging features among LR-M observations. <i>PLoS ONE</i> , 2021 , 16, e0257308	3.7	o
13	A Case Report of Cholangiocarcinoma Arising from a Biliary Adenofibroma: Radiologic Findings and Review of Literature. <i>Journal of the Korean Society of Radiology</i> , 2021 , 82, 721	0.2	o
12	Impact of Reference Standard on CT, MRI, and Contrast-enhanced US LI-RADS Diagnosis of Hepatocellular Carcinoma: A Meta-Analysis.. <i>Radiology</i> , 2022 , 212340	20.5	o
11	Extrasosseous Ewing's Sarcoma Presented as a Rectal Subepithelial Tumor: Radiological and Pathological Features 2017 , 21, 51		
10	Ferucarbotran-enhanced T2-weighted magnetic resonance imaging: differentiation of hepatic cavernous hemangiomas from malignant solid lesions. <i>Abdominal Imaging</i> , 2009 , 34, 494-501		
9	A growing solitary mass in the spleen. <i>Gastroenterology</i> , 2009 , 137, 1896, 2184	13.3	
8	A multi-institutional, single-arm, phase II trial of neoadjuvant modified-FOLFIRINOX for resectable pancreatic ductal adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2022 , 40, TPS624-TPS624	2.2	
7	Diffuse Large B-Cell Lymphoma in the Portal Vein. <i>Journal of the Korean Society of Radiology</i> , 2020 , 81, 707	0.2	
6	MRI Findings of Rectal Submucosal Tumors. <i>Korean Journal of Radiology</i> , 2011 , 12, 496	6.9	
5	Pancreatic Cystic Neoplasm: Radiologic Evaluation and Differential Diagnosis. <i>The Korean Journal of Pancreas and Biliary Tract</i> , 2019 , 24, 141-146	0.2	
4	Anal Metastasis Originating from Colorectal Cancer: Report of Two Cases. <i>Journal of the Korean Society of Radiology</i> , 2016 , 75, 501	0.2	
3	Gadoxetic Acid-Enhanced and Diffusion-Weighted Magnetic Resonance Imaging of Histologically Defined Early Hepatocellular Carcinoma. <i>Korean Journal of Abdominal Radiology</i> , 2021 , 5, 17-31		
2	Radiologic Evaluation for Resectability of Pancreatic Adenocarcinoma. <i>Journal of the Korean Society of Radiology</i> , 2021 , 82, 315	0.2	
1	A prospective study on the use of ultralow-dose computed tomography with iterative reconstruction for the follow-up of patients liver and renal abscess. <i>PLoS ONE</i> , 2021 , 16, e0246532	3.7	