

Mi-Suk Park

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/391132/mi-suk-park-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

196
papers

4,161
citations

34
h-index

54
g-index

202
ext. papers

4,910
ext. citations

6.3
avg, IF

5.18
L-index

#	Paper	IF	Citations
196	MRI-diagnosis of category LR-M observations in the Liver Imaging Reporting and Data System v2018: a systematic review and meta-analysis.. <i>European Radiology</i> , 2022 , 1	8	0
195	A multi-institutional, single-arm, phase II trial of neoadjuvant modified-FOLFIRINOX for resectable pancreatic ductal adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2022 , 40, TPS624-TPS624	2.2	
194	Retrospective Evaluation of Treatment Response in Patients with Nonmetastatic Pancreatic Cancer Using CT and CA 19-9.. <i>Radiology</i> , 2022 , 212236	20.5	2
193	Impact of Reference Standard on CT, MRI, and Contrast-enhanced US LI-RADS Diagnosis of Hepatocellular Carcinoma: A Meta-Analysis.. <i>Radiology</i> , 2022 , 212340	20.5	0
192	CT/MRI and CEUS LI-RADS Major Features Association with Hepatocellular Carcinoma: Individual Patient Data Meta-Analysis. <i>Radiology</i> , 2021 , 211244	20.5	4
191	Preoperative prediction of postsurgical outcomes in mass-forming intrahepatic cholangiocarcinoma based on clinical, radiologic, and radiomics features. <i>European Radiology</i> , 2021 , 31, 8638-8648	8	6
190	Diagnostic Performance of Liver Imaging Reporting and Data System Version 2017 Versus Version 2018 for Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis of Comparative Studies. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 1912-1919	5.6	2
189	Characteristics and Early Recurrence of Hepatocellular Carcinomas Categorized as LR-M: Comparison with Those Categorized as LR-4 or 5. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 1446-1454	5.6	2
188	Contrast-enhanced ultrasound Liver Imaging Reporting and Data System category M: a systematic review and meta-analysis. <i>Ultrasonography</i> , 2021 ,	4.3	1
187	LI-RADS treatment response algorithm for detecting incomplete necrosis in hepatocellular carcinoma after locoregional treatment: a systematic review and meta-analysis using individual patient data. <i>Abdominal Radiology</i> , 2021 , 46, 3717-3728	3	3
186	Gadoxetic Acid-Enhanced and Diffusion-Weighted Magnetic Resonance Imaging of Histologically Defined Early Hepatocellular Carcinoma. <i>Korean Journal of Abdominal Radiology</i> , 2021 , 5, 17-31		
185	Initial experience of irreversible electroporation for locally advanced pancreatic cancer in a Korean population. <i>Acta Radiologica</i> , 2021 , 62, 164-171	2	3
184	Liver MRI with amide proton transfer imaging: feasibility and accuracy for the characterization of focal liver lesions. <i>European Radiology</i> , 2021 , 31, 222-231	8	1
183	Prognostic factors in patients with locally advanced or borderline resectable pancreatic ductal adenocarcinoma: chemotherapy vs. chemoradiotherapy. <i>Abdominal Radiology</i> , 2021 , 46, 655-666	3	2
182	Subcentimeter hepatocellular carcinoma in treatment-naïve patients: noninvasive diagnostic criteria and tumor staging on gadoxetic acid-enhanced MRI. <i>European Radiology</i> , 2021 , 31, 2321-2331	8	0
181	Gadoxetic acid-enhanced MRI of macrotrabecular-massive hepatocellular carcinoma and its prognostic implications. <i>Journal of Hepatology</i> , 2021 , 74, 109-121	13.4	15
180	Extended application of subtraction arterial phase imaging in LI-RADS version 2018: a strategy to improve the diagnostic performance for hepatocellular carcinoma on gadoxetate disodium-enhanced MRI. <i>European Radiology</i> , 2021 , 31, 1620-1629	8	2

179	Radiologic Evaluation for Resectability of Pancreatic Adenocarcinoma. <i>Journal of the Korean Society of Radiology</i> , 2021 , 82, 315	0.2	
178	Should Threshold Growth Be Considered a Major Feature in the Diagnosis of Hepatocellular Carcinoma Using LI-RADS?. <i>Korean Journal of Radiology</i> , 2021 , 22, 1628-1639	6.9	1
177	Noninvasive evaluation of liver fibrosis: comparison of the stretched exponential diffusion-weighted model to other diffusion-weighted MRI models and transient elastography. <i>European Radiology</i> , 2021 , 31, 4813-4823	8	2
176	Effectiveness of Hepatocellular Carcinoma Surveillance and an Optimal Surveillance Interval: Nationwide Cohort of Korea. <i>Yonsei Medical Journal</i> , 2021 , 62, 758-766	3	1
175	Comparison of the current guidelines for diagnosing hepatocellular carcinoma using gadoxetic acid-enhanced magnetic resonance imaging. <i>European Radiology</i> , 2021 , 31, 4492-4503	8	4
174	LI-RADS Major Features on MRI for Diagnosing Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 518-525	5.6	4
173	A prospective study on the use of ultralow-dose computed tomography with iterative reconstruction for the follow-up of patients liver and renal abscess. <i>PLoS ONE</i> , 2021 , 16, e0246532	3.7	
172	Systematic review and meta-analysis of diagnostic performance of CT imaging for assessing resectability of pancreatic ductal adenocarcinoma after neoadjuvant therapy: importance of CT criteria. <i>Abdominal Radiology</i> , 2021 , 46, 5201-5217	3	2
171	Clinical Staging of Mass-Forming Intrahepatic Cholangiocarcinoma: Computed Tomography Versus Magnetic Resonance Imaging. <i>Hepatology Communications</i> , 2021 , 5, 2009-2018	6	3
170	Hepatobiliary phase signal intensity: A potential method of diagnosing HCC with atypical imaging features among LR-M observations. <i>PLoS ONE</i> , 2021 , 16, e0257308	3.7	0
169	A Case Report of Cholangiocarcinoma Arising from a Biliary Adenofibroma: Radiologic Findings and Review of Literature. <i>Journal of the Korean Society of Radiology</i> , 2021 , 82, 721	0.2	0
168	The Role of Imaging in Current Treatment Strategies for Pancreatic Adenocarcinoma. <i>Korean Journal of Radiology</i> , 2021 , 22, 23-40	6.9	12
167	Diagnostic performance of the LR-M criteria and spectrum of LI-RADS imaging features among primary hepatic carcinomas. <i>Abdominal Radiology</i> , 2020 , 45, 3743-3754	3	7
166	Diagnostic performance of Liver Imaging Reporting and Data System in patients at risk of both hepatocellular carcinoma and metastasis. <i>Abdominal Radiology</i> , 2020 , 45, 3789-3799	3	3
165	Is there association between statin usage and contrast-associated acute kidney injury after intravenous administration of iodine-based contrast media in enhanced computed tomography?. <i>European Radiology</i> , 2020 , 30, 5261-5271	8	3
164	Diagnostic Performance of CT/MRI Liver Imaging Reporting and Data System v2017 for Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis. <i>Liver International</i> , 2020 , 40, 1488-1497	7.9	21
163	Gadoxetic acid-enhanced MRI of hepatocellular carcinoma: Diagnostic performance of category-adjusted LR-5 using modified criteria. <i>PLoS ONE</i> , 2020 , 15, e0242344	3.7	5
162	The AFSUMB Consensus Statements and Recommendations for the Clinical Practice of Contrast-Enhanced Ultrasound using Sonazoid. <i>Ultrasonography</i> , 2020 , 39, 191-220	4.3	15

161	Diffuse Large B-Cell Lymphoma in the Portal Vein. <i>Journal of the Korean Society of Radiology</i> , 2020 , 81, 707	0.2	
160	FOLFIRINOX gemcitabine/nab-paclitaxel for treatment of metastatic pancreatic cancer: Single-center cohort study. <i>World Journal of Gastrointestinal Oncology</i> , 2020 , 12, 182-194	3.4	23
159	The Modified Response Evaluation Criteria in Solid Tumors (RECIST) Yield a More Accurate Prognoses Than the RECIST 1.1 in Hepatocellular Carcinoma Treated with Transarterial Radioembolization. <i>Gut and Liver</i> , 2020 , 14, 765-774	4.8	8
158	Retrospective comparison of EASL 2018 and LI-RADS 2018 for the noninvasive diagnosis of hepatocellular carcinoma using magnetic resonance imaging. <i>Hepatology International</i> , 2020 , 14, 70-79	8.8	22
157	MRI Ancillary Features for LI-RADS Category 3 and 4 Observations: Improved Categorization to Indicate the Risk of Hepatic Malignancy. <i>American Journal of Roentgenology</i> , 2020 , 215, 1354-1362	5.4	4
156	Radiographic portal or superior mesenteric vein invasion is an independent prognostic factor in non-metastatic pancreatic ductal adenocarcinoma: A missing block of clinical T staging?. <i>Pancreatology</i> , 2020 , 20, 952-959	3.8	1
155	Contrast-enhanced ultrasound liver imaging reporting and data system for diagnosing hepatocellular carcinoma: A meta-analysis. <i>Liver International</i> , 2020 , 40, 2345-2352	7.9	8
154	Evaluation of treatment response in hepatocellular carcinoma in the explanted liver with Liver Imaging Reporting and Data System version 2017. <i>European Radiology</i> , 2020 , 30, 261-271	8	34
153	The AFSUMB Consensus Statements and Recommendations for the Clinical Practice of Contrast-Enhanced Ultrasound using Sonazoid. <i>Journal of Medical Ultrasound</i> , 2020 , 28, 59-82	0.8	5
152	Metal implants influence CT scan parameters leading to increased local radiation exposure: A proposal for correction techniques. <i>PLoS ONE</i> , 2019 , 14, e0221692	3.7	1
151	Multimodality imaging studies of intraductal tubulopapillary neoplasms of the pancreas. <i>Diagnostic and Interventional Radiology</i> , 2019 , 25, 251-256	3.2	2
150	Optimal lexicon of gadoxetic acid-enhanced magnetic resonance imaging for the diagnosis of hepatocellular carcinoma modified from LI-RADS. <i>Abdominal Radiology</i> , 2019 , 44, 3078-3088	3	9
149	Characterization of focal liver lesions using the stretched exponential model: comparison with monoexponential and biexponential diffusion-weighted magnetic resonance imaging. <i>European Radiology</i> , 2019 , 29, 5111-5120	8	9
148	Optimal criteria for hepatocellular carcinoma diagnosis using CT in patients undergoing liver transplantation. <i>European Radiology</i> , 2019 , 29, 1022-1031	8	5
147	Contrast-enhanced US with Perfluorobutane for Hepatocellular Carcinoma Surveillance: A Multicenter Diagnostic Trial (SCAN). <i>Radiology</i> , 2019 , 292, 638-646	20.5	12
146	Hepatobiliary versus Extracellular MRI Contrast Agents in Hepatocellular Carcinoma Detection: Hepatobiliary Phase Features in Relation to Disease-free Survival. <i>Radiology</i> , 2019 , 293, 594-604	20.5	5
145	Preoperative Clinical and Computed Tomography (CT)-Based Nomogram to Predict Oncologic Outcomes in Patients with Pancreatic Head Cancer Resected with Curative Intent: A Retrospective Study. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	5
144	Gadoxetic acid-enhanced magnetic resonance imaging: Hepatocellular carcinoma and mimickers. <i>Clinical and Molecular Hepatology</i> , 2019 , 25, 223-233	6.9	21

143	Incremental Role of Pancreatic Magnetic Resonance Imaging after Staging Computed Tomography to Evaluate Patients with Pancreatic Ductal Adenocarcinoma. <i>Cancer Research and Treatment</i> , 2019 , 51, 24-33	5.2	10
142	Noninvasive Biomarker for Predicting Treatment Response to Concurrent Chemoradiotherapy in Patients with Hepatocellular Carcinoma 2019 , 23, 351		2
141	Pancreatic Cystic Neoplasm: Radiologic Evaluation and Differential Diagnosis. <i>The Korean Journal of Pancreas and Biliary Tract</i> , 2019 , 24, 141-146	0.2	
140	Imaging Features of Hepatocellular Carcinoma: Quantitative and Qualitative Comparison Between MRI-Enhanced With Gd-EOB-DTPA and Gd-DTPA. <i>Investigative Radiology</i> , 2019 , 54, 494-499	10.1	9
139	Radiological patterns of secondary sclerosing cholangitis in patients after lung transplantation. <i>Abdominal Radiology</i> , 2019 , 44, 1361-1366	3	1
138	Clinical Feasibility of MR Elastography in Patients With Biliary Obstruction. <i>American Journal of Roentgenology</i> , 2018 , 210, 1273-1278	5.4	8
137	Magnetic Resonance Imaging for Colorectal Cancer Metastasis to the Liver: Comparative Effectiveness Research for the Choice of Contrast Agents. <i>Cancer Research and Treatment</i> , 2018 , 50, 60-70	5.2	7
136	Feasibility of radiation dose reduction with iterative reconstruction in abdominopelvic CT for patients with inappropriate arm positioning. <i>PLoS ONE</i> , 2018 , 13, e0209754	3.7	6
135	CT features of hepatic metastases from hepatoid adenocarcinoma. <i>Abdominal Radiology</i> , 2017 , 42, 2402-2409	7	
134	Contrast-enhanced US with Perfluorobutane(Sonazoid) used as a surveillance test for Hepatocellular Carcinoma (HCC) in Cirrhosis (SCAN): an exploratory cross-sectional study for a diagnostic trial. <i>BMC Cancer</i> , 2017 , 17, 279	4.8	13
133	Rectal Mucinous Adenocarcinoma: MR Imaging Assessment of Response to Concurrent Chemotherapy and Radiation Therapy-A Hypothesis-generating Study. <i>Radiology</i> , 2017 , 285, 124-133	20.5	20
132	Feasibility of 3D navigator-triggered magnetic resonance cholangiopancreatography with combined parallel imaging and compressed sensing reconstruction at 3T. <i>Journal of Magnetic Resonance Imaging</i> , 2017 , 46, 1289-1297	5.6	33
131	Use of Imaging to Predict Complete Response of Colorectal Liver Metastases after Chemotherapy: MR Imaging versus CT Imaging. <i>Radiology</i> , 2017 , 284, 423-431	20.5	24
130	Extraosseous Ewing Sarcoma Presented as a Rectal Subepithelial Tumor: Radiological and Pathological Features 2017 , 21, 51		
129	Long-Term Follow-Up of Patients After Autologous Bone Marrow Cell Infusion for Decompensated Liver Cirrhosis. <i>Cell Transplantation</i> , 2017 , 26, 1059-1066	4	9
128	Added value of smooth hypointense rim in the hepatobiliary phase of gadoxetic acid-enhanced MRI in identifying tumour capsule and diagnosing hepatocellular carcinoma. <i>European Radiology</i> , 2017 , 27, 2610-2618	8	33
127	Efficacy and treatment-related adverse events of gemcitabine plus nab-paclitaxel for treatment of metastatic pancreatic cancer "in a Korean" population: A single-center cohort study. <i>Seminars in Oncology</i> , 2017 , 44, 420-427	5.5	12
126	A lexicon for hepatocellular carcinoma surveillance ultrasonography: benign versus malignant lesions. <i>Clinical and Molecular Hepatology</i> , 2017 , 23, 57-65	6.9	1

125	Lack of anti-tumor activity by anti-VEGF treatments in hepatic hemangiomas. <i>Angiogenesis</i> , 2016 , 19, 147-53	10.6	10
124	Dynamic contrast-enhanced MRI coupled with a subtraction technique is useful for treatment response evaluation of malignant melanoma hepatic metastasis. <i>Oncotarget</i> , 2016 , 7, 38513-38522	3.3	6
123	Magnetization-tagged MRI is a simple method for predicting liver fibrosis. <i>Clinical and Molecular Hepatology</i> , 2016 , 22, 140-5	6.9	1
122	Dynamic Contrast-Enhanced Magnetic Resonance Imaging as a Surrogate Biomarker for Bevacizumab in Colorectal Cancer Liver Metastasis: A Single-Arm, Exploratory Trial. <i>Cancer Research and Treatment</i> , 2016 , 48, 1210-1221	5.2	9
121	Portal venous perfusion steal causing graft dysfunction after orthotopic liver transplantation: serial imaging findings in a successfully treated patient. <i>Ultrasonography</i> , 2016 , 35, 78-82	4.3	0
120	Anal Metastasis Originating from Colorectal Cancer: Report of Two Cases. <i>Journal of the Korean Society of Radiology</i> , 2016 , 75, 501	0.2	
119	Antiangiogenic Therapy Induces Hepatic Tumor Vascular Network Rearrangement to Receive Perfusion via the Portal Vein and Hepatic Artery. <i>Journal of Vascular Research</i> , 2016 , 53, 72-82	1.9	1
118	Comparison of MR imaging features of solid pseudopapillary neoplasm of pancreas between male and female patients. <i>European Journal of Radiology</i> , 2015 , 84, 2065-70	4.7	14
117	Quantitative analysis of the effect of iterative reconstruction using a phantom: determining the appropriate blending percentage. <i>Yonsei Medical Journal</i> , 2015 , 56, 253-61	3	19
116	MRI Findings and Prediction of Time to Progression of Patients with Hepatocellular Carcinoma Treated with Drug-eluting Bead Transcatheter Arterial Chemoembolization. <i>Journal of Korean Medical Science</i> , 2015 , 30, 965-73	4.7	10
115	Pain palliation in patients with bone metastases using magnetic resonance-guided focused ultrasound with conformal bone system: a preliminary report. <i>Yonsei Medical Journal</i> , 2015 , 56, 503-9	3	28
114	Inter-observer variability of response evaluation criteria for hepatocellular carcinoma treated with chemoembolization. <i>Digestive and Liver Disease</i> , 2015 , 47, 682-8	3.3	5
113	Complete response at first chemoembolization is still the most robust predictor for favorable outcome in hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2015 , 62, 1304-10	13.4	108
112	Mouse Hepatic Tumor Vascular Imaging by Experimental Selective Angiography. <i>PLoS ONE</i> , 2015 , 10, e0131687	3.7	3
111	Is Non-Contrast CT Adequate for the Evaluation of Hepatic Metastasis in Patients Who Cannot Receive Iodinated Contrast Media?. <i>PLoS ONE</i> , 2015 , 10, e0134133	3.7	10
110	Necrotic lymphoma in a patient with post-transplantation lymphoproliferative disorder: ultrasonography and CT findings with pathologic correlation. <i>Ultrasonography</i> , 2015 , 34, 148-52	4.3	1
109	Growth rate of early-stage hepatocellular carcinoma in patients with chronic liver disease. <i>Clinical and Molecular Hepatology</i> , 2015 , 21, 279-86	6.9	53
108	Diffusion and perfusion MRI prediction of progression-free survival in patients with hepatocellular carcinoma treated with concurrent chemoradiotherapy. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 39, 286-92	5.6	17

107	Dynamic enhancement pattern of HCC smaller than 3 cm in diameter on gadoteric acid-enhanced MRI: comparison with multiphasic MDCT. <i>Liver International</i> , 2014 , 34, 1593-602	7.9	20
106	Correlation between 18F-fluorodeoxyglucose positron emission tomography and pathologic differentiation in pancreatic cancer. <i>Annals of Nuclear Medicine</i> , 2014 , 28, 430-5	2.5	22
105	Liver stiffness assessment with tagged MRI of cardiac-induced liver motion in cirrhosis patients. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 39, 1301-7	5.6	16
104	Prediction of postoperative pancreatic fistulas after pancreatectomy: assessment with acoustic radiation force impulse elastography. <i>Journal of Ultrasound in Medicine</i> , 2014 , 33, 781-6	2.9	20
103	Intraindividual comparison of diagnostic performance in patients with hepatic metastasis of full-dose standard and half-dose iterative reconstructions with dual-source abdominal computed tomography. <i>Investigative Radiology</i> , 2014 , 49, 195-200	10.1	25
102	Detection of recurrent hepatocellular carcinoma on post-operative surveillance: comparison of MDCT and gadoteric acid-enhanced MRI. <i>Abdominal Imaging</i> , 2014 , 39, 291-9		19
101	Added value of subtraction imaging in detecting arterial enhancement in small (. <i>European Radiology</i> , 2013 , 23, 924-30	8	38
100	Comparison of breathhold, navigator-triggered, and free-breathing diffusion-weighted MRI for focal hepatic lesions. <i>Journal of Magnetic Resonance Imaging</i> , 2013 , 38, 109-18	5.6	52
99	Radiation dose reduction via sinogram affirmed iterative reconstruction and automatic tube voltage modulation (CARE kV) in abdominal CT. <i>Korean Journal of Radiology</i> , 2013 , 14, 886-93	6.9	24
98	Characterization of incidental liver lesions: comparison of multidetector CT versus Gd-EOB-DTPA-enhanced MR imaging. <i>PLoS ONE</i> , 2013 , 8, e66141	3.7	27
97	¹⁸ F-FDG PET metabolic parameters and MRI perfusion and diffusion parameters in hepatocellular carcinoma: a preliminary study. <i>PLoS ONE</i> , 2013 , 8, e71571	3.7	14
96	Correlations of dynamic contrast-enhanced magnetic resonance imaging with morphologic, angiogenic, and molecular prognostic factors in rectal cancer. <i>Yonsei Medical Journal</i> , 2013 , 54, 123-30	3	25
95	Preoperative prediction of the microvascular invasion of hepatocellular carcinoma with diffusion-weighted imaging. <i>Liver Transplantation</i> , 2012 , 18, 1171-8	4.5	71
94	Liver trauma diagnosis with contrast-enhanced ultrasound: interobserver variability between radiologist and emergency physician in an animal study. <i>American Journal of Emergency Medicine</i> , 2012 , 30, 1229-34	2.9	6
93	Detection of liver metastases using gadoteric-enhanced dynamic and 10- and 20-minute delayed phase MR imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 35, 635-43	5.6	31
92	Quantitative evaluation of liver cirrhosis using T1 relaxation time with 3 tesla MRI before and after oxygen inhalation. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 36, 405-10	5.6	39
91	Distal common bile duct metastasis from hepatocellular carcinoma. <i>Hepatology</i> , 2012 , 55, 1638-9	11.2	4
90	Hepatocellular carcinoma: detection with diffusion-weighted versus contrast-enhanced magnetic resonance imaging in pretransplant patients. <i>Hepatology</i> , 2012 , 56, 140-8	11.2	87

89	Hepatocellular carcinoma: clinical and radiological findings in patients with chronic B viral hepatitis and chronic C viral hepatitis. <i>Abdominal Imaging</i> , 2012 , 37, 591-4		8
88	Accuracy of gadoxetic acid-enhanced magnetic resonance imaging for the diagnosis of sinusoidal obstruction syndrome in patients with chemotherapy-treated colorectal liver metastases. <i>European Radiology</i> , 2012 , 22, 864-71	8	85
87	Prediction of the histopathological grade of hepatocellular carcinoma using qualitative diffusion-weighted, dynamic, and hepatobiliary phase MRI. <i>European Radiology</i> , 2012 , 22, 1701-8	8	45
86	Response evaluation in patients with colorectal liver metastases: RECIST version 1.1 versus modified CT criteria. <i>American Journal of Roentgenology</i> , 2012 , 199, 809-15	5-4	60
85	The impact of CT follow-up interval on stages of hepatocellular carcinomas detected during the surveillance of patients with liver cirrhosis. <i>American Journal of Roentgenology</i> , 2012 , 199, 816-21	5-4	3
84	Enhancement patterns of hepatocellular carcinoma after transarterial chemoembolization using drug-eluting beads on arterial phase CT images: a pilot retrospective study. <i>American Journal of Roentgenology</i> , 2012 , 199, 349-59	5-4	19
83	Malignant Mixed Müllerian Tumor with Small Bowel Metastasis: A Case Report. <i>Journal of the Korean Society of Magnetic Resonance in Medicine</i> , 2012 , 16, 257		2
82	Differentiation of benign and malignant ampullary obstructions on MR imaging. <i>European Journal of Radiology</i> , 2011 , 80, 198-203	4-7	27
81	Using multi-detector-row CT to diagnose ampullary adenoma or adenocarcinoma in situ. <i>European Journal of Radiology</i> , 2011 , 80, e340-5	4-7	10
80	MRI findings of rectal submucosal tumors. <i>Korean Journal of Radiology</i> , 2011 , 12, 487-98	6-9	19
79	Pancreatic tumors: emphasis on CT findings and pathologic classification. <i>Korean Journal of Radiology</i> , 2011 , 12, 731-9	6-9	28
78	Prognostic factors and 10-year survival in patients with hepatocellular carcinoma after curative hepatectomy. <i>Journal of Gastrointestinal Surgery</i> , 2011 , 15, 598-607	3-3	26
77	Comparison of gadoxetic acid-enhanced dynamic imaging and diffusion-weighted imaging for the preoperative evaluation of colorectal liver metastases. <i>Journal of Magnetic Resonance Imaging</i> , 2011 , 34, 345-53	5-6	77
76	Differentiation of hepatic hyperintense lesions seen on gadoxetic acid-enhanced hepatobiliary phase MRI. <i>American Journal of Roentgenology</i> , 2011 , 197, W44-52	5-4	64
75	MRI Findings of Rectal Submucosal Tumors. <i>Korean Journal of Radiology</i> , 2011 , 12, 496		6-9
74	Intraductal ultrasonography combined with percutaneous transhepatic cholangioscopy for the preoperative evaluation of longitudinal tumor extent in hilar cholangiocarcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2010 , 25, 286-92	4	22
73	Differential features of pancreatobiliary- and intestinal-type ampullary carcinomas at MR imaging. <i>Radiology</i> , 2010 , 257, 384-93	20-5	21
72	Optimal T2-weighted MR cholangiopancreatographic images can be obtained after administration of gadoxetic acid. <i>Radiology</i> , 2010 , 256, 475-84	20-5	18

71	Comparison of MRI and endoscopic ultrasound in the characterization of pancreatic cystic lesions. <i>American Journal of Roentgenology</i> , 2010 , 195, 947-52	5.4	69
70	Autologous bone marrow infusion activates the progenitor cell compartment in patients with advanced liver cirrhosis. <i>Cell Transplantation</i> , 2010 , 19, 1237-46	4	71
69	Potential conditions causing impairment of selective hepatobiliary enhancement of gadobenate dimeglumine-enhanced delayed magnetic resonance imaging. <i>Journal of Computer Assisted Tomography</i> , 2010 , 34, 113-20	2.2	7
68	Diffusion-weighted MR imaging of liver on 3.0-Tesla system: effect of intravenous administration of gadoxetic acid disodium. <i>European Radiology</i> , 2010 , 20, 1052-60	8	90
67	Characterisation of small hypoattenuating hepatic lesions in multi-detector CT (MDCT) in patients with underlying extrahepatic malignancy: added value of contrast-enhanced MR images. <i>European Radiology</i> , 2010 , 20, 2853-61	8	2
66	Primary hepatic schwannoma mimicking malignancy on fluorine-18 2-fluoro-2-deoxy-D-glucose positron emission tomography-computed tomography. <i>Hepatology</i> , 2010 , 51, 1080-1	11.2	10
65	Quantification of superparamagnetic iron oxide-mediated signal intensity change in patients with liver cirrhosis using T2 and T2* mapping: a preliminary report. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 31, 1379-86	5.6	6
64	Preoperative radiologic and postoperative pathologic risk factors for early intra-hepatic recurrence in hepatocellular carcinoma patients who underwent curative resection. <i>Yonsei Medical Journal</i> , 2009 , 50, 789-95	3	22
63	Hepatocellular carcinoma variants: radiologic-pathologic correlation. <i>American Journal of Roentgenology</i> , 2009 , 193, W7-13	5.4	34
62	Effects of neoadjuvant combined chemotherapy and radiation therapy on the CT evaluation of resectability and staging in patients with pancreatic head cancer. <i>Radiology</i> , 2009 , 250, 758-65	20.5	62
61	Perfusion CT: noninvasive surrogate marker for stratification of pancreatic cancer response to concurrent chemo- and radiation therapy. <i>Radiology</i> , 2009 , 250, 110-7	20.5	123
60	Ferucarbotran-enhanced T2-weighted magnetic resonance imaging: differentiation of hepatic cavernous hemangiomas from malignant solid lesions. <i>Abdominal Imaging</i> , 2009 , 34, 494-501		
59	Role of EUS and MDCT in the diagnosis of gastric submucosal tumors according to the revised pathologic concept of gastrointestinal stromal tumors. <i>European Radiology</i> , 2009 , 19, 924-34	8	24
58	Can microvessel invasion of hepatocellular carcinoma be predicted by pre-operative MRI?. <i>European Radiology</i> , 2009 , 19, 1744-51	8	118
57	A growing solitary mass in the spleen. <i>Gastroenterology</i> , 2009 , 137, 1896, 2184	13.3	
56	Metastasis versus focal eosinophilic infiltration of the liver in patients with extrahepatic abdominal cancer: an evaluation with gadobenate dimeglumine-enhanced magnetic resonance imaging. <i>Journal of Computer Assisted Tomography</i> , 2009 , 33, 119-24	2.2	15
55	Paraortic lymph node metastasis in patients with intra-abdominal malignancies: CT vs PET. <i>World Journal of Gastroenterology</i> , 2009 , 15, 4434-8	5.6	17
54	Electronic clinical challenges and images in GI. Hepatic venoocclusive disease. <i>Gastroenterology</i> , 2008 , 135, e3-4	13.3	1

53	Peritoneal mesotheliomas: clinicopathologic features, CT findings, and differential diagnosis. <i>American Journal of Roentgenology</i> , 2008 , 191, 814-25	5-4	64
52	MR cholangiography for evaluation of hilar branching anatomy in transplantation of the right hepatic lobe from a living donor. <i>American Journal of Roentgenology</i> , 2008 , 191, 537-45	5-4	28
51	Choledochal cyst and anomalous pancreaticobiliary ductal union in adults: radiological spectrum and complications. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 17-22	2-2	21
50	Radiological and clinical features of sarcomatoid hepatocellular carcinoma in 11 cases. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 745-9	2-2	33
49	Hepatic splenosis preoperatively diagnosed as hepatocellular carcinoma in a patient with chronic hepatitis B: a case report. <i>Journal of Korean Medical Science</i> , 2008 , 23, 336-41	4-7	26
48	Intestinal parasitic infection. <i>Abdominal Imaging</i> , 2008 , 33, 166-71		20
47	Double common bile duct: curved-planar reformatted computed tomography (CT) and gadobenate dimeglumine-enhanced MR cholangiography. <i>Journal of Magnetic Resonance Imaging</i> , 2008 , 27, 209-11	5-6	7
46	Lymphoma presenting as an echogenic periportal mass: sonographic findings. <i>Journal of Clinical Ultrasound</i> , 2008 , 36, 437-9	1	7
45	Comparison of CT and MRI for presurgical characterization of paraaortic lymph nodes in patients with pancreatico-biliary carcinoma. <i>World Journal of Gastroenterology</i> , 2008 , 14, 2208-12	5-6	12
44	Spontaneous regression of a cystic tumor in a postpartum woman; is it a cystic lymphangioma?. <i>Yonsei Medical Journal</i> , 2007 , 48, 715-8	3	10
43	Value of manganese-enhanced T1- and T2-weighted MR cholangiography for differentiating cystic parenchymal lesions from cystic abnormalities which communicate with bile ducts. <i>Yonsei Medical Journal</i> , 2007 , 48, 1072-4	3	7
42	Solitary fibrous tumor arising from stomach: CT findings. <i>Yonsei Medical Journal</i> , 2007 , 48, 1056-60	3	10
41	Gadobenate dimeglumine-enhanced MRI of intraductal papillary mucinous tumor of the bile ducts. <i>Journal of Magnetic Resonance Imaging</i> , 2007 , 25, 625-7	5-6	9
40	Imaging-guided minimally invasive laparoscopic resection of intraluminal small-bowel tumor: report of two cases. <i>American Journal of Roentgenology</i> , 2007 , 189, 56-60	5-4	11
39	MR cholangiography of accessory bile duct connected to the stomach. <i>American Journal of Roentgenology</i> , 2007 , 189, W344-7	5-4	9
38	Hepatic MRI using the double-echo chemical shift phase-selective gradient-echo technique. <i>American Journal of Roentgenology</i> , 2007 , 188, W49-56	5-4	9
37	Value of "patent track" sign on Doppler sonography after percutaneous liver biopsy in detection of postbiopsy bleeding: a prospective study in 352 patients. <i>American Journal of Roentgenology</i> , 2007 , 189, 109-16	5-4	29
36	Imaging findings of leukemic involvement of the pancreaticobiliary system in adults. <i>American Journal of Roentgenology</i> , 2007 , 188, 1589-95	5-4	15

35	Diagnostic accuracy of multidetector-row computed tomography for common bile duct calculi: is it necessary to add non-contrast-enhanced images to contrast-enhanced images?. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 508-12	2.2	5
34	Biliary ductal involvement of hilar cholangiocarcinoma: multidetector computed tomography versus magnetic resonance cholangiography. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 72-8	2.2	41
33	Gastric true leiomyoma: computed tomographic findings and pathological correlation. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 204-8	2.2	34
32	FDG uptake in PET by bladder hernia simulating inguinal metastasis. <i>Yonsei Medical Journal</i> , 2007 , 48, 886-90	3	7
31	Non-contiguous Multi-organ Involvement of an Inflammatory Myofibroblastic Tumor: A Case Report. <i>Journal of the Korean Radiological Society</i> , 2007 , 57, 265		1
30	Primary Malignant Melanoma of the Esophagus: A Case Report. <i>Journal of the Korean Radiological Society</i> , 2007 , 57, 37		1
29	Hyperintense nodules on non-enhanced T1-weighted gradient-echo magnetic resonance imaging of cirrhotic liver: fate and clinical implications. <i>Journal of Magnetic Resonance Imaging</i> , 2006 , 24, 630-6	5.6	13
28	Hepatic hemangiomas with arterioportal shunt: sonographic appearances with CT and MRI correlation. <i>American Journal of Roentgenology</i> , 2006 , 187, W406-14	5.4	32
27	CT and PET in stomach cancer: preoperative staging and monitoring of response to therapy. <i>Radiographics</i> , 2006 , 26, 143-56	5.4	133
26	Comparison of CT and 18F-FDG pet for detecting peritoneal metastasis on the preoperative evaluation for gastric carcinoma. <i>Korean Journal of Radiology</i> , 2006 , 7, 249-56	6.9	72
25	Preoperative staging accuracy of multidetector row computed tomography for extrahepatic bile duct carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 362-7	2.2	26
24	Advanced gastric carcinoma with signet ring cell carcinoma versus non-signet ring cell carcinoma: differentiation with multidetector CT. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 880-4	2.2	22
23	Imaging findings of biliary and nonbiliary complications following laparoscopic surgery. <i>European Radiology</i> , 2006 , 16, 1906-14	8	4
22	Focal eosinophilic necrosis versus metastasis in the liver: the usefulness of two-phase dynamic CT. <i>American Journal of Roentgenology</i> , 2005 , 184, 1085-90	5.4	12
21	Eosinophilic hepatic necrosis: magnetic resonance imaging and computed tomography comparison. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 765-71	2.2	16
20	Unusual cystic neoplasms in the pancreas: radiologic-pathologic correlation. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 610-6	2.2	18
19	Capecitabine combined with gemcitabine (CapGem) as first-line treatment in patients with advanced/metastatic biliary tract carcinoma. <i>Cancer</i> , 2005 , 104, 2753-8	6.4	86
18	A Phase II study of capecitabine combined with gemcitabine in patients with advanced gallbladder carcinoma. <i>Yonsei Medical Journal</i> , 2005 , 46, 526-31	3	14

17	Abdominal sparganosis presenting as an abscess with fistulous communication to the bowel. <i>American Journal of Roentgenology</i> , 2005 , 185, 1084-5	5.4	13
16	Preoperative MRI of potential living-donor-related liver transplantation using a single dose of gadobenate dimeglumine. <i>American Journal of Roentgenology</i> , 2005 , 185, 424-31	5.4	30
15	Abdominal cocoon: preoperative diagnostic clues from radiologic imaging with pathologic correlation. <i>American Journal of Roentgenology</i> , 2004 , 182, 639-41	5.4	66
14	Scirrhus gastric carcinoma: endoscopy versus upper gastrointestinal radiography. <i>Radiology</i> , 2004 , 231, 421-6	20.5	26
13	Early biliary complications of laparoscopic cholecystectomy: evaluation on T2-weighted MR cholangiography in conjunction with mangafodipir trisodium-enhanced 3D T1-weighted MR cholangiography. <i>American Journal of Roentgenology</i> , 2004 , 183, 1559-66	5.4	38
12	Primary malignant melanoma of the rectum: CT findings in eight patients. <i>Radiology</i> , 2004 , 232, 181-6	20.5	35
11	Small (. <i>American Journal of Roentgenology</i> , 2004 , 182, 1233-40	5.4	38
10	Differentiation of extrahepatic bile duct cholangiocarcinoma from benign stricture: findings at MRCP versus ERCP. <i>Radiology</i> , 2004 , 233, 234-40	20.5	203
9	Acute cholecystitis at T2-weighted and manganese-enhanced T1-weighted MR cholangiography: preliminary study. <i>Radiology</i> , 2003 , 227, 580-4	20.5	45
8	Predisposing factors of bile duct injury after transcatheter arterial chemoembolization (TACE) for hepatic malignancy. <i>CardioVascular and Interventional Radiology</i> , 2002 , 25, 270-4	2.7	59
7	Mucinous versus nonmucinous gastric carcinoma: differentiation with helical CT. <i>Radiology</i> , 2002 , 223, 540-6	20.5	30
6	Radiographic findings of primary B-cell lymphoma of the stomach: low-grade versus high-grade malignancy in relation to the mucosa-associated lymphoid tissue concept. <i>American Journal of Roentgenology</i> , 2002 , 179, 1297-304	5.4	31
5	Transient peritumoral enhancement during dynamic MRI of the liver: cavernous hemangioma versus hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 411-7	2.2	19
4	Radiologic findings of gastrointestinal tract involvement in hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 95-101	2.2	34
3	Recurrent pyogenic cholangitis: comparison between MR cholangiography and direct cholangiography. <i>Radiology</i> , 2001 , 220, 677-82	20.5	50
2	Bile duct injuries leading to portal vein obliteration after transcatheter arterial chemoembolization in the liver: CT findings and initial observations. <i>Radiology</i> , 2001 , 221, 429-36	20.5	33
1	Hepatic cavernous hemangioma in cirrhotic liver: imaging findings. <i>Korean Journal of Radiology</i> , 2000 , 1, 185-90	6.9	11