

Whady Hueb

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3911920/whady-hueb-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

126
papers

6,446
citations

30
h-index

79
g-index

174
ext. papers

8,052
ext. citations

5.7
avg, IF

4.86
L-index

#	Paper	IF	Citations
126	Death and Myocardial Infarction Following Initial Revascularization Versus Optimal Medical Therapy in Chronic Coronary Syndromes With Myocardial Ischemia: A Systematic Review and Meta-Analysis of Contemporary Randomized Controlled Trials. <i>Journal of the American Heart Association</i> , 2021, 10, e019114	6	5
125	Surgical and percutaneous revascularization outcomes based on SYNTAX I, II, and residual scores: a long-term follow-up study. <i>Journal of Cardiothoracic Surgery</i> , 2021, 16, 248	1.6	0
124	Initial Invasive or Conservative Strategy for Stable Coronary Disease. <i>New England Journal of Medicine</i> , 2020, 382, 1395-1407	59.2	642
123	Significant association of SYNTAX score on release of cardiac biomarkers in uncomplicated post-revascularization procedures among patients with stable multivessel disease: MASS-V Study group. <i>Medicine (United States)</i> , 2020, 99, e18973	1.8	1
122	Association of Longitudinal Values of Glycated Hemoglobin With Cardiovascular Events in Patients With Type 2 Diabetes and Multivessel Coronary Artery Disease. <i>JAMA Network Open</i> , 2020, 3, e1919666	10.4	4
121	Stress Testing and Risk Prediction in People With Known Symptomatic Multivessel Coronary Artery Disease-Reply. <i>JAMA Internal Medicine</i> , 2020, 180, 166-167	11.5	
120	Long-term prognostic value of late gadolinium enhancement and periprocedural myocardial infarction after uncomplicated revascularization: MASS-V follow-up. <i>European Heart Journal Cardiovascular Imaging</i> , 2020,	4.1	1
119	Long-term outcomes of patients with stable coronary disease and chronic kidney dysfunction: 10-year follow-up of the Medicine, Angioplasty, or Surgery Study II Trial. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 1369-1376	4.3	6
118	Lipid transfer to HDL, CETP and HDL composition in coronary artery disease patients with or without type 2 diabetes mellitus. <i>European Journal of Preventive Cardiology</i> , 2020, 27, 2223-2225	3.9	
117	Hypotheses, rationale, design, and methods for prognostic evaluation of a randomized comparison between patients with coronary artery disease associated with ischemic cardiomyopathy who undergo medical or surgical treatment: MASS-VI (HF). <i>Trials</i> , 2020, 21, 337	2.8	0
116	Clinical significance of chronic myocardial ischemia in coronary artery disease patients. <i>Journal of Thoracic Disease</i> , 2019, 11, 1005-1015	2.6	20
115	Association Between Stress Testing-Induced Myocardial Ischemia and Clinical Events in Patients With Multivessel Coronary Artery Disease. <i>JAMA Internal Medicine</i> , 2019, 179, 1345-1351	11.5	8
114	Large Bilateral Coronary Artery Fistula: 10-year Follow-up in Clinical Treatment. <i>Arquivos Brasileiros De Cardiologia</i> , 2019, 112, 211-213	1.2	
113	Effect of ischemic preconditioning on cardiovascular outcomes in patients with symptomatic coronary artery disease: a cohort study. <i>Coronary Artery Disease</i> , 2019, 30, 536-541	1.4	
112	Long-Term Survival Following Multivessel Revascularization in Patients With Diabetes: The FREEDOM Follow-On Study. <i>Journal of the American College of Cardiology</i> , 2019, 73, 629-638	15.1	109
111	Ten-Year Follow-Up of Off-Pump and On-Pump Multivessel Coronary Artery Bypass Grafting: MASS III. <i>Angiology</i> , 2019, 70, 337-344	2.1	4
110	Mortality after coronary artery bypass grafting versus percutaneous coronary intervention with stenting for coronary artery disease: a pooled analysis of individual patient data. <i>Lancet, The</i> , 2018, 391, 939-948	40	290

109	Stroke Rates Following Surgical Versus Percutaneous Coronary Revascularization. <i>Journal of the American College of Cardiology</i> , 2018 , 72, 386-398	15.1	59
108	Cost-effectiveness of on-pump and off-pump coronary artery bypass grafting for patients with coronary artery disease: Results from the MASS III trial. <i>International Journal of Cardiology</i> , 2018 , 273, 63-68	3.2	1
107	Behavior of Ultrasensitive C-Reactive Protein in Myocardial Revascularization with and without Cardiopulmonary Bypass. <i>Brazilian Journal of Cardiovascular Surgery</i> , 2018 , 33, 535-541	1.1	
106	Chronic Kidney Disease and Coronary Artery Disease 2018 ,		1
105	Comparative cost-effectiveness of surgery, angioplasty, or medical therapy in patients with multivessel coronary artery disease: MASS II trial. <i>Cost Effectiveness and Resource Allocation</i> , 2018 , 16, 55	2.4	4
104	SYNTAX Score in Patients With Diabetes Undergoing Coronary Revascularization in the FREEDOM Trial. <i>Journal of the American College of Cardiology</i> , 2018 , 72, 2826-2837	15.1	29
103	SYNTAX Score and Long-Term Outcomes: The BARI-2D Trial. <i>Journal of the American College of Cardiology</i> , 2017 , 69, 395-403	15.1	38
102	Biomarker release after percutaneous coronary intervention in patients without established myocardial infarction as assessed by cardiac magnetic resonance with late gadolinium enhancement. <i>Catheterization and Cardiovascular Interventions</i> , 2017 , 90, 87-93	2.7	3
101	Abnormal elevation of myocardial necrosis biomarkers after coronary artery bypass grafting without established myocardial infarction assessed by cardiac magnetic resonance. <i>Journal of Cardiothoracic Surgery</i> , 2017 , 12, 122	1.6	2
100	Myocardial injury in diabetic patients with multivessel coronary artery disease after revascularization interventions. <i>Diabetology and Metabolic Syndrome</i> , 2017 , 9, 92	5.6	2
99	Significant elevation of biomarkers of myocardial necrosis after coronary artery bypass grafting without myocardial infarction established assessed by cardiac magnetic resonance. <i>Medicine (United States)</i> , 2017 , 96, e6053	1.8	2
98	Utility and quality-adjusted life-years in coronary artery disease: Five-year follow-up of the MASS II trial. <i>Medicine (United States)</i> , 2017 , 96, e9113	1.8	9
97	Reduced risk of myocardial infarct and revascularization following coronary artery bypass grafting compared with percutaneous coronary intervention in patients with chronic kidney disease. <i>Kidney International</i> , 2016 , 90, 411-421	9.9	25
96	Impact of Chronic Kidney Disease on Long-Term Outcomes in Type 2 Diabetic Patients With Coronary Artery Disease on Surgical, Angioplasty, or Medical Treatment. <i>Annals of Thoracic Surgery</i> , 2016 , 101, 1735-44	2.7	13
95	Accuracy of Myocardial Biomarkers in the Diagnosis of Myocardial Infarction After Revascularization as Assessed by Cardiac Resonance: The Medicine, Angioplasty, Surgery Study V (MASS-V) Trial. <i>Annals of Thoracic Surgery</i> , 2016 , 101, 2202-8	2.7	13
94	The challenge of treating elderly coronary artery disease patients. <i>Journal of Thoracic Disease</i> , 2016 , 8, 1434-6	2.6	2
93	The Influence of Diabetes Mellitus in Myocardial Ischemic Preconditioning. <i>Journal of Diabetes Research</i> , 2016 , 2016, 8963403	3.9	2
92	Effects of glycemic control upon serum lipids and lipid transfers to HDL in patients with type 2 diabetes mellitus: novel findings in unesterified cholesterol status. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015 , 123, 232-9	2.3	8

91	Recurrent angina caused by coronary subclavian steal syndrome confirmed by positron emission tomography. <i>Annals of Thoracic Surgery</i> , 2015 , 99, e111-4	2.7	2
90	Troponin in diabetic patients with and without chronic coronary artery disease. <i>BMC Cardiovascular Disorders</i> , 2015 , 15, 72	2.3	21
89	Comparison between off-pump and on-pump coronary artery bypass grafting in patients with severe lesions at the circumflex artery territory: 5-year follow-up of the MASS III trial. <i>European Journal of Cardio-thoracic Surgery</i> , 2015 , 47, 455-8	3	6
88	Role of Trimetazidine in Ischemic Preconditioning in Patients With Symptomatic Coronary Artery Disease. <i>Medicine (United States)</i> , 2015 , 94, e1161	1.8	4
87	Alterations in lipid transfers to HDL associated with the presence of coronary artery disease in patients with type 2 diabetes mellitus. <i>Cardiovascular Diabetology</i> , 2015 , 14, 107	8.7	24
86	Type 2 diabetes mellitus and myocardial ischemic preconditioning in symptomatic coronary artery disease patients. <i>Cardiovascular Diabetology</i> , 2015 , 14, 66	8.7	15
85	Conservative strategy for treatment of stable coronary artery disease. <i>World Journal of Clinical Cases</i> , 2015 , 3, 163-70	1.6	2
84	On-pump versus off-pump coronary artery bypass surgery in patients older than 60 years: five-year follow-up of MASS III trial. <i>Journal of Cardiothoracic Surgery</i> , 2014 , 9,	1.6	3
83	On-pump versus off-pump coronary artery bypass surgery in patients older than 60 years: five-year follow-up of MASS III trial. <i>Journal of Cardiothoracic Surgery</i> , 2014 , 9, 127	1.6	3
82	A case of mid-apical obstructive hypertrophic cardiomyopathy treated with a transapical myectomy approach: a case report. <i>Journal of Medical Case Reports</i> , 2014 , 8, 364	1.2	1
81	The cost-effectiveness of strategies in coronary artery disease. <i>Expert Review of Pharmacoeconomics and Outcomes Research</i> , 2014 , 14, 805-13	2.2	3
80	Percutaneous coronary intervention outcomes in patients with stable obstructive coronary artery disease and myocardial ischemia: a collaborative meta-analysis of contemporary randomized clinical trials. <i>JAMA Internal Medicine</i> , 2014 , 174, 232-40	11.5	166
79	Long-term follow-up of a randomized, controlled clinical trial of three therapeutic strategies for multivessel stable coronary artery disease in women. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2014 , 19, 997-1001	1.8	1
78	Impact of hypoglycemic agents on myocardial ischemic preconditioning. <i>World Journal of Diabetes</i> , 2014 , 5, 258-66	4.7	9
77	Association between UCP2 A55V polymorphism and risk of cardiovascular events in patients with multi-vessel coronary arterial disease. <i>BMC Medical Genetics</i> , 2013 , 14, 40	2.1	13
76	Impact of diabetes on 10-year outcomes of patients with multivessel coronary artery disease in the Medicine, Angioplasty, or Surgery Study II (MASS II) trial. <i>American Heart Journal</i> , 2013 , 166, 250-7	4.9	45
75	Ten-year outcomes of patients randomized to surgery, angioplasty, or medical treatment for stable multivessel coronary disease: effect of age in the Medicine, Angioplasty, or Surgery Study II trial. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2013 , 146, 1105-12	1.5	10
74	Hypotheses, rationale, design, and methods for evaluation of ischemic preconditioning assessed by sequential exercise tests in diabetic and non-diabetic patients with stable coronary artery disease--a prospective study. <i>BMC Cardiovascular Disorders</i> , 2013 , 13, 117	2.3	4

73	Long-term analysis of left ventricular ejection fraction in patients with stable multivessel coronary disease undergoing medicine, angioplasty or surgery: 10-year follow-up of the MASS II trial. <i>European Heart Journal</i> , 2013 , 34, 3370-7	9.5	9
72	Effect of hypoglycemic agents on ischemic preconditioning in patients with type 2 diabetes and symptomatic coronary artery disease. <i>Diabetes Care</i> , 2013 , 36, 1654-9	14.6	25
71	Effect of complete revascularization on 10-year survival of patients with stable multivessel coronary artery disease: MASS II trial. <i>Circulation</i> , 2012 , 126, S158-63	16.7	44
70	Higher incidence of death in multi-vessel coronary artery disease patients associated with polymorphisms in chromosome 9p21. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 61	2.3	15
69	Hypotheses, rationale, design, and methods for prognostic evaluation of cardiac biomarker elevation after percutaneous and surgical revascularization in the absence of manifest myocardial infarction. A comparative analysis of biomarkers and cardiac magnetic resonance. The MASS-V Trial. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 65	2.3	7
68	Strategies for multivessel revascularization in patients with diabetes. <i>New England Journal of Medicine</i> , 2012 , 367, 2375-84	59.2	1140
67	The Future REvascularization Evaluation in patients with Diabetes mellitus: optimal management of Multivessel disease (FREEDOM) trial: clinical and angiographic profile at study entry. <i>American Heart Journal</i> , 2012 , 164, 591-9	4.9	24
66	The effect of age on outcomes of coronary artery bypass surgery compared with balloon angioplasty or bare-metal stent implantation among patients with multivessel coronary disease. A collaborative analysis of individual patient data from 10 randomized trials. <i>Journal of the American College of Cardiology</i> , 2012 , 49, 2172-7	15.1	35
65	Cost-effectiveness analysis for surgical, angioplasty, or medical therapeutics for coronary artery disease: 5-year follow-up of medicine, angioplasty, or surgery study (MASS) II trial. <i>Circulation</i> , 2012 , 126, S145-50	16.7	28
64	Cancer-related deaths among different treatment options in chronic coronary artery disease: results of a 6-year follow-up of the MASS II study. <i>Coronary Artery Disease</i> , 2012 , 23, 79-84	1.4	19
63	Coronary to bronchial artery fistula: are we treating it right?. <i>Journal of Invasive Cardiology</i> , 2012 , 24, E303-4	0.7	3
62	New prognostic score for stable coronary disease evaluation. <i>Arquivos Brasileiros De Cardiologia</i> , 2011 , 96, 411-8	1.2	1
61	Genetic variants of diabetes risk and incident cardiovascular events in chronic coronary artery disease. <i>PLoS ONE</i> , 2011 , 6, e16341	3.7	6
60	The effect of internal thoracic artery grafts on long-term clinical outcomes after coronary bypass surgery. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2011 , 142, 829-35	1.5	13
59	Teratoma of the mediastinum: a case report. <i>Journal of Medical Case Reports</i> , 2011 , 5, 193	1.2	5
58	Effects of optimal medical treatment with or without coronary revascularization on angina and subsequent revascularizations in patients with type 2 diabetes mellitus and stable ischemic heart disease. <i>Circulation</i> , 2011 , 123, 1492-500	16.7	88
57	Preoperative B-type natriuretic peptide, and not the inflammation status, predicts an adverse outcome for patients undergoing heart surgery. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2011 , 12, 778-83	1.8	9
56	Efficacy of aneurysmectomy in patients with severe left ventricular dysfunction: favorable short-and long-term results in ischemic cardiomyopathy. <i>Clinics</i> , 2010 , 65, 947-52	2.3	3

55	Five-year follow-up of a randomized comparison between off-pump and on-pump stable multivessel coronary artery bypass grafting. The MASS III Trial. <i>Circulation</i> , 2010 , 122, S48-52	16.7	92
54	Five-year follow-up of angiographic disease progression after medicine, angioplasty, or surgery. <i>Journal of Cardiothoracic Surgery</i> , 2010 , 5, 91	1.6	24
53	Association between ADAMTS13 polymorphisms and risk of cardiovascular events in chronic coronary disease. <i>Thrombosis Research</i> , 2010 , 125, 61-6	8.2	27
52	Ten-year follow-up survival of the Medicine, Angioplasty, or Surgery Study (MASS II): a randomized controlled clinical trial of 3 therapeutic strategies for multivessel coronary artery disease. <i>Circulation</i> , 2010 , 122, 949-57	16.7	214
51	Reduced expression of systemic proinflammatory and myocardial biomarkers after off-pump versus on-pump coronary artery bypass surgery: a prospective randomized study. <i>Journal of Critical Care</i> , 2010 , 25, 305-12	4	41
50	Hypotheses, rationale, design, and methods for prognostic evaluation in type 2 diabetic patients with angiographically normal coronary arteries. The MASS IV-DM Trial. <i>BMC Cardiovascular Disorders</i> , 2010 , 10, 47	2.3	
49	TCF7L2 polymorphism rs7903146 is associated with coronary artery disease severity and mortality. <i>PLoS ONE</i> , 2009 , 4, e7697	3.7	50
48	Mild chronic kidney dysfunction and treatment strategies for stable coronary artery disease. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2009 , 137, 1443-9	1.5	14
47	In vitro simultaneous transfer of lipids to HDL in coronary artery disease and in statin treatment. <i>Lipids</i> , 2009 , 44, 917-24	1.6	47
46	Factors related to the selection of surgical versus percutaneous revascularization in diabetic patients with multivessel coronary artery disease in the BARI 2D (Bypass Angioplasty Revascularization Investigation in Type 2 Diabetes) trial. <i>JACC: Cardiovascular Interventions</i> , 2009 , 2, 384-92	5	26
45	Coronary artery bypass surgery compared with percutaneous coronary interventions for multivessel disease: a collaborative analysis of individual patient data from ten randomised trials. <i>Lancet, The</i> , 2009 , 373, 1190-7	40	531
44	Hemostatic changes and clinical sequelae after on-pump compared with off-pump coronary artery bypass surgery: a prospective randomized study. <i>Coronary Artery Disease</i> , 2009 , 20, 100-5	1.4	15
43	Large bilateral coronary artery fistula: the choice of clinical treatment. <i>Arquivos Brasileiros De Cardiologia</i> , 2009 , 93, e48-9	1.2	6
42	Long-term safety and efficacy of percutaneous coronary intervention with stenting and coronary artery bypass surgery for multivessel coronary artery disease: a meta-analysis with 5-year patient-level data from the ARTS, ERACI-II, MASS-II, and SoS trials. <i>Circulation</i> , 2008 , 118, 1146-54	16.7	222
41	Impact of number of vessels disease on outcome of patients with stable coronary artery disease: 5-year follow-up of the Medical, Angioplasty, and bypass Surgery study (MASS). <i>European Journal of Cardio-thoracic Surgery</i> , 2008 , 33, 349-54	3	15
40	Impact of metabolic syndrome on the outcome of patients with stable coronary artery disease: 2-year follow-up of the MASS II study. <i>Coronary Artery Disease</i> , 2008 , 19, 383-8	1.4	8
39	TCF7L2 variant genotypes and type 2 diabetes risk in Brazil: significant association, but not a significant tool for risk stratification in the general population. <i>BMC Medical Genetics</i> , 2008 , 9, 106	2.1	26
38	A randomized comparative study of patients undergoing myocardial revascularization with or without cardiopulmonary bypass surgery: The MASS III Trial. <i>Trials</i> , 2008 , 9, 52	2.8	8

37	Quality of life after on-pump and off-pump coronary artery bypass grafting surgery. <i>Arquivos Brasileiros De Cardiologia</i> , 2008 , 91, 217-22, 238-44	1.2	14
36	Comparative costs between myocardial revascularization with or without extracorporeal circulation. <i>Arquivos Brasileiros De Cardiologia</i> , 2008 , 91, 340-6	1.2	8
35	Effects of epinephrine in local dental anesthesia in patients with coronary artery disease. <i>Arquivos Brasileiros De Cardiologia</i> , 2007 , 88, 545-51	1.2	21
34	Quality of life after surgical myocardial revascularization, angioplasty or medical treatment. <i>Arquivos Brasileiros De Cardiologia</i> , 2007 , 88, 537-44	1.2	9
33	Five-year follow-up of the Medicine, Angioplasty, or Surgery Study (MASS II): a randomized controlled clinical trial of 3 therapeutic strategies for multivessel coronary artery disease. <i>Circulation</i> , 2007 , 115, 1082-9	16.7	262
32	Effect of a hypoglycemic agent on ischemic preconditioning in patients with type 2 diabetes and stable angina pectoris. <i>Coronary Artery Disease</i> , 2007 , 18, 55-9	1.4	20
31	Dynamic regulation of MTHFR mRNA expression and C677T genotype modulate mortality in coronary artery disease patients after revascularization. <i>Thrombosis Research</i> , 2007 , 121, 25-32	8.2	17
30	Impact of diabetes on five-year outcomes of patients with multivessel coronary artery disease. <i>Annals of Thoracic Surgery</i> , 2007 , 83, 93-9	2.7	33
29	Quality of life in patients with symptomatic multivessel coronary artery disease: a comparative post hoc analyses of medical, angioplasty or surgical strategies-MASS II trial. <i>International Journal of Cardiology</i> , 2007 , 116, 364-70	3.2	57
28	Coronary revascularization (surgical or percutaneous) decreases mortality after the first year in diabetic subjects but not in nondiabetic subjects with multivessel disease: an analysis from the Medicine, Angioplasty, or Surgery Study (MASS II). <i>Circulation</i> , 2006 , 114, 1420-4	16.7	36
27	Clinical judgment and treatment options in stable multivessel coronary artery disease: results from the one-year follow-up of the MASS II (Medicine, Angioplasty, or Surgery Study II). <i>Journal of the American College of Cardiology</i> , 2006 , 48, 948-53	15.1	24
26	Association between platelet P2Y12 haplotype and risk of cardiovascular events in chronic coronary disease. <i>Thrombosis Research</i> , 2006 , 118, 679-83	8.2	20
25	Complete, incomplete or no myocardial revascularization. <i>Arquivos Brasileiros De Cardiologia</i> , 2006 , 87, e144-6	1.2	1
24	Exercise stress testing before and after successful multivessel percutaneous transluminal coronary angioplasty. <i>Brazilian Journal of Medical and Biological Research</i> , 2006 , 39, 475-82	2.8	2
23	Culprit lesion revascularisation initially costs less and is more successful than complete revascularisation in multivessel disease, but results equalise over time. Commentary. <i>Evidence-based Cardiovascular Medicine</i> , 2005 , 9, 52-4		
22	The effects of glibenclamide, a K(ATP) channel blocker, on the warm-up phenomenon. <i>Annals of Noninvasive Electrocardiology</i> , 2005 , 10, 356-62	1.5	16
21	One-year outcomes of coronary artery bypass graft surgery versus percutaneous coronary intervention with multiple stenting for multisystem disease: a meta-analysis of individual patient data from randomized clinical trials. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2005 , 130, 512-9	1.5	123
20	Effect of glycoprotein IIIa PLA2 polymorphism on outcome of patients with stable coronary artery disease and effect of smoking. <i>American Journal of Cardiology</i> , 2004 , 93, 1469-72	3	16

19	The medicine, angioplasty, or surgery study (MASS-II): a randomized, controlled clinical trial of three therapeutic strategies for multivessel coronary artery disease: one-year results. <i>Journal of the American College of Cardiology</i> , 2004 , 43, 1743-51	15.1	286
18	Impaired intravascular triglyceride lipolysis constitutes a marker of clinical outcome in patients with stable angina undergoing secondary prevention treatment: a long-term follow-up study. <i>Journal of the American College of Cardiology</i> , 2004 , 43, 2225-32	15.1	26
17	Plasma kinetics of a cholesterol-rich emulsion in subjects with or without coronary artery disease. <i>Journal of Lipid Research</i> , 2003 , 44, 464-9	6.3	25
16	Relative cost comparison of treatments for coronary artery disease: the First Year Follow-Up of MASS II Study. <i>Circulation</i> , 2003 , 108 Suppl 1, I121-3	16.7	10
15	Coronary heart disease clinical manifestation and risk factors in Japanese immigrants and their descendents in the city of S Paulo. <i>Arquivos Brasileiros De Cardiologia</i> , 2003 , 81, 229-8	1.2	2
14	Detection of retained surgical sponge by transthoracic and transesophageal echocardiography. <i>Journal of the American Society of Echocardiography</i> , 2003 , 16, 1191-3	5.8	10
13	Acute pleuropulmonary complications detected by computed tomography following myocardial revascularization. <i>Revista Do Hospital Das Clinicas</i> , 2002 , 57, 135-42		18
12	LDL concentration is correlated with the removal from the plasma of a chylomicron-like emulsion in subjects with coronary artery disease. <i>Atherosclerosis</i> , 2002 , 161, 447-53	3.1	12
11	Five-year follow-up of the medicine, angioplasty, or surgery study (MASS): A prospective, randomized trial of medical therapy, balloon angioplasty, or bypass surgery for single proximal left anterior descending coronary artery stenosis. <i>Circulation</i> , 1999 , 100, I1107-13	16.7	47
10	Pulmonary function after coronary artery bypass surgery. <i>Respiratory Medicine</i> , 1997 , 91, 629-33	4.6	53
9	VATS for complete dissection of LIMA in minimally invasive coronary artery bypass grafting. <i>Annals of Thoracic Surgery</i> , 1997 , 63, S110-3	2.7	14
8	The Medicine, Angioplasty or Surgery Study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenoses. <i>Journal of the American College of Cardiology</i> , 1995 , 26, 1600-5	15.1	246
7	Meta-analysis of randomised trials comparing coronary angioplasty with bypass surgery. <i>Lancet, The</i> , 1995 , 346, 1184-9	40	392
6	Relationship between pleural effusion and pericardial involvement after myocardial revascularization. <i>Chest</i> , 1994 , 105, 1748-52	5.3	42
5	Influence of atelectasis on pulmonary function after coronary artery bypass grafting. <i>Chest</i> , 1993 , 104, 434-7	5.3	28
4	Ventricular arrhythmias induced by programmed ventricular stimulation after uncomplicated myocardial infarction. <i>Angiology</i> , 1992 , 43, 578-84	2.1	
3	Relationship between pleural changes after myocardial revascularization and pulmonary mechanics. <i>Chest</i> , 1992 , 102, 1333-6	5.3	39
2	Two- to eight-year survival rates in patients who refused coronary artery bypass grafting. <i>American Journal of Cardiology</i> , 1989 , 63, 155-9	3	10

- 1 Coronary bypass surgery for patients with renal transplantation. *Cardiology*, **1986**, 73, 151-5 1.6 3