

Micha? Ambroziak

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3914546/publications.pdf>

Version: 2024-02-01

10
papers

168
citations

1163117

8
h-index

1281871

11
g-index

14
all docs

14
docs citations

14
times ranked

242
citing authors

#	ARTICLE	IF	CITATIONS
1	Younger age of patients with myocardial infarction is associated with a higher number of relatives with a history of premature atherosclerosis. BMC Cardiovascular Disorders, 2020, 20, 410.	1.7	8
2	Increased coagulation factor XIII activity but not genetic variants of coagulation factors is associated with myocardial infarction in young patients. Journal of Thrombosis and Thrombolysis, 2019, 48, 519-527.	2.1	10
3	Adiponectin gene variants and decreased adiponectin plasma levels are associated with the risk of myocardial infarction in young age. Gene, 2018, 642, 498-504.	2.2	14
4	ESR2 gene G1730A variant is associated with triglycerides level and myocardial infarction in young men but not in women. Gene, 2018, 677, 83-88.	2.2	2
5	The $\epsilon^{351A/G}$ polymorphism of ESR1 is associated with risk of myocardial infarction but not with extreme longevity. Clinica Chimica Acta, 2010, 411, 1883-1887.	1.1	11
6	Human Breast Cancer Tissue Expresses High Level of Type 1 ϵ^2 -Deiodinase. Thyroid, 2007, 17, 3-10.	4.5	26
7	Disturbed Expression of Type 1 and Type 2 Iodothyronine Deiodinase As Well As Titf1/Nkx2-1 and Pax-8 Transcription Factor Genes in Papillary Thyroid Cancer. Thyroid, 2005, 15, 1137-1146.	4.5	32
8	Thyroid Sialyltransferase mRNA Level and Activity Are Increased in Graves' Disease. Thyroid, 2005, 15, 645-652.	4.5	15
9	Pax-8 Expression Correlates with Type II ϵ^2 Deiodinase Expression in Thyroids from Patients with Graves' Disease. Thyroid, 2003, 13, 141-148.	4.5	7
10	Type I ϵ^2 -iodothyronine deiodinase activity and mRNA are remarkably reduced in renal clear cell carcinoma. Journal of Endocrinological Investigation, 2001, 24, 253-261.	3.3	42