

Jarmo Hietala

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3918169/publications.pdf>

Version: 2024-02-01

137
papers

7,323
citations

57758

44
h-index

62596

80
g-index

148
all docs

148
docs citations

148
times ranked

7883
citing authors

#	ARTICLE	IF	CITATIONS
1	Age-related dopamine D2/D3 receptor loss in extrastriatal regions of the human brain. <i>Neurobiology of Aging</i> , 2000, 21, 683-688.	3.1	313
2	Lack of Efficacy of the Substance P (Neurokinin1 Receptor) Antagonist Aprepitant in the Treatment of Major Depressive Disorder. <i>Biological Psychiatry</i> , 2006, 59, 216-223.	1.3	287
3	Prediction Models of Functional Outcomes for Individuals in the Clinical High-Risk State for Psychosis or With Recent-Onset Depression. <i>JAMA Psychiatry</i> , 2018, 75, 1156.	11.0	251
4	Depressive symptoms and presynaptic dopamine function in neuroleptic-naïve schizophrenia. <i>Schizophrenia Research</i> , 1999, 35, 41-50.	2.0	239
5	Striatal D2 dopamine receptor binding characteristics in vivo in patients with alcohol dependence. <i>Psychopharmacology</i> , 1994, 116, 285-290.	3.1	235
6	Common and distinct neural correlates of emotional processing in Bipolar Disorder and Major Depressive Disorder: A voxel-based meta-analysis of functional magnetic resonance imaging studies. <i>European Neuropsychopharmacology</i> , 2012, 22, 100-113.	0.7	206
7	Striatal dopamine D2 receptors in modulation of pain in humans: a review. <i>European Journal of Pharmacology</i> , 2004, 500, 187-192.	3.5	199
8	Human positron emission tomography studies of brain neurokinin 1 receptor occupancy by aprepitant. <i>Biological Psychiatry</i> , 2004, 55, 1007-1012.	1.3	187
9	Decrease in Human Striatal Dopamine D ₂ Receptor Density with Age: A PET Study with [¹¹ C]Raclopride. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1993, 13, 310-314.	4.3	186
10	Sex differences in striatal presynaptic dopamine synthesis capacity in healthy subjects. <i>Biological Psychiatry</i> , 2002, 52, 759-763.	1.3	181
11	Mesolimbic dopamine release is linked to symptom severity in pathological gambling. <i>NeuroImage</i> , 2012, 60, 1992-1999.	4.2	181
12	Striatal D2 Dopamine Receptor Characteristics in Neuroleptic-Naïve Schizophrenic Patients Studied With Positron Emission Tomography. <i>Archives of General Psychiatry</i> , 1994, 51, 116.	12.3	176
13	A PET-study of [¹¹ C]FLB 457 binding to extrastriatal D ₂ -dopamine receptors in healthy subjects and antipsychotic drug-treated patients. <i>Psychopharmacology</i> , 1997, 133, 396-404.	3.1	174
14	C957T polymorphism of dopamine D2 receptor gene affects striatal DRD2 in vivo availability by changing the receptor affinity. <i>Synapse</i> , 2009, 63, 907-912.	1.2	156
15	Decreased brain serotonin 5-HT _{1A} receptor availability in medication-naïve patients with major depressive disorder: an in-vivo imaging study using PET and [carbonyl- ¹¹ C]WAY-100635. <i>International Journal of Neuropsychopharmacology</i> , 2008, 11, 465-76.	2.1	150
16	Comparison of the Transient Equilibrium and Continuous Infusion Method for Quantitative PET Analysis of [¹¹ C]Raclopride Binding. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1998, 18, 941-950.	4.3	144
17	Increased Caudate Dopamine D2 Receptor Availability as a Genetic Marker for Schizophrenia. <i>Archives of General Psychiatry</i> , 2005, 62, 371.	12.3	142
18	Sex Differences in Extrastriatal Dopamine D2-Like Receptors in the Human Brain. <i>American Journal of Psychiatry</i> , 2001, 158, 308-311.	7.2	138

#	ARTICLE	IF	CITATIONS
19	Dopamine D2 receptor binding in the human brain is associated with the response to painful stimulation and pain modulatory capacity. <i>Pain</i> , 2002, 99, 273-279.	4.2	129
20	Multimodal Machine Learning Workflows for Prediction of Psychosis in Patients With Clinical High-Risk Syndromes and Recent-Onset Depression. <i>JAMA Psychiatry</i> , 2021, 78, 195.	11.0	125
21	Striatal Dopamine Synthesis in First-degree Relatives of Patients with Schizophrenia. <i>Biological Psychiatry</i> , 2008, 63, 114-117.	1.3	110
22	C957T polymorphism of the human dopamine D2 receptor gene predicts extrastriatal dopamine receptor availability in vivo. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 630-636.	4.8	106
23	Personality Traits and Striatal Dopamine Synthesis Capacity in Healthy Subjects. <i>American Journal of Psychiatry</i> , 2003, 160, 904-910.	7.2	104
24	Measurement of Striatal D2 Dopamine Receptor Density and Affinity with [¹¹ C]-Raclopride in Vivo: A Test-Retest Analysis. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1999, 19, 210-217.	4.3	80
25	Prolonged Central μ -Opioid Receptor Occupancy after Single and Repeated Nalmefene Dosing. <i>Neuropsychopharmacology</i> , 2005, 30, 2245-2253.	5.4	80
26	The amygdala and schizophrenia: a volumetric magnetic resonance imaging study in first-episode, neuroleptic-naïve patients. <i>Biological Psychiatry</i> , 2003, 54, 1302-1304.	1.3	76
27	Short-Term Psychodynamic Psychotherapy and Fluoxetine in Major Depressive Disorder: A Randomized Comparative Study. <i>Psychotherapy and Psychosomatics</i> , 2008, 77, 351-357.	8.8	75
28	Striatal dopamine D2/D3 receptor availability correlates with individual response characteristics to pain. <i>European Journal of Neuroscience</i> , 2004, 20, 1587-1592.	2.6	74
29	Right secondary somatosensory cortex—a promising novel target for the treatment of drug-resistant neuropathic orofacial pain with repetitive transcranial magnetic stimulation. <i>Pain</i> , 2015, 156, 1276-1283.	4.2	73
30	Variation in the dopamine D2 receptor gene plays a key role in human pain and its modulation by transcranial magnetic stimulation. <i>Pain</i> , 2014, 155, 2180-2187.	4.2	70
31	Brain Dopamine D ₁ Receptors in Twins Discordant for Schizophrenia. <i>American Journal of Psychiatry</i> , 2006, 163, 1747-1753.	7.2	68
32	Measurement of Serotonin 5-HT _{1A} Receptor Binding Using Positron Emission Tomography and [carbonyl- ¹¹ C]WAY-100635—Considerations on the Validity of Cerebellum as a Reference Region. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2007, 27, 185-195.	4.3	66
33	Sex difference in brain CB1 receptor availability in man. <i>NeuroImage</i> , 2019, 184, 834-842.	4.2	65
34	Measurement of Striatal and Extrastriatal Dopamine Transporter Binding with High-Resolution PET and [¹¹ C]PE2I: Quantitative Modeling and Test-Retest Reproducibility. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2008, 28, 1059-1069.	4.3	63
35	Striatal dopamine D2 receptors in medication-naïve patients with major depressive disorder as assessed with [¹¹ C]raclopride PET. <i>Psychopharmacology</i> , 2008, 197, 581-590.	3.1	61
36	Aberrant mesolimbic dopamine–opiate interaction in obesity. <i>NeuroImage</i> , 2015, 122, 80-86.	4.2	61

#	ARTICLE	IF	CITATIONS
37	Lack of association between the functional variant of the catechol-o-methyltransferase (COMT) gene and early-onset alcoholism associated with severe antisocial behavior. American Journal of Medical Genetics Part A, 2000, 96, 348-352.	2.4	60
38	Interindividual variability and lateralization of μ -opioid receptors in the human brain. NeuroImage, 2020, 217, 116922.	4.2	60
39	[18F]CFT ([18F]WIN 35,428), a radioligand to study the dopamine transporter with PET: Characterization in human subjects. , 1998, 28, 244-250.		58
40	Association of striatal dopamine D2/D3 receptor binding potential with pain but not tactile sensitivity or placebo analgesia. Neuroscience Letters, 2005, 376, 149-153.	2.1	57
41	μ -Receptor agonism with alfentanil increases striatal dopamine D2 receptor binding in man. Synapse, 2002, 45, 25-30.	1.2	56
42	Trends in the long-term use of benzodiazepine anxiolytics and hypnotics: A national register study for 2006 to 2014. Pharmacoepidemiology and Drug Safety, 2018, 27, 674-682.	1.9	53
43	Childhood physical abuse and emotional neglect are specifically associated with adult mental disorders. Journal of Mental Health, 2020, 29, 376-384.	1.9	52
44	Dopamine in Schizophrenia. Annals of Medicine, 1996, 28, 557-561.	3.8	51
45	Visualization and Quantification of Neurokinin-1 (NK1) Receptors in the Human Brain. Molecular Imaging and Biology, 2005, 7, 262-272.	2.6	51
46	Measurement of extrastriatal D2-like receptor binding with [11C]FLB 457 â€” a test-retest analysis. European Journal of Nuclear Medicine and Molecular Imaging, 2000, 27, 1666-1673.	6.4	50
47	In Vivo Availability of Cannabinoid 1 Receptor Levels in Patients With First-Episode Psychosis. JAMA Psychiatry, 2019, 76, 1074.	11.0	50
48	Dopamine D2/D3 receptor binding in the anterior cingulate cortex and executive functioning. Psychiatry Research - Neuroimaging, 2007, 156, 69-74.	1.8	48
49	[18F]CFT ([18F]WIN 35,428), a radioligand to study the dopamine transporter with PET: Biodistribution in rats. , 1996, 23, 321-327.		45
50	Mapping neurotransmitter networks with PET: An example on serotonin and opioid systems. Human Brain Mapping, 2014, 35, 1875-1884.	3.6	45
51	Synthesis and characterization of a potent, selective, radiolabeled substance-P antagonist for NK receptor quantitation: ([F]SPA-RQ). Molecular Imaging and Biology, 2004, 6, 373-384.	2.6	44
52	Dopaminergic and serotonergic mechanisms in the modulation of pain: In vivo studies in human brain. European Journal of Pharmacology, 2018, 834, 337-345.	3.5	44
53	Lowered endogenous mu-opioid receptor availability in subclinical depression and anxiety. Neuropsychopharmacology, 2020, 45, 1953-1959.	5.4	44
54	Gender differences in brain serotonin transporter availability in panic disorder. Journal of Psychopharmacology, 2011, 25, 952-959.	4.0	41

#	ARTICLE	IF	CITATIONS
55	Magia: Robust Automated Image Processing and Kinetic Modeling Toolbox for PET Neuroinformatics. <i>Frontiers in Neuroinformatics</i> , 2020, 14, 3.	2.5	41
56	Lack of efficacy of L-759274, a novel neurokinin 1 (substance P) receptor antagonist, for the treatment of generalized anxiety disorder. <i>International Journal of Neuropsychopharmacology</i> , 2013, 16, 1-11.	2.1	40
57	Shape variability of the human striatum—Effects of age and gender. <i>NeuroImage</i> , 2007, 34, 85-93.	4.2	38
58	Correlation of human cold pressor pain responses with 5-HT _{1A} receptor binding in the brain. <i>Brain Research</i> , 2007, 1172, 21-31.	2.2	37
59	Dopamine 2 receptor C957T and catechol-o-methyltransferase Val158Met polymorphisms are associated with treatment response in electroconvulsive therapy. <i>Neuroscience Letters</i> , 2008, 448, 79-83.	2.1	37
60	Temperament trait Harm Avoidance associates with μ -opioid receptor availability in frontal cortex: A PET study using [¹¹ C]carfentanil. <i>NeuroImage</i> , 2012, 61, 670-676.	4.2	37
61	Meta-analysis of the Glial Marker TSPO in Psychosis Revisited: Reconciling Inconclusive Findings of Patient—Control Differences. <i>Biological Psychiatry</i> , 2021, 89, e5-e8.	1.3	36
62	Receptor binding profile and anxiolytic-type activity of deramciclane (EGIS-3886) in animal models. <i>Drug Development Research</i> , 1997, 40, 333-348.	2.9	34
63	Effects of childhood and adolescence physical activity patterns on psychosis risk—a general population cohort study. <i>NPJ Schizophrenia</i> , 2017, 3, 5.	3.6	34
64	The effects of d-amphetamine on extrastriatal dopamine D ₂ /D ₃ receptors: a randomized, double-blind, placebo-controlled PET study with [¹¹ C]FLB 457 in healthy subjects. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2009, 36, 475-483.	6.4	33
65	Effects of antidepressant drug treatment and psychotherapy on striatal and thalamic dopamine D _{2/3} receptors in major depressive disorder studied with [¹¹ C]raclopride PET. <i>Journal of Psychopharmacology</i> , 2011, 25, 1329-1336.	4.0	33
66	Gender and age affect NK ₁ receptors in the human brain—a positron emission tomography study with [¹⁸ F]SPA-RQ. <i>International Journal of Neuropsychopharmacology</i> , 2007, 10, 219.	2.1	32
67	Toward Generalizable and Transdiagnostic Tools for Psychosis Prediction: An Independent Validation and Improvement of the NAPLS-2 Risk Calculator in the Multisite PRONIA Cohort. <i>Biological Psychiatry</i> , 2021, 90, 632-642.	1.3	32
68	Sertindole is a serotonin 5-HT _{2c} inverse agonist and decreases agonist but not antagonist binding to 5-HT _{2c} receptors after chronic treatment. <i>Psychopharmacology</i> , 2001, 157, 180-187.	3.1	30
69	Striatal dopamine D ₁ and D ₂ receptor balance in twins at increased genetic risk for schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , 2006, 146, 13-20.	1.8	30
70	COMT Val158Met Genotype Does Not Alter Cortical or Striatal Dopamine D ₂ Receptor Availability In Vivo. <i>Molecular Imaging and Biology</i> , 2010, 12, 192-197.	2.6	30
71	Effects of fluoxetine on dopamine D ₂ receptors in the human brain: a positron emission tomography study with [¹¹ C]raclopride. <i>International Journal of Neuropsychopharmacology</i> , 2004, 7, 431-439.	2.1	29
72	Striatal μ -opioid receptor availability predicts cold pressor pain threshold in healthy human subjects. <i>Neuroscience Letters</i> , 2012, 521, 11-14.	2.1	29

#	ARTICLE	IF	CITATIONS
73	Measurement of Cortical Dopamine D1 Receptor Binding with [11C]SCH 23390: A Testâ€“Retest Analysis. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2001, 21, 1146-1150.	4.3	28
74	Dysfunctional Brain Networks and Genetic Risk for Schizophrenia: Specific Neurotransmitter Systems. <i>CNS Neuroscience and Therapeutics</i> , 2011, 17, 89-96.	3.9	28
75	Automatic cerebral and cerebellar hemisphere segmentation in 3D MRI: Adaptive disconnection algorithm. <i>Medical Image Analysis</i> , 2010, 14, 360-372.	11.6	26
76	Neuroticism and serotonin 5-HT1A receptors in healthy subjects. <i>Psychiatry Research - Neuroimaging</i> , 2015, 234, 1-6.	1.8	26
77	The Cannabinoid Receptor-1 Is an Imaging Biomarker of Brown Adipose Tissue. <i>Journal of Nuclear Medicine</i> , 2015, 56, 1937-1941.	5.0	24
78	A study on the association of psychiatric diagnoses and childhood adversities with suicide risk. <i>Nordic Journal of Psychiatry</i> , 2019, 73, 125-131.	1.3	24
79	Incidence of and Characteristics Associated With Long-term Benzodiazepine Use in Finland. <i>JAMA Network Open</i> , 2020, 3, e2019029.	5.9	24
80	Links between central CB1-receptor availability and peripheral endocannabinoids in patients with first episode psychosis. <i>NPJ Schizophrenia</i> , 2020, 6, 21.	3.6	23
81	Increased Risk of Parkinson's Disease in Patients With Schizophrenia Spectrum Disorders. <i>Movement Disorders</i> , 2021, 36, 1353-1361.	3.9	23
82	Amygdala subnucleus volumes in psychosis high-risk state and first-episode psychosis. <i>Schizophrenia Research</i> , 2020, 215, 284-292.	2.0	22
83	A PET Study on the Acute Effect of Ethanol on Striatal D2 Dopamine Receptors with [11C]Raclopride in Healthy Males. <i>Human Psychopharmacology</i> , 1997, 12, 145-152.	1.5	20
84	Automatic statistical shape analysis of cerebral asymmetry in 3D T1-weighted magnetic resonance images at vertex-level: Application to neuroleptic-naïve schizophrenia. <i>Magnetic Resonance Imaging</i> , 2013, 31, 676-687.	1.8	20
85	Depression predicts persistence of paranoia in clinical high-risk patients to psychosis: results of the EPOS project. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 247-257.	3.1	20
86	Quantification of [Carbonyl-11C]WAY-100635 binding: considerations on the cerebellum. <i>Nuclear Medicine and Biology</i> , 2000, 27, 483-486.	0.6	19
87	Neuroticism Associates with Cerebral in Vivo Serotonin Transporter Binding Differently in Males and Females. <i>International Journal of Neuropsychopharmacology</i> , 2017, 20, 963-970.	2.1	19
88	Differential tolerance to cataleptic effects of SCH 23390 and haloperidol after repeated administration. <i>Psychopharmacology</i> , 1989, 98, 472-475.	3.1	18
89	Ligandâ€“receptor interactions as studied by PET: implications for drug development. <i>Annals of Medicine</i> , 1999, 31, 438-443.	3.8	17
90	Adult Attachment System Links With Brain Mu Opioid Receptor Availability Inâ€“Vivo. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2021, 6, 360-369.	1.5	17

#	ARTICLE	IF	CITATIONS
91	Increased Serotonin Receptor 1A Binding in Major Depressive Disorder after Psychotherapy, but Not after SSRI Pharmacotherapy, Is Related to Improved Social Functioning Capacity. <i>Psychotherapy and Psychosomatics</i> , 2013, 82, 260-261.	8.8	16
92	Elevated serum chemokine CCL22 levels in first-episode psychosis: associations with symptoms, peripheral immune state and in vivo brain glial cell function. <i>Translational Psychiatry</i> , 2020, 10, 94.	4.8	16
93	Serotonin transporter in attention-deficit hyperactivity disorder – preliminary results from a positron emission tomography study. <i>Psychiatry Research - Neuroimaging</i> , 2013, 212, 164-165.	1.8	15
94	Platform for systems medicine research and diagnostic applications in psychotic disorders – The METSY project. <i>European Psychiatry</i> , 2018, 50, 40-46.	0.2	14
95	Childhood adversity predicts persistence of suicidal thoughts differently in females and males at clinical high-risk patients of psychosis. Results of the EPOS project. <i>Microbial Biotechnology</i> , 2019, 13, 935-942.	1.7	14
96	Seasonal Variation in the Brain μ -Opioid Receptor Availability. <i>Journal of Neuroscience</i> , 2021, 41, 1265-1273.	3.6	14
97	Antagonist Binding Characteristics of the Ser311 \rightarrow Cys Variant of Human Dopamine D2 Receptor in Vivo and in Vitro. <i>Biochemical and Biophysical Research Communications</i> , 1997, 232, 143-146.	2.1	13
98	Verbal memory and 5-HT _{1A} receptors in healthy volunteers – A PET study with [carbonyl- ¹¹ C]WAY-100635. <i>European Neuropsychopharmacology</i> , 2016, 26, 570-577.	0.7	13
99	Brain 5-HT _{2A} receptor occupancy of deramciclane in humans after a single oral administration - a positron emission tomography study. <i>Psychopharmacology</i> , 1999, 145, 76-81.	3.1	10
100	DRD2-Related TaqIA Genotype Is Associated With Dopamine Release During a Gambling Task. <i>Journal of Addiction Medicine</i> , 2014, 8, 294-295.	2.6	10
101	Effect of childhood adversities on alcohol problems is mainly mediated by depression. <i>American Journal on Addictions</i> , 2018, 27, 391-399.	1.4	10
102	The neural and molecular basis of working memory function in psychosis: a multimodal PET-fMRI study. <i>Molecular Psychiatry</i> , 2021, 26, 4464-4474.	7.9	10
103	Personality traits and recovery from major depressive disorder. <i>Nordic Journal of Psychiatry</i> , 2011, 65, 52-57.	1.3	9
104	Association Between Circulating Lipids and Future Weight Gain in Individuals With an At-Risk Mental State and in First-Episode Psychosis. <i>Schizophrenia Bulletin</i> , 2021, 47, 160-169.	4.3	9
105	Cerebral grey matter density is associated with neuroreceptor and neurotransmitter availability: A combined PET and MRI study. <i>NeuroImage</i> , 2021, 235, 117968.	4.2	9
106	Evaluation of the automatic three-dimensional delineation of caudate and putamen for PET receptor occupancy studies. <i>Nuclear Medicine Communications</i> , 2008, 29, 53-65.	1.1	8
107	Adverse childhood experiences leads to perceived negative attitude of others and the effect of adverse childhood experiences on depression in adulthood is mediated via negative attitude of others. <i>European Psychiatry</i> , 2018, 54, 27-34.	0.2	8
108	Longitudinal brain morphology in anti-NMDA receptor encephalitis: a case report with controls. <i>BMC Psychiatry</i> , 2019, 19, 145.	2.6	8

#	ARTICLE	IF	CITATIONS
109	Body mass index trajectories in childhood and adolescence - Risk for non-affective psychosis. Schizophrenia Research, 2019, 206, 313-317.	2.0	8
110	Atlas of type 2 dopamine receptors in the human brain: Age and sex dependent variability in a large PET cohort. NeuroImage, 2022, 255, 119149.	4.2	8
111	Cortical dopamine D2/D3 receptors and verbal memory in man. NeuroImage, 2010, 51, 918-922.	4.2	7
112	Multimodal prognosis of negative symptom severity in individuals at increased risk of developing psychosis. Translational Psychiatry, 2021, 11, 312.	4.8	7
113	Brain neurokinin-1 receptor availability in never-medicated patients with major depression – A pilot study. Journal of Affective Disorders, 2019, 242, 188-194.	4.1	6
114	Implementation of CYP2D6 copy-number imputation panel and frequency of key pharmacogenetic variants in Finnish individuals with a psychotic disorder. Pharmacogenomics Journal, 2022, 22, 166-172.	2.0	6
115	Magical thinking in individuals with high polygenic risk for schizophrenia but no non-affective psychoses – a general population study. Molecular Psychiatry, 2022, 27, 3286-3293.	7.9	6
116	Clinical, Brain, and Multilevel Clustering in Early Psychosis and Affective Stages. JAMA Psychiatry, 2022, 79, 677.	11.0	6
117	Voxel-based NK1 Receptor Occupancy Measurements with [18F]SPA-RQ and Positron Emission Tomography: A Procedure for Assessing Errors from Image Reconstruction and Physiological Modeling. Molecular Imaging and Biology, 2007, 9, 284-294.	2.6	5
118	Basic Symptoms Are Associated With Age in Patients With a Clinical High-Risk State for Psychosis: Results From the PRONIA Study. Frontiers in Psychiatry, 2020, 11, 552175.	2.6	5
119	Segmentation of Striatal Brain Structures from High Resolution PET Images. International Journal of Biomedical Imaging, 2009, 2009, 1-12.	3.9	4
120	Short-term functional outcome in psychotic patients: results of the Turku early psychosis study (TEPS). BMC Psychiatry, 2021, 21, 602.	2.6	4
121	42.4 THE ENDOCANNABINOID SYSTEM IN FIRST-EPISODE PSYCHOSIS. Schizophrenia Bulletin, 2018, 44, S69-S69.	4.3	3
122	Reaction Time and Visual Memory in Connection with Alcohol Use in Schizophrenia and Schizoaffective Disorder. Brain Sciences, 2021, 11, 688.	2.3	3
123	Chronic D1-receptor blockade: effects on D2-receptor agonist-induced yawning in rats. Journal of Pharmacy and Pharmacology, 2011, 43, 278-279.	2.4	2
124	Dopamine Receptor Imaging in Schizophrenia. , 2014, , 341-360.		2
125	F135. BODY MASS INDEX TRAJECTORIES IN CHILDHOOD AND RISK FOR NON-AFFECTIVE PSYCHOSIS – A GENERAL POPULATION COHORT STUDY. Schizophrenia Bulletin, 2018, 44, S272-S272.	4.3	2
126	Patient-Made Videos as a Tool of Self-Observation Enhancing Self-Reflection in Psychotherapy. Journal of Contemporary Psychotherapy, 2019, 49, 187-195.	1.2	2

#	ARTICLE	IF	CITATIONS
127	A method for automatic extraction of striatal structures for dose-finding studies in PET. , 2006, , .		1
128	S170. AMYGDALA SUBNUCLEI VOLUMES IN FIRST-EPISODE PSYCHOSIS: ASSOCIATION WITH CHILDHOOD ADVERSITY. Schizophrenia Bulletin, 2018, 44, S391-S391.	4.3	1
129	[18F]SPA-RQ/PET Study of NK1 receptors in the Whole Body of Guinea Pig and Rat. Scientific Reports, 2019, 9, 20412.	3.3	1
130	Reaction Time and Visual Memory in Connection to Alcohol Use in Persons with Bipolar Disorder. Brain Sciences, 2021, 11, 1154.	2.3	1
131	Selection bias in clinical studies of first-episode psychosis: A follow-up study. Schizophrenia Research, 2022, 246, 235-240.	2.0	1
132	Author reply to: "Depressive symptomatology, serotonergic activity, and neuroticism: A methodological recommendation". Psychiatry Research - Neuroimaging, 2015, 234, 391.	1.8	0
133	SA88. Hippocampal Subfield Volumes in First-Episode Psychosis: Association With Glucose Metabolism. Schizophrenia Bulletin, 2017, 43, S144-S145.	4.3	0
134	O5.5. SLEEP IN MAJOR PSYCHIATRIC DISORDERS: RESULTS FROM NATIONWIDE SUPER FINLAND STUDY. Schizophrenia Bulletin, 2018, 44, S88-S88.	4.3	0
135	T7. UPDATED INDIVIDUAL PARTICIPANT DATA META-ANALYSIS CONFIRMS LOWER LEVELS OF THE GLIAL MARKER TSPO IN PSYCHOSIS PATIENTS. Schizophrenia Bulletin, 2020, 46, S233-S233.	4.3	0
136	Reaction Time and Visual Memory in Connection to Hazardous Drinking Polygenic Scores in Schizophrenia, Schizoaffective Disorder and Bipolar Disorder. Brain Sciences, 2021, 11, 1422.	2.3	0
137	Evidence of discontinuity between psychosis-risk and non-clinical samples in the neuroanatomical correlates of social function. Schizophrenia Research: Cognition, 2022, 29, 100252.	1.3	0