
Hongtao Liu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3923427/publications.pdf Version: 2024-02-01



Номстло Ци

#	Article	IF	CITATIONS
1	Geriatric assessment to predict survival in older allogeneic hematopoietic cell transplantation recipients. Haematologica, 2014, 99, 1373-1379.	3.5	213
2	Reduced-intensity conditioning with combined haploidentical and cord blood transplantation results in rapid engraftment, low GVHD, and durable remissions. Blood, 2011, 118, 6438-6445.	1.4	158
3	Next-generation immuno-oncology agents: current momentum shifts in cancer immunotherapy. Journal of Hematology and Oncology, 2020, 13, 29.	17.0	146
4	Emerging agents and regimens for AML. Journal of Hematology and Oncology, 2021, 14, 49.	17.0	104
5	Posttransplant cyclophosphamide is associated with increased cytomegalovirus infection: a CIBMTR analysis. Blood, 2021, 137, 3291-3305.	1.4	85
6	The past, present, and future of CRM1/XPO1 inhibitors. Stem Cell Investigation, 2019, 6, 6-6.	3.0	77
7	Results from a multidisciplinary clinic guided by geriatric assessment before stem cell transplantation in older adults. Blood Advances, 2019, 3, 3488-3498.	5.2	62
8	Gal9/Tim-3 expression level is higher in AML patients who fail chemotherapy. , 2019, 7, 175.		59
9	A phase I study of selinexor in combination with high-dose cytarabine and mitoxantrone for remission induction in patients with acute myeloid leukemia. Journal of Hematology and Oncology, 2018, 11, 4.	17.0	52
10	Umbilical Cord Blood Transplantation Supported by Third-Party Donor Cells: Rationale, Results, andÂApplications. Biology of Blood and Marrow Transplantation, 2013, 19, 682-691.	2.0	35
11	Haploidentical vs sibling, unrelated, or cord blood hematopoietic cell transplantation for acute lymphoblastic leukemia. Blood Advances, 2022, 6, 339-357.	5.2	35
12	Haploidentical vs haplo-cord transplant in adults under 60 years receiving fludarabine and melphalan conditioning. Blood Advances, 2019, 3, 1858-1867.	5.2	25
13	WT1 peptide vaccine in Montanide in contrast to poly ICLC, is able to induce WT1-specific immune response with TCR clonal enrichment in myeloid leukemia. Experimental Hematology and Oncology, 2018, 7, 1.	5.0	24
14	Immune reconstitution after combined haploidentical and umbilical cord blood transplant. Leukemia and Lymphoma, 2013, 54, 1242-1249.	1.3	23
15	Outcomes following second allogeneic stem cell transplant for disease relapse after T cell depleted transplant correlate with remission status and remission duration after the first transplant. Experimental Hematology and Oncology, 2019, 8, 1.	5.0	21
16	Frequency and Risk Factors Associated with Cord Graft Failure after Transplant with Single-Unit Umbilical Cord Cells Supplemented by Haploidentical Cells with Reduced-Intensity Conditioning. Biology of Blood and Marrow Transplantation, 2016, 22, 1065-1072.	2.0	20
17	Combined Haploidentical and Umbilical Cord Blood Allogeneic Stem Cell Transplantation for High-Risk Lymphoma and Chronic Lymphoblastic Leukemia. Biology of Blood and Marrow Transplantation, 2018, 24, 359-365.	2.0	20
18	Unexpected Toxicities When Nivolumab Was Given as Maintenance Therapy following Allogeneic Stem Cell Transplantation. Biology of Blood and Marrow Transplantation, 2020, 26, 1025-1027.	2.0	20

IF # ARTICLE CITATIONS Reduced-Intensity Allogeneic Transplant for Acute Myeloid Leukemia and Myelodysplastic Syndrome Using Combined CD34-Selected Haploidentical Graft and a Single Umbilical Cord Unit Compared with Matched Unrelated Donor Stem Cells in Older Adults. Biology of Blood and Marrow Transplantation, 2018.24.997-1004 Reduced intensity conditioning for acute myeloid leukemia using melphalan- vs busulfan-based 20 5.2 18 regimens: a CIBMTR report. Blood Advances, 2020, 4, 3180-3190. Novel strategies for immuno-oncology breakthroughs with cell therapy. Biomarker Research, 2021, 9, 6.8 62. Preliminary Results of Balli-01: A Phase I Study of UCART22 (allogeneic engineered T-cells expressing) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 22 1.4 15 Acute Lymphoblastic Leukemia (B-ALL). Blood, 2020, 136, 7-8. No Exit: Identifying Avoidable Terminal Oncology Intensive Care Unit Hospitalizations. Journal of 2.5 Oncology Practice, 2016, 12, e901-e911. Chronic Graft-versus-Host Disease, Nonrelapse Mortality, and Disease Relapse in Older versus Younger Adults Undergoing Matched Allogeneic Peripheral Blood Hematopoietic Cell Transplantation: A Center for International Blood and Marrow Transplant Research Analysis. 24 1.2 13 Transplantation and Cellular Therapy, 2022, 28, 34-42. Expanded indications for allogeneic stem cell transplantation in patients with myeloid malignancies. Current Opinion in Hematology, 2013, 20, 115-122. 2.5 Dose escalation prophylactic donor lymphocyte infusion after T-cell depleted matched related donor allogeneic hematopoietic cell transplantation is feasible and results in higher donor chimerism, 26 2.4 11 faster immune re-constitution, and prolonged progression-free survival. Bone Marrow Transplantation, 2020, 55, 1161-1168. Pembrolizumab for the Treatment of Disease Relapse Following Allogeneic Hematopoietic Cell 1.4 Transplantation. Blood, 2018, 132, 3415-3415. A Phase 1 Study of NKX019, a CD19 Chimeric Antigen Receptor Natural Killer (CAR NK) Cell Therapy, in 28 1.4 11 Subjects with B-Cell Malignancies. Blood, 2021, 138, 3868-3868. Alternative Donor Transplantation—"Mixing and Matching― the Role of Combined Cord Blood and Haplo-Identical Donor Transplantation (Haplo-Cord SCT) as a Treatment Strategy for Patients Lacking 2.3 10 Standard Donors?. Current Hematologic Malignancy Reports, 2015, 10, 1-7. Recommendations and outcomes from a geriatric assessment guided multidisciplinary clinic prior to 30 1.0 10 autologous stem cell transplant in older patients. Journal of Geriatric Oncology, 2021, 12, 585-591. Incidence and predictors of respiratory viral infections by multiplex PCR in allogeneic hematopoietic cell transplant recipients 50 years and older including gériatric assessment. Leukemia and Lymphoma, 2016, 57, 1807-1813. 1.3 A phase 1 study of azacitidine with high-dose cytarabine and mitoxantrone in high-risk acute myeloid 32 5.2 9 leukemia. Blood Advances, 2020, 4, 599-606. A 2:1 randomized, open-label, phase II study of selinexor vs. physician's choice in older patients with 1.3 relapsed or refractory acute myeloid leukemia. Leukemia and Lymphoma, 2021, 62, 1-12. Reduction of Imatinib Concentration After Gastric Bypass Surgery. Blood, 2010, 116, 4948-4948. 34 1.4 9 Preliminary Results from the Flu/Cy/Alemtuzumab Arm of the Phase I BALLI-01 Trial of UCART22, an Anti-CD22 Allogeneic CAR-T Cell Product, in Adult Patients with Relapsed or Refractory (R/R) CD22+ 1.4 B-Cell Acute Lymphoblastic Leukemia (B-ALL). Blood, 2021, 138, 1746-1746. Phase I trial of maintenance selinexor after allogeneic hematopoietic stem cell transplantation for 36 patients with acute myeloid leukemia and myelodysplastic syndrome. Bone Marrow Transplantation, 2.4 5

ΗΟΝGTAO LIU

2020, 55, 2204-2206.

Ηονςτάο Liu

#	Article	IF	CITATIONS
37	Risk classification at diagnosis predicts post-HCT outcomes in intermediate-, adverse-risk, and <i>KMT2A</i> -rearranged AML. Blood Advances, 2022, 6, 828-847.	5.2	5
38	A phase I study of the WT2725 dosing emulsion in patients with advanced malignancies. Scientific Reports, 2021, 11, 22355.	3.3	5
39	Characterization of cancer comorbidity prior to allogeneic hematopoietic cell transplantation. Leukemia and Lymphoma, 2019, 60, 629-638.	1.3	4
40	Efficacy and tolerability of a modified pediatricâ€inspired intensive regimen for acute lymphoblastic leukemia in older adults. EJHaem, 2021, 2, 413-420.	1.0	4
41	Haplo-Cord UCB SCT with Low Cell Dose, Well Matched UCB Units. a Prospective Multicenter Study. Blood, 2014, 124, 1093-1093.	1.4	4
42	Geriatric Assessment (GA) to Predict Survival in Older Allogeneic Hematopoietic Cell Transplantation (HCT) Recipients. Biology of Blood and Marrow Transplantation, 2014, 20, S39-S40.	2.0	3
43	Outcomes of IDH-Mutated Advanced Phase Ph-Negative Myeloproliferative Neoplasms Treated with IDH Inhibitors. Blood, 2019, 134, 4176-4176.	1.4	3
44	Efficacy of Single-Agent Decitabine in Relapsed and Primary Refractory (rel/ref) Acute Myeloid Leukemia (AML). Blood, 2015, 126, 2518-2518.	1.4	3
45	A phase 1 trial utilizing TMI with fludarabine-melphalan in patients with hematologic malignancies undergoing second allo-SCT. Blood Advances, 0, , .	5.2	3
46	Unexpected Toxicities When Nivolumab Was Given after Allogeneic Stem Cell Transplantation. Blood, 2019, 134, 1956-1956.	1.4	2
47	Expanding Use of a Modified Pediatric Intensive Regimen for Acute Lymphoblastic Leukemia (ALL) into an Older Adult Population: Feasibility and Efficacy Results. Blood, 2020, 136, 41-42.	1.4	2
48	The Outcomes of Second Allogeneic Stem Cell Transplantation for Disease Relapse after T Cell Depleted Allogeneic Stem Cell Transplantation: A Single Center Experience-University of Chicago. Blood, 2014, 124, 2509-2509.	1.4	2
49	ls it time to use minimal residual disease to stratify post-remission treatment for acute myeloid leukemia?. Leukemia and Lymphoma, 2015, 56, 3005-3007.	1.3	1
50	Bloodless chimeric antigen receptor (CAR) T-cell therapy in Jehovah's Witnesses. Leukemia and Lymphoma, 2021, 62, 1497-1501.	1.3	1
51	WT1 Peptide Vaccine Is Able to Induce WT1-Specifc Immune Response with TCR Clonal Enrichment to Control Minimal Residual Disease in Patients with Myeloid Leukemia. Blood, 2016, 128, 3984-3984.	1.4	1
52	Comorbidity from Solid Tumor or Hematologic Malignancy Prior to Allogeneic Hematopoietic Cell Transplantation (HCT) May Both Increase Non-Relapse Mortality. Blood, 2016, 128, 5844-5844.	1.4	1
53	A Phase II Prospective Feasibility Study of Clofarabine Cytoreduction Prior to Allogeneic Hematopoietic Cell Transplantation (HCT) for Patients with Relapsed or Refractory Acute Leukemias and Advanced Myelodysplastic Syndromes. Blood, 2011, 118, 496-496.	1.4	0
54	Frequency and Risk Factors of Cord Graft Failure (CGF) Following Reduced Intensity Conditioning Haplo-Cord Hematopoietic Stem Cell Transplantation. Blood, 2014, 124, 2463-2463.	1.4	0

Ηονςτάο Liu

#	Article	IF	CITATIONS
55	Incidence and Predictors of Respiratory Viral Infections By Multi-Plex PCR in Allogeneic Hematopoietic Cell Transplant (HCT) Recipients 50 Years and Older Including Geriatric Assessment (GA). Blood, 2014, 124, 2464-2464.	1.4	0
56	Dose-Escalation Study of Azacitidine Followed By High-Dose Cytarabine (HiDAC) and Mitoxantrone (Mito) for Remission Induction in High-Risk Acute Myeloid Leukemia (AML). Blood, 2015, 126, 3777-3777.	1.4	0
57	Evaluation of a pre-transplant serum biomarker score for allogeneic hematopoietic stem cell transplant (HCT) and association with clinical factors Journal of Clinical Oncology, 2016, 34, e18537-e18537.	1.6	0
58	Final Results from a Phase I Trial Combining Selinexor with High-Dose Cytarabine (HiDAC) and Mitoxantrone (Mito) for Remission Induction in Acute Myeloid Leukemia (AML). Blood, 2018, 132, 4073-4073.	1.4	0
59	Feasibility and Outcomes of T-Cell Depleted Hematopoietic Stem Cell Transplantation in Patients with Relapsed or Refractory AML and High Risk MDS. Blood, 2019, 134, 3324-3324.	1.4	Ο
60	Phase I Trial of a Novel Conditioning Regimen Utilizing Total Marrow Irradiation (TMI) with Fludarabine-Melphalan for Patients with Relapsed Hematologic Malignancies Undergoing Second Allogeneic Stem Cell Transplantation (Allo-SCT). Blood, 2020, 136, 39-40.	1.4	0