

# Tadeusz Osadnik

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3929097/publications.pdf>

Version: 2024-02-01

48  
papers

935  
citations

516710

16  
h-index

501196

28  
g-index

51  
all docs

51  
docs citations

51  
times ranked

1588  
citing authors

#	ARTICLE	IF	CITATIONS
1	A comparison of ST elevation versus non-ST elevation myocardial infarction outcomes in a large registry database. <i>International Journal of Cardiology</i> , 2011, 152, 70-77.	1.7	87
2	The Prognostic Role of Red Blood Cell Distribution Width in Coronary Artery Disease: A Review of the Pathophysiology. <i>Disease Markers</i> , 2015, 2015, 1-12.	1.3	68
3	Red cell distribution width is associated with long-term prognosis in patients with stable coronary artery disease. <i>BMC Cardiovascular Disorders</i> , 2013, 13, 113.	1.7	64
4	Impact of chronic total occlusion artery on 12-month mortality in patients with non-ST-segment elevation myocardial infarction treated by percutaneous coronary intervention (From the PL-ACS). <i>Tj ETQq0 0 0 rgBT / Overlock 10 Tf 50</i>		
5	“Obesity and Insulin Resistance” Is the Component of the Metabolic Syndrome Most Strongly Associated with Oxidative Stress. <i>Antioxidants</i> , 2022, 11, 79.	5.1	49
6	Oxidative Stress in Association with Metabolic Health and Obesity in Young Adults. <i>Oxidative Medicine and Cellular Longevity</i> , 2021, 2021, 1-19.	4.0	40
7	Temporal Trends in the Treatment and Outcomes of Patients With Non-ST-Segment Elevation Myocardial Infarction in Poland from 2004–2010 (from the Polish Registry of Acute Coronary). <i>Tj ETQq1 1 0.784314 rgBT / Overlock 10 Tf 50</i>		
8	Prognostic implications of mean platelet volume on short- and long-term outcomes among patients with non-ST-segment elevation myocardial infarction treated with percutaneous coronary intervention: A single-center large observational study. <i>Platelets</i> , 2016, 27, 452-458.	2.3	33
9	The platelet-to-lymphocyte ratio as a predictor of all-cause mortality in patients with coronary artery disease undergoing elective percutaneous coronary intervention and stent implantation. <i>Journal of the Saudi Heart Association</i> , 2015, 27, 144-151.	0.4	31
10	The Relationships between Polymorphisms in Genes Encoding the Growth Factors TGF- $\beta$ 1, PDGFB, EGF, bFGF and VEGF-A and the Restenosis Process in Patients with Stable Coronary Artery Disease Treated with Bare Metal Stent. <i>PLoS ONE</i> , 2016, 11, e0150500.	2.5	31
11	The prevalence and management of familial hypercholesterolemia in patients with acute coronary syndrome in the Polish tertiary centre: Results from the TERCET registry with 19,781 individuals. <i>Atherosclerosis</i> , 2019, 288, 33-41.	0.8	28
12	Characteristics of lipid profile and effectiveness of management of dyslipidaemia in patients with acute coronary syndromes “ Data from the TERCET registry with 19,287 patients. <i>Pharmacological Research</i> , 2019, 139, 460-466.	7.1	28
13	Metabolically healthy obese and metabolic syndrome of the lean: the importance of diet quality. Analysis of MAGNETIC cohort. <i>Nutrition Journal</i> , 2020, 19, 19.	3.4	27
14	Outcomes of invasive treatment in very elderly Polish patients with non-ST-segment-elevation myocardial infarction from 2003&#8211;2009 (from the PL-ACS registry). <i>Cardiology Journal</i> , 2013, 20, 34-43.	1.2	25
15	Evaluation of dyslipidaemia and the impact of hypolipidemic therapy on prognosis in high and very high risk patients through the Hyperlipidaemia Therapy in tERtiary Cardiological cEnTer (TERCET) Registry. <i>Pharmacological Research</i> , 2018, 132, 204-210.	7.1	20
16	In-Hospital and 12-Month Outcomes After Acute Coronary Syndrome Treatment in Patients Aged <40 Years of Age (from the Polish Registry of Acute Coronary Syndromes). <i>American Journal of Cardiology</i> , 2014, 114, 175-180.	1.6	19
17	Risk-factors associated with extremely high cardiovascular risk of mid- and long-term mortality following myocardial infarction: Analysis of the Hyperlipidaemia Therapy in tERtiary Cardiological cEnTer (TERCET) registry. <i>Atherosclerosis</i> , 2021, 333, 16-23.	0.8	19
18	Predominant location of coronary artery atherosclerosis in the left anterior descending artery. The impact of septal perforators and the myocardial bridging effect. <i>Kardiologia i Torakochirurgia Polska</i> , 2015, 4, 379-385.	0.1	16

#	ARTICLE	IF	CITATIONS
19	Association of Metabolically Healthy and Unhealthy Obesity Phenotypes with Oxidative Stress Parameters and Telomere Length in Healthy Young Adult Men. Analysis of the MAGNETIC Study. <i>Antioxidants</i> , 2021, 10, 93.	5.1	16
20	Comparison of Invasive and Non-Invasive Treatment Strategies in Older Patients With Acute Myocardial Infarction Complicated by Cardiogenic Shock (from the Polish Registry of Acute Coronary) <i>Tj ETQq0 0 OrgBT /Overlock 10 TF</i>		
21	The association of functional polymorphisms in genes encoding growth factors for endothelial cells and smooth muscle cells with the severity of coronary artery disease. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 218.	1.7	14
22	Family History of Premature Coronary Artery Disease (P-CAD) – A Non-Modifiable Risk Factor? Dietary Patterns of Young Healthy Offspring of P-CAD Patients: A Case-Control Study (MAGNETIC Project). <i>Nutrients</i> , 2018, 10, 1488.	4.1	14
23	Metabolic and genetic profiling of young adults with and without a family history of premature coronary heart disease (MAGNETIC). Study design and methodology. <i>Archives of Medical Science</i> , 2019, 15, 590-597.	0.9	14
24	Renal function on admission affects both treatment strategy and long-term outcomes of patients with myocardial infarction (from the Polish Registry of Acute Coronary Syndromes). <i>Kardiologia Polska</i> , 2017, 75, 332-343.	0.6	14
25	Association of Metabolically Healthy and Unhealthy Obesity Phenotype with Markers Related to Obesity, Diabetes among Young, Healthy Adult Men. Analysis of MAGNETIC Study. <i>Life</i> , 2021, 11, 1350.	2.4	14
26	Novel inflammatory biomarkers may reflect subclinical inflammation in young healthy adults with obesity. <i>Endokrynologia Polska</i> , 2019, 70, 135-142.	1.0	12
27	Functional polymorphism rs710218 in the gene coding GLUT1 protein is associated with in-stent restenosis. <i>Biomarkers in Medicine</i> , 2015, 9, 743-750.	1.4	11
28	Prognostic value of neutrophil to lymphocyte ratio in predicting long-term mortality in patients with ischemic and nonischemic heart failure. <i>Polish Archives of Internal Medicine</i> , 2016, 126, 166-173.	0.4	11
29	Comparison of modification of diet in renal disease and chronic kidney disease epidemiology collaboration formulas in predicting long-term outcomes in patients undergoing stent implantation due to stable coronary artery disease. <i>Clinical Research in Cardiology</i> , 2014, 103, 569-576.	3.3	10
30	The Role of Septal Perforators and Myocardial Bridging Effect in Atherosclerotic Plaque Distribution in the Coronary Artery Disease. <i>Polski Przegląd Radiologii i Medycyny Nuklearnej</i> , 2015, 80, 195-201.	1.0	10
31	CARDIAC SURGERY Risk factors for paravalvular leak after transcatheter aortic valve implantation. <i>Kardiochirurgia i Torakochirurgia Polska</i> , 2015, 2, 89-94.	0.1	9
32	Relationship of the rs1799752 polymorphism of the angiotensin-converting enzyme gene and the rs699 polymorphism of the angiotensinogen gene to the process of in-stent restenosis in a population of Polish patients with stable coronary artery disease. <i>Advances in Medical Sciences</i> , 2016, 61, 276-281.	2.1	9
33	Calcium and Phosphate Levels are Among Other Factors Associated with Metabolic Syndrome in Patients with Normal Weight. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020, Volume 13, 1281-1288.	2.4	9
34	Design and rationale of a nationwide screening analysis from the LIPIDOGRAM2015 and LIPIDOGEN2015 studies. <i>Archives of Medical Science</i> , 2020, 18, 604-616.	0.9	9
35	Średnia objętość pętyek krwi i wskaźnik duży jako czynniki prognostyczne choroby wieńcowej i zawału serca. <i>Folia Cardiologica</i> , 2016, 10, 418-422.	0.1	9
36	Who is eligible for randomized trials? A comparison between the exclusion criteria defined by the ISCHEMIA trial and 3102 real-world patients with stable coronary artery disease undergoing stent implantation in a single cardiology center. <i>Trials</i> , 2015, 16, 411.	1.6	8

#	ARTICLE	IF	CITATIONS
37	The Relationship between <i>VEGFA</i> and <i>TGFB1</i> Polymorphisms and Target Lesion Revascularization after Elective Percutaneous Coronary Intervention. <i>Disease Markers</i> , 2017, 2017, 1-8.	1.3	7
38	The role of MGMT polymorphisms (Rs12917 and Rs11016879) in head and neck cancer risk and prognosis. <i>Acta Biochimica Polonica</i> , 2018, 65, 87-92.	0.5	7
39	Genetic and environmental factors associated with homocysteine concentrations in a population of healthy young adults. Analysis of the MAGNETIC study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 939-947.	2.6	7
40	VDR Gene Polymorphisms in Healthy Individuals with Family History of Premature Coronary Artery Disease. <i>Disease Markers</i> , 2021, 2021, 1-9.	1.3	7
41	High progesterone levels are associated with family history of premature coronary artery disease in young healthy adult men. <i>PLoS ONE</i> , 2019, 14, e0215302.	2.5	5
42	Comparative effect of nutraceuticals on lipid profile: a protocol for systematic review and network meta-analysis. <i>BMJ Open</i> , 2020, 10, e032755.	1.9	5
43	High baseline fibrinogen concentration as a risk factor of no tissue reperfusion in ST-segment elevation acute myocardial infarction treated with successful primary percutaneous coronary intervention. <i>Kardiologia Polska</i> , 2006, 64, 967-72; discussion 973-4.	0.6	5
44	New Variants of the Cytochrome P450 2R1 (CYP2R1) Gene in Individuals with Severe Vitamin D-Activating Enzyme 25(OH)D Deficiency. <i>Biomolecules</i> , 2021, 11, 1867.	4.0	5
45	The CTGF gene -945 G/C polymorphism is associated with target lesion revascularization for in-stent restenosis. <i>Experimental and Molecular Pathology</i> , 2021, 118, 104598.	2.1	4
46	Is GLUT1 a potential target for in-stent restenosis treatment?. <i>International Journal of Cardiology</i> , 2016, 223, 199-200.	1.7	2
47	Polymorphisms of genes coding for telomerase reverse transcriptase and telomerase RNA component and the need for target lesion revascularization after percutaneous coronary intervention. <i>Polish Archives of Internal Medicine</i> , 2021, 131, 299-301.	0.4	0
48	Genotype-phenotype correlations in Polish patients with hypertrophic cardiomyopathy: Preliminary report. <i>Kardiologia Polska</i> , 2022, 80, 482-484.	0.6	0