Ademir Araujo

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3933738/ademir-araujo-publications-by-year.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

144 2,250 25 42 g-index

161 2,762 3 5.14 ext. papers ext. citations avg, IF L-index

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 144 | Ecosystem functions in different physiognomies of Cerrado through the Rapid Ecosystem Function Assessment (REFA) <i>Anais Da Academia Brasileira De Ciencias</i> , 2022 , 94, e20200457 | 1.4 | O |
| 143 | Dataset for effects of the transition from dry forest to pasture on diversity and structure of bacterial communities in Northeastern Brazil <i>Data in Brief</i> , 2022 , 41, 107842 | 1.2 | |
| 142 | Cowpea nodules host a similar bacterial community regardless of soil properties. <i>Applied Soil Ecology</i> , 2022 , 172, 104354 | 5 | O |
| 141 | Bacillus subtilis rhizobacteria ameliorate heat stress in the common bean. Rhizosphere, 2022, 21, 10047 | 23.5 | 2 |
| 140 | Land degradation affects the microbial communities in the Brazilian Caatinga biome. <i>Catena</i> , 2022 , 211, 105961 | 5.8 | O |
| 139 | Genetically related genotypes of cowpea present similar bacterial community in the rhizosphere <i>Scientific Reports</i> , 2022 , 12, 3472 | 4.9 | 0 |
| 138 | Rhizobacteria and arbuscular mycorrhizal fungus presented distinct and specific effects on soybean growth when inoculated with organic compost. <i>Rhizosphere</i> , 2022 , 22, 100513 | 3.5 | O |
| 137 | Forest-to-pasture conversion modifies the soil bacterial community in Brazilian dry forest Caatinga. <i>Science of the Total Environment</i> , 2021 , 810, 151943 | 10.2 | 0 |
| 136 | Distinct taxonomic composition of soil bacterial community across a native gradient of Cerrado-Ecotone-Caatinga. <i>Applied Soil Ecology</i> , 2021 , 161, 103874 | 5 | 3 |
| 135 | Plant growth-promoting bacteria improve growth and nitrogen metabolism in maize and sorghum. <i>Theoretical and Experimental Plant Physiology</i> , 2021 , 33, 249-260 | 2.4 | 1 |
| 134 | Bacillus subtilis changes the root architecture of soybean grown on nutrient-poor substrate. <i>Rhizosphere</i> , 2021 , 18, 100348 | 3.5 | 3 |
| 133 | Response of soil bacterial communities to the application of the herbicides imazethapyr and flumyzin. <i>European Journal of Soil Biology</i> , 2021 , 102, 103252 | 2.9 | 7 |
| 132 | Distinct bacterial community structure and composition along different cowpea producing ecoregions in Northeastern Brazil. <i>Scientific Reports</i> , 2021 , 11, 831 | 4.9 | 4 |
| 131 | Seed size influences the promoting activity of rhizobia on plant growth, nodulation and N fixation in lima bean. <i>Ciencia Rural</i> , 2021 , 51, | 1.3 | 2 |
| 130 | Structure and diversity of bacterial community in semiarid soils cultivated with prickly-pear cactus (Opuntia ficus-indica (L.) Mill.). <i>Anais Da Academia Brasileira De Ciencias</i> , 2021 , 93, e20190183 | 1.4 | 1 |
| 129 | Grazing exclusion regulates bacterial community in highly degraded semiarid soils from the Brazilian Caatinga biome. <i>Land Degradation and Development</i> , 2021 , 32, 2210-2225 | 4.4 | 5 |
| 128 | Dynamics of bacterial and archaeal communities along the composting of tannery sludge. <i>Environmental Science and Pollution Research</i> , 2021 , 28, 64295-64306 | 5.1 | 2 |

| 127 | Characterization of edaphic fauna in different monocultures in Savanna of Piau Darazilian Journal of Biology, 2021 , 81, 657-664 | 1.5 | 2 | |
|-----|---|------|----|--|
| 126 | Arbuscular mycorrhizal community in soil from different Brazilian Cerrado physiognomies. <i>Rhizosphere</i> , 2021 , 19, 100375 | 3.5 | 2 | |
| 125 | Microbial co-occurrence network and its key microorganisms in soil with permanent application of composted tannery sludge. <i>Science of the Total Environment</i> , 2021 , 789, 147945 | 10.2 | 4 | |
| 124 | Diversity, structure, and composition of plant growth-promoting bacteria in soil from Brazilian Cerrado. <i>Rhizosphere</i> , 2021 , 20, 100435 | 3.5 | 1 | |
| 123 | Cover crops shape the soil bacterial community in a tropical soil under no-till. <i>Applied Soil Ecology</i> , 2021 , 168, 104166 | 5 | 2 | |
| 122 | Responses of soil microbial biomass and enzyme activity to herbicides imazethapyr and flumioxazin. <i>Scientific Reports</i> , 2020 , 10, 7694 | 4.9 | 10 | |
| 121 | Phytotoxicity and cytogenotoxicity of composted tannery sludge. <i>Environmental Science and Pollution Research</i> , 2020 , 27, 34495-34502 | 5.1 | 3 | |
| 120 | Polyphasic characterization of nitrogen-fixing and co-resident bacteria in nodules of Phaseolus lunatus inoculated with soils from Piau [^] [State, Northeast Brazil. <i>Symbiosis</i> , 2020 , 80, 279-292 | 3 | 7 | |
| 119 | Nodule microbiome from cowpea and lima bean grown in composted tannery sludge-treated soil. <i>Applied Soil Ecology</i> , 2020 , 151, 103542 | 5 | 12 | |
| 118 | Inoculation of rhizobia increases lima bean (Phaseolus lunatus) yield in soils from Piau [^] Land Cear [^] Land Cear States, Brazil. <i>Revista Ceres</i> , 2020 , 67, 419-423 | 0.7 | 2 | |
| 117 | Soil microbial biomass and enzyme activity in six Brazilian oxisols under cropland and native vegetation. <i>Bragantia</i> , 2020 , 79, 623-629 | 1.2 | 1 | |
| 116 | Changes in Soil Properties and Crop Yield as a Function of Early Desiccation of Pastures. <i>Journal of Soil Science and Plant Nutrition</i> , 2020 , 20, 840-848 | 3.2 | 4 | |
| 115 | Diversity and structure of bacterial community in rhizosphere of lima bean. <i>Applied Soil Ecology</i> , 2020 , 150, 103490 | 5 | 11 | |
| 114 | Analysis and advanced characterization of municipal solid waste vermicompost maturity for a green environment. <i>Journal of Environmental Management</i> , 2020 , 255, 109914 | 7.9 | 36 | |
| 113 | Long-term effect of composted tannery sludge on soil chemical and biological parameters. <i>Environmental Science and Pollution Research</i> , 2020 , 27, 41885-41892 | 5.1 | 6 | |
| 112 | Bacillus subtilis can modulate the growth and root architecture in soybean through volatile organic compounds. <i>Theoretical and Experimental Plant Physiology</i> , 2020 , 32, 99-108 | 2.4 | 14 | |
| 111 | Maize rhizosphere soil stimulates greater soil microbial biomass and enzyme activity leading to subsequent enhancement of cowpea growth. <i>Environmental Sustainability</i> , 2019 , 2, 89-94 | 2.9 | 3 | |
| 110 | Bacterial community associated with rhizosphere of maize and cowpea in a subsequent cultivation. <i>Applied Soil Ecology</i> , 2019 , 143, 26-34 | 5 | 16 | |

| 109 | Microbial biomass and organic matter in an oxisol under application of biochar. <i>Bragantia</i> , 2019 , 78, 10 | 9-1.18 | 7 |
|-----|--|--------|----|
| 108 | Dynamics of archaeal community in soil with application of composted tannery sludge. <i>Scientific Reports</i> , 2019 , 9, 7347 | 4.9 | 6 |
| 107 | Capability of plant growth-promoting bacteria in chromium-contaminated soil after application of composted tannery sludge. <i>Annals of Microbiology</i> , 2019 , 69, 665-671 | 3.2 | 4 |
| 106 | Edaphic fauna in a vegetation gradient in the Sete Cidades National Park. <i>Brazilian Journal of Biology</i> , 2019 , 79, 45-51 | 1.5 | 5 |
| 105 | Bacillus subtilis ameliorates water stress tolerance in maize and common bean. <i>Journal of Plant Interactions</i> , 2019 , 14, 432-439 | 3.8 | 14 |
| 104 | Changes on microbial C and enzyme activities in soil with amendment of composted tannery sludge after 9 years. <i>International Journal of Recycling of Organic Waste in Agriculture</i> , 2019 , 8, 501-505 | 3.1 | 1 |
| 103 | Sugarcane inoculated with endophytic diazotrophic bacteria: effects on yield, biological nitrogen fixation and industrial characteristics. <i>Anais Da Academia Brasileira De Ciencias</i> , 2019 , 91, e20180990 | 1.4 | 8 |
| 102 | Soil microbial C:N:P ratio across physiognomies of Brazilian Cerrado Soil microbial biomass across a gradient of preserved native Cerrado. <i>Anais Da Academia Brasileira De Ciencias</i> , 2019 , 91, e20190049 | 1.4 | 4 |
| 101 | Nodulation, nitrogen uptake and growth of lima bean in a composted tannery sludge-treated soil. <i>Ciencia Rural</i> , 2019 , 49, | 1.3 | 2 |
| 100 | Archaea diversity in vegetation gradients from the Brazilian Cerrado. <i>Brazilian Journal of Microbiology</i> , 2018 , 49, 522-528 | 2.2 | 16 |
| 99 | Responses of soil bacterial community after seventh yearly applications of composted tannery sludge. <i>Geoderma</i> , 2018 , 318, 1-8 | 6.7 | 24 |
| 98 | The Impact of Pasture Systems on Soil Microbial Biomass and Community-level Physiological Profiles. <i>Land Degradation and Development</i> , 2018 , 29, 284-291 | 4.4 | 16 |
| 97 | Chemical variables influencing microbial properties in composted tannery sludge-treated soil. <i>International Journal of Environmental Science and Technology</i> , 2018 , 15, 1793-1800 | 3.3 | 3 |
| 96 | Less abundant bacterial groups are more affected than the most abundant groups in composted tannery sludge-treated soil. <i>Scientific Reports</i> , 2018 , 8, 11755 | 4.9 | 8 |
| 95 | Bacillus subtilis improves maize tolerance to salinity. Ciencia Rural, 2018, 48, | 1.3 | 10 |
| 94 | Chromium accumulation in maize and cowpea after successive applications of composted tannery sludge. <i>Acta Scientiarum - Agronomy</i> , 2018 , 40, 35361 | 0.6 | 10 |
| 93 | Soil Microbial Biomass Across a Gradient of Preserved Native Cerrado. <i>Floresta E Ambiente</i> , 2018 , 25, | 1 | 2 |
| 92 | Protist species richness and soil microbiome complexity increase towards climax vegetation in the Brazilian Cerrado. <i>Communications Biology</i> , 2018 , 1, 135 | 6.7 | 31 |

(2016-2018)

| 91 | Responses of microbial biomass, available phosphorus, and sugarcane yield after filter cake amendment in a tropical soil. <i>Australian Journal of Crop Science</i> , 2018 , 12, 552-556 | 0.5 | 1 |
|----|--|------|-----|
| 90 | Ten years of application of sewage sludge on tropical soil. A balance sheet on agricultural crops and environmental quality. <i>Science of the Total Environment</i> , 2018 , 643, 1493-1501 | 10.2 | 46 |
| 89 | Distinct bacterial communities across a gradient of vegetation from a preserved Brazilian Cerrado. <i>Antonie Van Leeuwenhoek</i> , 2017 , 110, 457-469 | 2.1 | 26 |
| 88 | Symbiotic performance, nitrogen flux and growth of lima bean (Phaseolus lunatus L.) varieties inoculated with different indigenous strains of rhizobia. <i>Symbiosis</i> , 2017 , 73, 117-124 | 3 | 7 |
| 87 | Two new begomoviruses that infect non-cultivated malvaceae in Brazil. <i>Archives of Virology</i> , 2017 , 162, 1795-1797 | 2.6 | 4 |
| 86 | Fungal diversity in soils across a gradient of preserved Brazilian Cerrado. <i>Journal of Microbiology</i> , 2017 , 55, 273-279 | 3 | 13 |
| 85 | Time-dependent effect of composted tannery sludge on the chemical and microbial properties of soil. <i>Ecotoxicology</i> , 2017 , 26, 1366-1377 | 2.9 | 9 |
| 84 | Rhizobial Diversity for Tropical Pulses and Forage and Tree Legumes in Brazil 2017 , 135-151 | | |
| 83 | Biological properties of disturbed and undisturbed Cerrado sensu stricto from Northeast Brazil. Brazilian Journal of Biology, 2017 , 77, 16-21 | 1.5 | 2 |
| 82 | Complete genome sequence of a new bipartite begomovirus infecting Macroptilium lathyroides in Brazil. <i>Archives of Virology</i> , 2017 , 162, 3551-3554 | 2.6 | 5 |
| 81 | Historical and recent land use affects ecosystem functions in subtropical grasslands in Brazil. <i>Ecosphere</i> , 2017 , 8, e02032 | 3.1 | 14 |
| 80 | Nodulation ability in different genotypes of Phaseolus lunatus by rhizobia from California agricultural soils. <i>Symbiosis</i> , 2017 , 73, 7-14 | 3 | 5 |
| 79 | Diversity of plant growth-promoting bacteria associated with sugarcane. <i>Genetics and Molecular Research</i> , 2017 , 16, | 1.2 | 11 |
| 78 | Agroecological Responses of Heavy Metal Pollution with Special Emphasis on Soil Health and Plant Performances. <i>Frontiers in Environmental Science</i> , 2017 , 5, | 4.8 | 111 |
| 77 | T-RFLP analysis of soil bacterial structure from Cerrado within the Sete Cidades National Park, Brazil. <i>Neotropical Biodiversity</i> , 2016 , 2, 163-170 | 0.7 | 1 |
| 76 | Land-Use Type Effects on Soil Organic Carbon and Microbial Properties in a Semi-arid Region of Northeast Brazil. <i>Land Degradation and Development</i> , 2016 , 27, 171-178 | 4.4 | 65 |
| 75 | Penetration resistance and density of a yellow oxissol under conventional management at different ages. <i>Bioscience Journal</i> , 2016 , 32, 115-122 | 2 | 2 |
| 74 | Short Communication: Soil carbon pools in different pasture systems. <i>Spanish Journal of Agricultural Research</i> , 2016 , 14, e11SC01 | 1.1 | 1 |

| 73 | Soil properties and cowpea yield after six years of consecutive amendment of composted tannery sludge. <i>Acta Scientiarum - Agronomy</i> , 2016 , 38, 407 | 0.6 | 4 |
|----|---|------|----|
| 72 | Caracteriza [^] [] B e Diverg [^] []cia Gen [^] []ica de Popula [^] [] Bs de Casearia grandiflora no Cerrado Piauiense. <i>Floresta E Ambiente</i> , 2016 , 23, 387-396 | 1 | 6 |
| 71 | Repeated application of composted tannery sludge affects differently soil microbial biomass, enzymes activity, and ammonia-oxidizing organisms. <i>Environmental Science and Pollution Research</i> , 2016 , 23, 19193-200 | 5.1 | 11 |
| 70 | HLA-B*15:04:04, a novel HLA allele identified during proficiency testing in Brazil. <i>Hla</i> , 2016 , 88, 200-1 | 1.9 | 3 |
| 69 | Biological response of using municipal solid waste compost in agriculture as fertilizer supplement. <i>Reviews in Environmental Science and Biotechnology</i> , 2016 , 15, 677-696 | 13.9 | 48 |
| 68 | Bradyrhizobium sp. inoculation ameliorates oxidative protection in cowpea subjected to long-term composted tannery sludge amendment. <i>European Journal of Soil Biology</i> , 2016 , 76, 35-45 | 2.9 | 6 |
| 67 | Soil Enzymatic Activity in Eucalyptus Grandis Plantations of Different Ages. <i>Land Degradation and Development</i> , 2016 , 27, 77-82 | 4.4 | 17 |
| 66 | Soil microbial properties after 5 years of consecutive amendment with composted tannery sludge. <i>Environmental Monitoring and Assessment</i> , 2015 , 187, 4153 | 3.1 | 24 |
| 65 | Soil Surface-Active Fauna in Degraded and Restored Lands of Northeast Brazil. <i>Land Degradation and Development</i> , 2015 , 26, 1-8 | 4.4 | 35 |
| 64 | CHROMIUM IN SOIL ORGANIC MATTER AND COWPEA AFTER FOUR CONSECUTIVE ANNUAL APPLICATIONS OF COMPOSTED TANNERY SLUDGE. <i>Revista Brasileira De Ciencia Do Solo</i> , 2015 , 39, 297- | 302 | 4 |
| 63 | Soil microbial biomass in an agroforestry system of Northeast Brazil. <i>Tropical Grasslands - Forrajes Tropicales</i> , 2015 , 3, 41 | 1.8 | 14 |
| 62 | Diversity of native rhizobia-nodulating Phaseolus lunatus in Brazil. Legume Research, 2015, 38, | 1 | 2 |
| 61 | LEITURAS DE CLOROFILA E TEORES DE N EM FASES FENOL [^] GICAS DO MILHO. <i>Colloquium Agrariae</i> , 2015 , 11, 57-63 | 0.2 | 2 |
| 60 | Effect of Utilization of Organic Waste as Agricultural Amendment on Soil Microbial Biomass. <i>Annual Research & Review in Biology</i> , 2015 , 7, 155-162 | 0.8 | 7 |
| 59 | Soil Microbial Biomass After Three-Year Consecutive Composted Tannery Sludge Amendment. <i>Pedosphere</i> , 2014 , 24, 469-475 | 5 | 13 |
| 58 | Nitrogen application and inoculation with Rhizobium tropici on common bean in the fall/winter. <i>African Journal of Agricultural Research Vol Pp</i> , 2014 , 9, 3156-3163 | 0.5 | 4 |
| 57 | Soil organic matter pools in a tropical savanna under agroforestry system in Northeastern Brazil. <i>Revista Arvore,</i> 2014 , 38, 711-723 | 1 | 9 |
| 56 | Soil microbial properties in Eucalyptus grandis plantations of different ages. <i>Journal of Soil Science and Plant Nutrition</i> , 2014 , 0-0 | 3.2 | 2 |

| 55 | Chromium, Cadmium, Nickel, and Lead in a Tropical Soil after 3 Years of Consecutive Applications of Composted Tannery Sludge. <i>Communications in Soil Science and Plant Analysis</i> , 2014 , 45, 1658-1666 | 1.5 | 3 |
|----|---|-------------|----|
| 54 | Soil microbial biomass after two years of the consecutive application of composted tannery sludge - doi: 10.4025/actasciagron.v36i1.17160. <i>Acta Scientiarum - Agronomy</i> , 2014 , 36, 35 | 0.6 | 9 |
| 53 | Heavy metals and yield of cowpea cultivated under composted tannery sludge amendment. <i>Acta Scientiarum - Agronomy</i> , 2014 , 36, 443 | 0.6 | 6 |
| 52 | Resposta do milho verde ^ [inocula ^ [i] com Azospirillum brasilense e n ^ [ileis de nitrog ^ [ilo. <i>Ciencia Rural</i> , 2014 , 44, 1556-1560 | 1.3 | 5 |
| 51 | Biofertilizers on soil microbial biomass and activity. <i>Revista Brasileirade Ciencias Agrarias</i> , 2014 , 9, 545-5 | 49 1 | 2 |
| 50 | Soil bacterial diversity in degraded and restored lands of Northeast Brazil. <i>Antonie Van Leeuwenhoek</i> , 2014 , 106, 891-9 | 2.1 | 27 |
| 49 | Estado nutricional e produ [^] [] [] da pimenteira com uso de biofertilizantes l [^] []quidos. <i>Revista Brasileira De Engenharia Agricola E Ambiental</i> , 2014 , 18, 1241-1246 | 0.9 | 5 |
| 48 | Short communication. Growth and nodulation of cowpea after 5 years of consecutive composted tannery sludge amendment. <i>Spanish Journal of Agricultural Research</i> , 2014 , 12, 1175 | 1.1 | 9 |
| 47 | Soil microbial properties and temporal stability in degraded and restored lands of Northeast Brazil. <i>Soil Biology and Biochemistry</i> , 2013 , 66, 175-181 | 7.5 | 74 |
| 46 | Biomassa e atividade microbiana do solo sob pastagem em sistemas de monocultura e silvipastoril. <i>Semina:Ciencias Agrarias</i> , 2013 , 34, 2727 | 0.6 | 3 |
| 45 | Biological Nitrogen Fixation: Importance, Associated Diversity, and Estimates 2013 , 267-289 | | 11 |
| 44 | Heavy metals in cowpea (Vigna unguiculata L.) after tannery sludge compost amendment. <i>Chilean Journal of Agricultural Research</i> , 2013 , 73, 282-287 | 1.9 | 9 |
| 43 | Microbiological process in agroforestry systems. A review. <i>Agronomy for Sustainable Development</i> , 2012 , 32, 215-226 | 6.8 | 26 |
| 42 | Impact of Land Degradation on Soil Microbial Biomass and Activity in Northeast Brazil. <i>Pedosphere</i> , 2012 , 22, 88-95 | 5 | 45 |
| 41 | Soil microbial biomass and organic matter fractions during transition from conventional to organic farming systems. <i>Geoderma</i> , 2012 , 170, 227-231 | 6.7 | 99 |
| 40 | Soil microbial biomass and activity under natural and regenerated forests and conventional sugarcane plantations in Brazil. <i>Geoderma</i> , 2012 , 189-190, 257-261 | 6.7 | 42 |
| 39 | Sistemas agroflorestais e seus efeitos sobre os atributos qu [^] Enicos em Argissolo Vermelho-Amarelo do Cerrado piauiense. <i>Revista Brasileira De Engenharia Agricola E Ambiental</i> , 2012 , 16, 730-738 | 0.9 | 18 |
| 38 | Fungos micorr [®] Zicos arbusculares como indicadores da recupera [®] D de [®] Deas degradadas no nordeste do Brasil. <i>Revista Ciencia Agronomica</i> , 2012 , 43, 648-657 | 1 | 9 |

| 37 | Tannery sludge compost amendment rates on soil microbial biomass of two different soils. European Journal of Soil Biology, 2011 , 47, 146-151 | 2.9 | 50 |
|----|---|-------|-----|
| 36 | Doses de paclobutrazol sobre a biomassa microbiana do solo. <i>Semina:Ciencias Agrarias</i> , 2011 , 31, 1349 | 0.6 | 1 |
| 35 | Emerg [^] Bcia e crescimento inicial de pl [^] Btulas de pimenta ornamental e celosia em substrato [^] Ibase de composto de lodo de curtume. <i>Ciencia Rural</i> , 2011 , 41, 412-417 | 1.3 | 3 |
| 34 | Genetic diversity among native isolates of rhizobia from Phaseolus lunatus. <i>Annals of Microbiology</i> , 2011 , 61, 437-444 | 3.2 | 10 |
| 33 | Management of urban solid waste: Vermicomposting a sustainable option. <i>Resources, Conservation and Recycling</i> , 2011 , 55, 719-729 | 11.9 | 116 |
| 32 | Efici [^] līcia simbi [^] līca de isolados de riz [^] līco noduladores de feij [^] līchava (Phaseolus lunatus L.). <i>Revista Brasileira De Ciencia Do Solo</i> , 2011 , 35, 751-757 | 1.5 | 9 |
| 31 | Growth, nodulation and nitrogen fixation of cowpea in soils amended with composted tannery sludge. <i>Revista Brasileira De Ciencia Do Solo</i> , 2011 , 35, 1865-1871 | 1.5 | 6 |
| 30 | Bacillus subtilis e aduba^ 🛘 🗗 nitrogenada na produtividade do milho. <i>Revista Brasileirade Ciencias Agrarias</i> , 2011 , 6, 657-66 | 1.1 | 5 |
| 29 | Coinocula [^] [] B̄ riz [^] B̄io e Bacillus subtilis em feij [^] B̄-caupi e leucena: efeito sobre a nodula [^] [] B̄, a fixa [^] [] B̄ de N2 e o crescimento das plantas. <i>Ciencia Rural</i> , 2010 , 40, 182-185 | 1.3 | 4 |
| 28 | Soil organic carbon and biological indicators in an Acrisol under tillage systems and organic management in north-eastern Brazil. <i>Soil Research</i> , 2010 , 48, 258 | 1.8 | 28 |
| 27 | Municipal solid waste compost amendment in agricultural soil: changes in soil microbial biomass. <i>Reviews in Environmental Science and Biotechnology</i> , 2010 , 9, 41-49 | 13.9 | 35 |
| 26 | Is the microwave irradiation a suitable method for measuring soil microbial biomass?. <i>Reviews in Environmental Science and Biotechnology</i> , 2010 , 9, 317-321 | 13.9 | 5 |
| 25 | The effect of converting tropical native savanna to Eucalyptus grandis forest on soil microbial biomass. <i>Land Degradation and Development</i> , 2010 , 21, 540-545 | 4.4 | 25 |
| 24 | Effect of different tannery sludge compost amendment rates on growth, biomass accumulation and yield responses of Capsicum plants. <i>Waste Management</i> , 2010 , 30, 1976-80 | 8.6 | 47 |
| 23 | Soil microbial biomass in organic farming system. <i>Ciencia Rural</i> , 2010 , 40, 2419-2426 | 1.3 | 13 |
| 22 | Changes in soil microbial biomass and activity in different Brazilian pastures. <i>Spanish Journal of Agricultural Research</i> , 2010 , 8, 1253 | 1.1 | 15 |
| 21 | Soil Microbial Activity in Conventional and Organic Agricultural Systems. Sustainability, 2009 , 1, 268-276 | 5 3.6 | 51 |
| 20 | Effect of paclobutrazol on microbial biomass, respiration and cellulose decomposition in soil. <i>European Journal of Soil Biology</i> , 2009 , 45, 235-238 | 2.9 | 14 |

| 19 | Responses of soil microbial biomass and activity for practices of organic and conventional farming systems in Piau [^] [state, Brazil. <i>European Journal of Soil Biology</i> , 2008 , 44, 225-230 | 2.9 | 92 |
|----|---|--------------|-----|
| 18 | Avalia^ 🛮 🖟 de indicadores biol^ ḡicos de qualidade do solo sob sistemas de cultivo convencional e org^ ਜico de frutas. <i>Ciencia E Agrotecnologia</i> , 2008 , 32, 353-359 | 1.6 | 17 |
| 17 | Inocula^ [] B e aduba^ [] B nitrogenada sobre a nodula^ [] B e a produtividade de gr^ Bs de feij^ B-caupi. <i>Ciencia Rural</i> , 2008 , 38, 2037-2041 | 1.3 | 6 |
| 16 | INOCULA [^] [] D E ADUBA [^] [] D MINERAL EM FEIJ [^] D-CAUPI: EFEITOS NA NODULA [^] [] D, CRESCIMENTO E PRODUTIVIDADE. <i>Scientia Agraria</i> , 2008 , 9, 469 | | 5 |
| 15 | Effect of composted textile sludge on growth, nodulation and nitrogen fixation of soybean and cowpea. <i>Bioresource Technology</i> , 2007 , 98, 1028-32 | 11 | 38 |
| 14 | Ontogenia da nodula^ 🛘 🗗 em duas cultivares de feij^ 🗗 caupi. <i>Ciencia Rural</i> , 2007 , 37, 561-564 | 1.3 | 7 |
| 13 | Microbial biomass and activity in a Brazilian soil amended with untreated and composted textile sludge. <i>Chemosphere</i> , 2006 , 64, 1043-6 | 8.4 | 27 |
| 12 | Efeito da adi^ 🛮 B̄ de lodo de curtume na fertilidade do solo, nodula^ 🗓 B̄ e rendimento de mat^ Ēia seca do Caupi. <i>Ciencia E Agrotecnologia</i> , 2006 , 30, 1071-1076 | 1.6 | 5 |
| 11 | Sobreviv^ ficia e nodula^ [] fi do Rhizobium tropici em sementes de feij^ fi tratadas com fungicidas. <i>Ciencia Rural</i> , 2006 , 36, 973-976 | 1.3 | 7 |
| 10 | Plant bioassays to assess toxicity of textile sludge compost. <i>Scientia Agricola</i> , 2005 , 62, 286-290 | 2.5 | 80 |
| 9 | Utiliza^ 🛮 🗗 de nitrog^ 🗟 io pelo trigo cultivado em solo fertilizado com adubo verde (Crotalaria juncea) e/ou ur^ 🗗 . <i>Ciencia Rural</i> , 2005 , 35, 284-289 | 1.3 | 10 |
| 8 | Composto de lodo t^ ⊠til em pl^ fitulas de soja e trigo. <i>Pesquisa Agropecuaria Brasileira</i> , 2005 , 40, 549-5 | 54 .8 | 7 |
| 7 | Effect of glyphosate on the microbial activity of two Brazilian soils. Chemosphere, 2003, 52, 799-804 | 8.4 | 207 |
| 6 | INFLU^ NCIA DE BACILLUS SUBTILIS NA ECLOS^ D, ORIENTA^ D D E INFEC^ D DE HETERODERA GLYCINES EM SOJA. <i>Ciencia Rural</i> , 2002 , 32, 197-203 | 1.3 | 13 |
| 5 | Plant growth-promoting endophytic bacteria on maize and sorghum1. <i>Pesquisa Agropecuaria Tropical</i> ,49, | 1.2 | 11 |
| 4 | Assessment of the phenotypic diversity in natural populations of Annona coriacea Mart.: implications for breeding. <i>Genetic Resources and Crop Evolution</i> ,1 | 2 | O |
| 3 | Plant growth-promoting rhizobacteria effect on maize growth and microbial biomass in a chromium-contaminated soil. <i>Bragantia</i> ,80, | 1.2 | 2 |
| 2 | Plant growth-promoting bacteria increase the yield of green maize and sweet sorghum. <i>Journal of Plant Nutrition</i> ,1-11 | 2.3 | |

Enzymatic Stoichiometry in Soils from Physiognomies of Brazilian Cerrado. *Journal of Soil Science and Plant Nutrition*,1

3.2 0