

# Kunio Ishii

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3935090/publications.pdf>

Version: 2024-02-01

43  
papers

652  
citations

516710

16  
h-index

610901

24  
g-index

44  
all docs

44  
docs citations

44  
times ranked

825  
citing authors

#	ARTICLE	IF	CITATIONS
1	Role of Epoxyeicosatrienoic Acids in Acetylcholine-Induced Dilation of Rat Retinal Arterioles &lt;i>in Vivo</i>. <i>Biological and Pharmaceutical Bulletin</i> , 2021, 44, 82-87.	1.4	5
2	L-Citrulline ameliorates the attenuation of acetylcholine-induced vasodilation of retinal arterioles in diabetic rats. <i>Heliyon</i> , 2021, 7, e06532.	3.2	3
3	Metformin Protects against NMDA-Induced Retinal Injury through the MEK/ERK Signaling Pathway in Rats. <i>International Journal of Molecular Sciences</i> , 2021, 22, 4439.	4.1	15
4	Involvement of Gap Junctions in Acetylcholine-Induced Endothelium-Derived Hyperpolarization-Type Dilation of Retinal Arterioles in Rats. <i>Biological and Pharmaceutical Bulletin</i> , 2021, 44, 1860-1865.	1.4	0
5	4-Aminopyridine, a Voltage-Gated K <sup>+</sup> Channel Inhibitor, Attenuates Nitric Oxide-Mediated Vasodilation of Retinal Arterioles in Rats. <i>Biological and Pharmaceutical Bulletin</i> , 2020, 43, 1123-1127.	1.4	4
6	Involvement of matrix metalloproteinases in capillary degeneration following NMDA-induced neurotoxicity in the neonatal rat retina. <i>Experimental Eye Research</i> , 2019, 182, 101-108.	2.6	4
7	Probucol Slows the Progression of Cataracts in Streptozotocin-Induced Hyperglycemic Rats. <i>Pharmacology</i> , 2019, 103, 212-219.	2.2	4
8	Iron-chelating agents attenuate NMDA-Induced neuronal injury via reduction of oxidative stress in the rat retina. <i>Experimental Eye Research</i> , 2018, 171, 30-36.	2.6	33
9	Retinal neuronal cell loss prevents abnormal retinal vascular growth in a rat model of retinopathy of prematurity. <i>Experimental Eye Research</i> , 2018, 168, 115-127.	2.6	8
10	Transient phenotypic changes in endothelial cells and pericytes in neonatal mouse retina following short-term blockade of vascular endothelial growth factor receptors. <i>Developmental Dynamics</i> , 2018, 247, 699-711.	1.8	3
11	Anti-angiogenic effects of valproic acid in a mouse model of oxygen-induced retinopathy. <i>Journal of Pharmacological Sciences</i> , 2018, 138, 203-208.	2.5	11
12	Anti-cataract Effect of Resveratrol in High-Glucose-Treated Streptozotocin-Induced Diabetic Rats. <i>Biological and Pharmaceutical Bulletin</i> , 2018, 41, 1586-1592.	1.4	29
13	A delay in vascularization induces abnormal astrocyte proliferation and migration in the mouse retina. <i>Developmental Dynamics</i> , 2017, 246, 186-200.	1.8	15
14	Stimulation of $\mu$ -opioid receptors dilates retinal arterioles by neuronal nitric oxide synthase-derived nitric oxide in rats. <i>European Journal of Pharmacology</i> , 2017, 803, 124-129.	3.5	10
15	MEK/ERK- and calcineurin/NFAT-mediated mechanism of cerebral hyperemia and brain injury following NMDA receptor activation. <i>Biochemical and Biophysical Research Communications</i> , 2017, 488, 329-334.	2.1	2
16	Stimulation of $\beta$ <sub>1</sub> - and $\beta$ <sub>2</sub> -adrenoceptors dilates retinal blood vessels in rats. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2017, 390, 527-533.	3.0	7
17	L-Citrulline ameliorates cerebral blood flow during cortical spreading depression in rats: Involvement of nitric oxide- and prostanoids-mediated pathway. <i>Journal of Pharmacological Sciences</i> , 2017, 133, 146-155.	2.5	8
18	Opioid receptor activation is involved in neuroprotection induced by TRPV1 channel activation against excitotoxicity in the rat retina. <i>European Journal of Pharmacology</i> , 2017, 812, 57-63.	3.5	12

#	ARTICLE	IF	CITATIONS
19	Probucol prevents the attenuation of $\text{I}^{22}$ -adrenoceptor-mediated vasodilation of retinal arterioles in diabetic rats. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2017, 390, 1247-1253.	3.0	5
20	Activation inhibitors of nuclear factor kappa B protect neurons against the NMDA-induced damage in the rat retina. <i>Journal of Pharmacological Sciences</i> , 2017, 135, 72-80.	2.5	24
21	Apelin-36 is protective against N-methyl-D-aspartic-acid-induced retinal ganglion cell death in the mice. <i>European Journal of Pharmacology</i> , 2016, 791, 213-220.	3.5	21
22	Protective effects of PF4708671 against N-methyl-D-aspartic acid-induced retinal damage in rats. <i>Fundamental and Clinical Pharmacology</i> , 2016, 30, 529-536.	1.9	4
23	Short-term treatment with VEGF receptor inhibitors induces retinopathy of prematurity-like abnormal vascular growth in neonatal rats. <i>Experimental Eye Research</i> , 2016, 143, 120-131.	2.6	16
24	Retinal region-dependent susceptibility of capillaries to high-concentration oxygen exposure and vascular endothelial growth factor receptor inhibition in neonatal mice. <i>Journal of Pharmacological Sciences</i> , 2015, 129, 107-118.	2.5	4
25	Impaired retinal vasodilator response to acetylcholine in a rat model of NMDA-induced retinal degeneration. <i>Journal of Pharmacological Sciences</i> , 2015, 127, 211-216.	2.5	7
26	Structural and functional changes in retinal vasculature induced by retinal ischemia-reperfusion in rats. <i>Experimental Eye Research</i> , 2015, 135, 134-145.	2.6	53
27	Age-Dependent Changes in the Severity of Capillary Degeneration in Rat Retina Following N-Methyl-D-Aspartate-Induced Neurotoxicity. <i>Current Eye Research</i> , 2015, 40, 549-553.	1.5	11
28	P2X7 receptor antagonists protect against N-methyl-d-aspartic acid-induced neuronal injury in the rat retina. <i>European Journal of Pharmacology</i> , 2015, 756, 52-58.	3.5	30
29	High-mobility group Box-1 is involved in NMDA-induced retinal injury the in rat retina. <i>Experimental Eye Research</i> , 2015, 137, 63-70.	2.6	17
30	L-Citrulline dilates rat retinal arterioles via nitric oxide- and prostaglandin-dependent pathways in vivo. <i>Journal of Pharmacological Sciences</i> , 2015, 127, 419-423.	2.5	18
31	Involvement of prostaglandin I2 in nitric oxide-induced vasodilation of retinal arterioles in rats. <i>European Journal of Pharmacology</i> , 2015, 764, 249-255.	3.5	16
32	Regression of retinal capillaries following N-methyl-D-aspartate-induced neurotoxicity in the neonatal rat retina. <i>Journal of Neuroscience Research</i> , 2015, 93, 380-390.	2.9	13
33	Effects of mTOR inhibition on normal retinal vascular development in the mouse. <i>Experimental Eye Research</i> , 2014, 129, 127-134.	2.6	18
34	Activation of the TRPV1 channel attenuates N-methyl-d-aspartic acid-induced neuronal injury in the rat retina. <i>European Journal of Pharmacology</i> , 2014, 733, 13-22.	3.5	31
35	Effects of pre- and post-natal treatment with KRN633, an inhibitor of vascular endothelial growth factor receptor tyrosine kinase, on retinal vascular development and patterning in mice. <i>Experimental Eye Research</i> , 2014, 120, 127-137.	2.6	18
36	Hydrogen sulfide attenuates NMDA-induced neuronal injury via its anti-oxidative activity in the rat retina. <i>Experimental Eye Research</i> , 2014, 120, 90-96.	2.6	41

#	ARTICLE	IF	CITATIONS
37	Small Molecule Cyclin-Dependent Kinase Inhibitors Protect Against Neuronal Cell Death in the Ischemic-Reperfused Rat Retina. <i>Journal of Ocular Pharmacology and Therapeutics</i> , 2011, 27, 419-425.	1.4	22
38	Histological Protection by Donepezil Against Neurodegeneration Induced by Ischemiaâ€“Reperfusion in the Rat Retina. <i>Journal of Pharmacological Sciences</i> , 2010, 112, 327-335.	2.5	27
39	Protective effect of all-trans retinoic acid on NMDA-induced neuronal cell death in rat retina. <i>European Journal of Pharmacology</i> , 2010, 635, 56-61.	3.5	26
40	Histological protection by cilnidipine, a dual L/N-type Ca <sup>2+</sup> channel blocker, against neurotoxicity induced by ischemiaâ€“reperfusion in rat retina. <i>Experimental Eye Research</i> , 2009, 88, 974-982.	2.6	39
41	Mexiletine inhibits pharmacological actions of salbutamol through blockade of Î² <sub>2</sub> -adrenoceptors in bovine tracheal smooth muscle. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2001, 364, 409-413.	3.0	2
42	Possible involvement of endothelium-derived hyperpolarizing factor (EDHF) in the depressor responses to platelet activating factor (PAF) in rats. <i>British Journal of Pharmacology</i> , 2000, 131, 1113-1120.	5.4	5
43	Lidocaine potentiates the relaxant effects of cAMP-elevating agents in bovine tracheal smooth muscle. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2000, 361, 605-609.	3.0	10