

Ida Tylleskar

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3937422/publications.pdf>

Version: 2024-02-01

11
papers

119
citations

1477746

6
h-index

1372195

10
g-index

12
all docs

12
docs citations

12
times ranked

83
citing authors

#	ARTICLE	IF	CITATIONS
1	Pharmacokinetics of a new, nasal formulation of naloxone. <i>European Journal of Clinical Pharmacology</i> , 2017, 73, 555-562.	0.8	31
2	REBOARREST, resuscitative endovascular balloon occlusion of the aorta in non-traumatic out-of-hospital cardiac arrest: a study protocol for a randomised, parallel group, clinical multicentre trial. <i>Trials</i> , 2021, 22, 511.	0.7	22
3	Pharmacokinetics and -dynamics of intramuscular and intranasal naloxone: an explorative study in healthy volunteers. <i>European Journal of Clinical Pharmacology</i> , 2018, 74, 873-883.	0.8	16
4	Pharmacokinetics of a novel, approved, 1.4 mg intranasal naloxone formulation for reversal of opioid overdose—a randomized controlled trial. <i>Addiction</i> , 2019, 114, 859-867.	1.7	15
5	Prehospital naloxone administration — what influences choice of dose and route of administration?. <i>BMC Emergency Medicine</i> , 2020, 20, 71.	0.7	10
6	Comparison of intranasal and intramuscular naloxone in opioid overdoses managed by ambulance staff: a double-blind, randomised, controlled trial. <i>Addiction</i> , 2022, 117, 1658-1667.	1.7	9
7	Pharmacodynamics and arteriovenous difference of intravenous naloxone in healthy volunteers exposed to remifentanyl. <i>European Journal of Clinical Pharmacology</i> , 2018, 74, 1547-1553.	0.8	8
8	The pharmacokinetic interaction between nasally administered naloxone and the opioid remifentanyl in human volunteers. <i>European Journal of Clinical Pharmacology</i> , 2021, 77, 1901-1908.	0.8	5
9	NTNU intranasal naloxone trial (NINA-1) study protocol for a double-blind, double-dummy, non-inferiority randomised controlled trial comparing intranasal 1.4 mg to intramuscular 0.8 mg naloxone for prehospital use. <i>BMJ Open</i> , 2020, 10, e041556.	0.8	1
10	NTNU intranasal naloxone trial (NINA-1) study protocol for a double-blind, double-dummy, non-inferiority randomised controlled trial comparing intranasal 1.4 mg to intramuscular 0.8 mg naloxone for prehospital use. <i>BMJ Open</i> , 2020, 10, e041556.	0.8	1
11	Naloxone administration—no balance without titration. <i>Addiction</i> , 2022, 117, 2750-2751.	1.7	1