

Koji Suzuki

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3944933/koji-suzuki-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

151
papers

3,577
citations

30
h-index

53
g-index

158
ext. papers

4,067
ext. citations

3.8
avg, IF

4.49
L-index

#	Paper	IF	Citations
151	Maternal High-Fructose Corn Syrup consumption causes insulin resistance and hyperlipidemia in offspring via DNA methylation of the Ppar α promoter region.. <i>Journal of Nutritional Biochemistry</i> , 2022 , 103, 108951	6.3	1
150	Association of drinking behaviors with DNA methylation levels in leukocytes among the general Japanese population.. <i>American Journal of Drug and Alcohol Abuse</i> , 2022 , 1-9	3.7	0
149	Increased risk of cancer mortality by smoking-induced aryl hydrocarbon receptor repressor DNA hypomethylation in Japanese population: A long-term cohort study.. <i>Cancer Epidemiology</i> , 2022 , 78, 102162	2.8	1
148	Circulating microRNA-27a and -133a are negatively associated with incident hypertension: A five-year longitudinal population-based study.. <i>Biomarkers</i> , 2022 , 1-16	2.6	2
147	DNA methylation is associated with muscle loss in community-dwelling older men -the Yakumo study- : a preliminary experimental study.. <i>Nagoya Journal of Medical Science</i> , 2022 , 84, 60-68	0.7	
146	High-fructose corn syrup intake has stronger effects on the transcription level of hepatic lipid metabolism-related genes, via DNA methylation modification, in childhood and adolescence than in other generations.. <i>Life Sciences</i> , 2022 , 120638	6.8	0
145	Maternal fructose intake predisposes rat offspring to metabolic disorders via abnormal hepatic programming. <i>FASEB Journal</i> , 2021 , 35, e22030	0.9	0
144	DNA methylation level of the gene encoding thioredoxin-interacting protein in peripheral blood cells is associated with metabolic syndrome in the Japanese general population. <i>Endocrine Journal</i> , 2021 ,	2.9	1
143	Dairy products and the risk of developing prostate cancer: A large-scale cohort study (JACC Study) in Japan. <i>Cancer Medicine</i> , 2021 , 10, 7298-7307	4.8	2
142	Association between circulating vascular-related microRNAs and an increase in blood pressure: a 5-year longitudinal population-based study. <i>Journal of Hypertension</i> , 2021 , 39, 84-89	1.9	7
141	A preliminary examination of the association between locomotive syndrome and circulating miRNA-199 in community-dwelling people: The Yakumo study. <i>Journal of Orthopaedic Science</i> , 2021 , 27, 696-696	1.6	
140	Global DNA hypermethylation in peripheral blood mononuclear cells and cardiovascular disease risk: a population-based propensity score-matched cohort study. <i>Journal of Epidemiology and Community Health</i> , 2021 , 75, 890-895	5.1	0
139	The development of knee osteoarthritis and serum carotenoid levels among community-dwelling people in Japan. <i>Modern Rheumatology</i> , 2021 , 1-8	3.3	0
138	Analysis of Repeated Measurements of Serum Carotenoid Levels and All-Cause and Cause-Specific Mortality in Japan. <i>JAMA Network Open</i> , 2021 , 4, e2113369	10.4	3
137	Dietary fish and Ω 3 polyunsaturated fatty acids are associated with leukocyte ABCA1 DNA methylation levels. <i>Nutrition</i> , 2021 , 81, 110951	4.8	8
136	Circulating miR-21, miR-29a, and miR-126 are associated with premature death risk due to cancer and cardiovascular disease: the JACC Study. <i>Scientific Reports</i> , 2021 , 11, 5298	4.9	11
135	Maternal fructose consumption downregulates hippocampal catalase expression via DNA methylation in rat offspring. <i>Nutrition Research</i> , 2021 , 92, 40-48	4	5

134	DNA methylation as a mediator of associations between the environment and chronic diseases: A scoping review on application of mediation analysis. <i>Epigenetics</i> , 2021 , 1-27	5.7	3
133	Association between circulating microRNAs and changes in kidney function: A five-year prospective study among Japanese adults without CKD. <i>Clinica Chimica Acta</i> , 2021 , 521, 97-103	6.2	1
132	Association of smoking habits with TXNIP DNA methylation levels in leukocytes among general Japanese population. <i>PLoS ONE</i> , 2020 , 15, e0235486	3.7	6
131	Association between dietary inflammatory index and serum C-reactive protein concentrations in the Japan Collaborative Cohort Study. <i>Nagoya Journal of Medical Science</i> , 2020 , 82, 237-249	0.7	4
130	Maternal fructose consumption down-regulates Lxra expression via miR-206-mediated regulation. <i>Journal of Nutritional Biochemistry</i> , 2020 , 82, 108386	6.3	9
129	Associations of Serum MicroRNA with Bone Mineral Density in Community-Dwelling Subjects: The Yakumo Study. <i>BioMed Research International</i> , 2020 , 2020, 5047243	3	3
128	Associations of Circulating MicroRNAs (miR-17, miR-21, and miR-150) and Chronic Kidney Disease in a Japanese Population. <i>Journal of Epidemiology</i> , 2020 , 30, 177-182	3.4	10
127	Associations between dietary vitamin intake, ABCA1 gene promoter DNA methylation, and lipid profiles in a Japanese population. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1213-1219	7	16
126	Carotid artery plaque screening using abdominal aortic calcification on lumbar radiographs. <i>PLoS ONE</i> , 2019 , 14, e0209175	3.7	5
125	Dietary Inflammatory Index Is Associated with Risk of All-Cause and Cardiovascular Disease Mortality but Not with Cancer Mortality in Middle-Aged and Older Japanese Adults. <i>Journal of Nutrition</i> , 2019 , 149, 1451-1459	4.1	14
124	Cluster of differentiation 36 gene polymorphism (rs1761667) is associated with dietary MUFA intake and hypertension in a Japanese population. <i>British Journal of Nutrition</i> , 2019 , 121, 1215-1222	3.6	6
123	Dietary vegetable intake is inversely associated with ATP-binding cassette protein A1 (ABCA1) DNA methylation levels among Japanese women. <i>Nutrition</i> , 2019 , 65, 1-5	4.8	8
122	Maternal high-fructose intake increases circulating corticosterone levels via decreased adrenal corticosterone clearance in adult offspring. <i>Journal of Nutritional Biochemistry</i> , 2019 , 67, 44-50	6.3	8
121	Associations of serum microRNA-20a, -27a, and -103a with cognitive function in a Japanese population: The Yakumo study. <i>Archives of Gerontology and Geriatrics</i> , 2019 , 82, 155-160	4	13
120	Establishment of a simpler method for measuring HDL-microRNAs. <i>Annals of Clinical Biochemistry</i> , 2019 , 56, 49-55	2.2	1
119	Maternal fructose-induced oxidative stress occurs and epigenetic regulation in offspring hippocampi. <i>FASEB Journal</i> , 2019 , 33, 11431-11442	0.9	12
118	Utility of the Serum Cystatin C Level for Diagnosis of Osteoporosis among Middle-Aged and Elderly People. <i>BioMed Research International</i> , 2019 , 2019, 5046852	3	9
117	Circulating microRNAs (miR-126, miR-197, and miR-223) are associated with chronic kidney disease among elderly survivors of the Great East Japan Earthquake. <i>BMC Nephrology</i> , 2019 , 20, 474	2.7	11

116	Association of circulating miR-20a, miR-27a, and miR-126 with non-alcoholic fatty liver disease in general population. <i>Scientific Reports</i> , 2019 , 9, 18856	4.9	11
115	Human serum albumin redox state is associated with decreased renal function in a community-dwelling population. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 316, F214-F218	4.3	3
114	Relationship between locomotive syndrome and body composition among community-dwelling middle-age and elderly individuals in Japan: The Yakumo study. <i>Modern Rheumatology</i> , 2019 , 29, 491-495	3.3	18
113	Maternal fructose intake disturbs ovarian estradiol synthesis in rats. <i>Life Sciences</i> , 2018 , 202, 117-123	6.8	12
112	Serum cystatin C level is associated with locomotive syndrome risk and can be an early predictor in community-living people: The Yakumo study. <i>Modern Rheumatology</i> , 2018 , 28, 1035-1040	3.3	17
111	Association of subcutaneous and visceral fat with circulating microRNAs in a middle-aged Japanese population. <i>Annals of Clinical Biochemistry</i> , 2018 , 55, 437-445	2.2	12
110	Oxidized human serum albumin as a possible correlation factor for atherosclerosis in a rural Japanese population: the results of the Yakumo Study. <i>Environmental Health and Preventive Medicine</i> , 2018 , 23, 1	4.2	19
109	Serum levels of carotenoids in patients with osteonecrosis of the femoral head are lower than in healthy, community-living people. <i>Journal of Orthopaedic Surgery</i> , 2018 , 26, 2309499018770927	1.4	3
108	Excess maternal fructose consumption impairs hippocampal function in offspring via epigenetic modification of BDNF promoter. <i>FASEB Journal</i> , 2018 , 32, 2549-2562	0.9	29
107	Association of Adiponectin With Cancer and All-Cause Mortality in a Japanese Community-Dwelling Elderly Cohort: A Case-Cohort Study. <i>Journal of Epidemiology</i> , 2018 , 28, 367-372	3.4	3
106	Relationship between Long Interspersed Nuclear Element-1 DNA Methylation in Leukocytes and Dyslipidemia in the Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 1231-1239	4	11
105	Fructose intake during gestation and lactation differentially affects the expression of hippocampal neurosteroidogenic enzymes in rat offspring. <i>Endocrine Research</i> , 2017 , 42, 71-77	1.9	12
104	Stability of serum high-density lipoprotein-microRNAs for preanalytical conditions. <i>Annals of Clinical Biochemistry</i> , 2017 , 54, 134-142	2.2	12
103	Alcohol consumption and mortality from aortic disease among Japanese men: The Japan Collaborative Cohort study. <i>Atherosclerosis</i> , 2017 , 266, 64-68	3.1	7
102	The impact of musculoskeletal diseases on mortality-comparison with internal diseases: A 15-year longitudinal study. <i>Journal of Orthopaedic Science</i> , 2017 , 22, 1126-1131	1.6	9
101	p53 and ki67 as biomarkers in determining response to chemoprevention for oral leukoplakia. <i>Journal of Oral Pathology and Medicine</i> , 2017 , 46, 346-352	3.3	9
100	Fructose consumption induces hypomethylation of hepatic mitochondrial DNA in rats. <i>Life Sciences</i> , 2016 , 149, 146-52	6.8	30
99	Diabetes Mellitus and Risk of Colorectal Cancer Mortality in Japan: the Japan Collaborative Cohort Study. <i>Asian Pacific Journal of Cancer Prevention</i> , 2016 , 17, 4681-4688	1.7	8

98	Longitudinal study of circulating miR-122 in a rat model of non-alcoholic fatty liver disease. <i>Clinica Chimica Acta</i> , 2015 , 446, 267-71	6.2	56
97	Carrier frequency of the GJB2 mutations that cause hereditary hearing loss in the Japanese population. <i>Journal of Human Genetics</i> , 2015 , 60, 613-7	4.3	12
96	High fructose consumption induces DNA methylation at PPAR α and CPT1A promoter regions in the rat liver. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 468, 185-9	3.4	53
95	Treatment of oral leukoplakia with a low-dose of beta-carotene and vitamin C supplements: a randomized controlled trial. <i>International Journal of Cancer</i> , 2015 , 136, 1708-17	7.5	29
94	Maternal fructose consumption alters messenger RNA expression of hippocampal StAR, PBR, P450(11 β) 11 β HSD, and 17 β HSD in rat offspring. <i>Nutrition Research</i> , 2015 , 35, 259-64	4	16
93	Low intake of vegetables and fruits and risk of colorectal cancer: the Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2014 , 24, 353-60	3.4	12
92	Association between serum levels of carotenoids and serum asymmetric dimethylarginine levels in Japanese subjects. <i>Journal of Epidemiology</i> , 2014 , 24, 250-7	3.4	7
91	Coffee consumption and risk of colorectal cancer: the Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2014 , 24, 370-8	3.4	30
90	A cross-sectional study to find out the relationship of methylenetetrahydrofolate reductase (MTHFR) C677T genotype with plasma levels of folate and total homocysteine by daily folate intake in Japanese. <i>Journal of Nutritional Science and Vitaminology</i> , 2014 , 60, 231-8	1.1	6
89	The expression of groups IIE and V phospholipase A2 is associated with an increased expression of osteogenic molecules in human calcified aortic valves. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 1308-25	4	8
88	Cigarette smoking and other risk factors for kidney cancer death in a Japanese population: Japan Collaborative Cohort Study for evaluation of cancer risk (JACC study). <i>Asian Pacific Journal of Cancer Prevention</i> , 2014 , 14, 6523-8	1.7	20
87	Risk factors for renal cell carcinoma in a Japanese population. <i>Asian Pacific Journal of Cancer Prevention</i> , 2014 , 15, 9065-70	1.7	18
86	Associations between circulating microRNAs (miR-21, miR-34a, miR-122 and miR-451) and non-alcoholic fatty liver. <i>Clinica Chimica Acta</i> , 2013 , 424, 99-103	6.2	216
85	Cohort profile of the Japan Collaborative Cohort Study at final follow-up. <i>Journal of Epidemiology</i> , 2013 , 23, 227-32	3.4	95
84	Body mass index and weight change during adulthood are associated with increased mortality from liver cancer: the JACC Study. <i>Journal of Epidemiology</i> , 2013 , 23, 219-26	3.4	9
83	Association of serum carotenoid levels with urinary albumin excretion in a general Japanese population: the Yakumo study. <i>Journal of Epidemiology</i> , 2013 , 23, 451-6	3.4	9
82	Association of serum carotenoid levels with N-terminal pro-brain-type natriuretic peptide: a cross-sectional study in Japan. <i>Journal of Epidemiology</i> , 2013 , 23, 163-8	3.4	8
81	The stiffness parameter β assessed by an ultrasonic phase-locked echo-tracking system is associated with plaque formation in the common carotid artery. <i>Journal of Medical Ultrasonics (2001)</i> , 2012 , 39, 3-9	1.4	6

80	SIRTUIN 1 gene polymorphisms are associated with cholesterol metabolism and coronary artery calcification in Japanese hemodialysis patients. <i>Journal of Renal Nutrition</i> , 2012 , 22, 114-9	3	46
79	Sirtuin 1 gene polymorphisms are associated with body fat and blood pressure in Japanese. <i>Translational Research</i> , 2011 , 157, 339-47	11	51
78	The effect of β -carotene on lumbar osteophyte formation. <i>Spine</i> , 2011 , 36, 2293-8	3.3	25
77	Inverse association of serum carotenoids with prevalence of metabolic syndrome among Japanese. <i>Clinical Nutrition</i> , 2011 , 30, 369-75	5.9	42
76	Reduction of liver stiffness by antiviral therapy in chronic hepatitis B. <i>Journal of Gastroenterology</i> , 2011 , 46, 1324-34	6.9	44
75	Do musculoskeletal degenerative diseases affect mortality and cause of death after 10 years in Japan?. <i>Journal of Bone and Mineral Metabolism</i> , 2011 , 29, 217-23	2.9	21
74	Relationship of sFas with metabolic risk factors and their clusters. <i>European Journal of Clinical Investigation</i> , 2010 , 40, 527-33	4.6	8
73	Association of serum carotenoids with high molecular weight adiponectin and inflammation markers among Japanese subjects. <i>Clinica Chimica Acta</i> , 2010 , 411, 1330-4	6.2	26
72	Relationship between serum levels of insulin-like growth factors and subsequent risk of cancer mortality: findings from a nested case-control study within the Japan Collaborative Cohort Study. <i>Cancer Epidemiology</i> , 2010 , 34, 279-84	2.8	7
71	Association of serum carotenoids, retinol, and tocopherols with radiographic knee osteoarthritis: possible risk factors in rural Japanese inhabitants. <i>Journal of Orthopaedic Science</i> , 2010 , 15, 477-84	1.6	19
70	A new index for non-invasive assessment of liver fibrosis. <i>World Journal of Gastroenterology</i> , 2010 , 16, 4809-16	5.6	19
69	Possible protective effect of serum beta-carotene levels on the association between interleukin-1B C-31T polymorphism and hypertension in a Japanese population. <i>Clinical Nutrition</i> , 2009 , 28, 198-202	5.9	7
68	Comparison of urinary concentrations of 3-phenoxybenzoic acid among general residents in rural and suburban areas and employees of pest control firms. <i>International Archives of Occupational and Environmental Health</i> , 2009 , 82, 1173-8	3.2	17
67	Relationship between dietary habits and urinary concentrations of 3-phenoxybenzoic acid in a middle-aged and elderly general population in Japan. <i>Environmental Health and Preventive Medicine</i> , 2009 , 14, 173-9	4.2	24
66	Common defects of ABCG2, a high-capacity urate exporter, cause gout: a function-based genetic analysis in a Japanese population. <i>Science Translational Medicine</i> , 2009 , 1, 5ra11	17.5	289
65	Urinary excretion of 3-phenoxybenzoic acid in middle-aged and elderly general population of Japan. <i>Environmental Research</i> , 2009 , 109, 175-80	7.9	48
64	Significant association between methylenetetrahydrofolate reductase 677T allele and hyperuricemia among adult Japanese subjects. <i>Nutrition Research</i> , 2009 , 29, 710-5	4	11
63	Cigarette smoking and serum soluble Fas levels: Findings from the JACC study. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2009 , 679, 79-83	3	1

62	Association between Interleukin-1B C-31T polymorphism and obesity in Japanese. <i>Journal of Epidemiology</i> , 2009 , 19, 131-5	3.4	7
61	Association of serum retinol and carotenoids with insulin-like growth factors and insulin-like growth factor binding protein-3 among control subjects of a nested case-control study in the Japan Collaborative Cohort Study. <i>Asian Pacific Journal of Cancer Prevention</i> , 2009 , 10 Suppl, 29-35	1.7	2
60	Prostate cancer risk in relation to insulin-like growth factor (IGF)-I and IGF-binding protein-3: A nested case-control study in large scale cohort study in Japan. <i>Asian Pacific Journal of Cancer Prevention</i> , 2009 , 10 Suppl, 57-61	1.7	12
59	Significant association between Helicobacter pylori infection and serum C-reactive protein. <i>International Journal of Medical Sciences</i> , 2008 , 5, 224-9	3.7	27
58	Association of serum NO(x) level with clustering of metabolic syndrome components in middle-aged and elderly general populations in Japan. <i>Environmental Health and Preventive Medicine</i> , 2008 , 13, 36-42	4.2	15
57	Smoking and serum CA19-9 levels according to Lewis and secretor genotypes. <i>International Journal of Cancer</i> , 2008 , 123, 2880-4	7.5	29
56	Lung cancer mortality and body mass index in a Japanese cohort: findings from the Japan Collaborative Cohort Study (JACC Study). <i>Cancer Causes and Control</i> , 2007 , 18, 229-34	2.8	11
55	Association between decreased kidney function and endotoxin receptor CD14 C-159T polymorphism among Japanese health check-up examinees. <i>Renal Failure</i> , 2007 , 29, 967-72	2.9	6
54	Dietary fiber and risk of colorectal cancer in the Japan collaborative cohort study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 668-75	4	61
53	Association of a Polymorphism in the Ornithine Decarboxylase Gene with Whole Blood Polyamine Concentrations in a Non-smoking Healthy Population. <i>Journal of Health Science</i> , 2007 , 53, 406-412		2
52	Associations between serum C-reactive protein (CRP) levels and polymorphisms of CRP, interleukin 1B, and tumor necrosis factor genes among Japanese health checkup examinees. <i>Asian Pacific Journal of Cancer Prevention</i> , 2007 , 8, 87-92	1.7	5
51	Health conditions and mortality in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). <i>Asian Pacific Journal of Cancer Prevention</i> , 2007 , 8 Suppl, 25-34	1.7	14
50	Association of abdominal circumference with serum nitric oxide concentration in healthy population. <i>Environmental Health and Preventive Medicine</i> , 2006 , 11, 321-5	4.2	7
49	Association of abdominal obesity with decreased serum levels of carotenoids in a healthy Japanese population. <i>Clinical Nutrition</i> , 2006 , 25, 780-9	5.9	44
48	Serum heat shock protein 70 levels and lung cancer risk: a case-control study nested in a large cohort study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006 , 15, 1733-7	4	42
47	Prospective cohort study of the risk of prostate cancer among rotating-shift workers: findings from the Japan collaborative cohort study. <i>American Journal of Epidemiology</i> , 2006 , 164, 549-55	3.8	291
46	Alcohol consumption and lung cancer mortality in Japanese men: results from Japan collaborative cohort (JACC) study. <i>Journal of Epidemiology</i> , 2006 , 16, 49-56	3.4	18
45	Association of Abdominal Circumference with Serum Nitric Oxide Concentration in Healthy Population. <i>Environmental Health and Preventive Medicine</i> , 2006 , 11, 321-325	4.2	

44	Blood Pressure, Levels of Serum Lipids, Liver Enzymes and Blood Glucose by Aldehyde Dehydrogenase 2 and Drinking Habit in Japanese Men. <i>Environmental Health and Preventive Medicine</i> , 2006 , 11, 82-88	4.2	
43	Genotype announcement to Japanese smokers who attended a health checkup examination. <i>Journal of Epidemiology</i> , 2006 , 16, 45-7	3.4	9
42	Cardiovascular disease mortality and serum carotenoid levels: a Japanese population-based follow-up study. <i>Journal of Epidemiology</i> , 2006 , 16, 154-60	3.4	82
41	Serum carotenoids and other antioxidative substances associated with urothelial cancer risk in a nested case-control study in Japanese men. <i>Journal of Urology</i> , 2005 , 173, 1502-6	2.5	10
40	Possible interactions of the endothelial constitutive nitric oxide synthase genotype with alcohol drinking and walking time for high serum uric acid levels among Japanese. <i>Metabolism: Clinical and Experimental</i> , 2005 , 54, 1302-8	12.7	6
39	Leptin is associated with an increased female colorectal cancer risk: a nested case-control study in Japan. <i>Oncology</i> , 2005 , 68, 454-61	3.6	87
38	Perceived psychologic stress and colorectal cancer mortality: findings from the Japan Collaborative Cohort Study. <i>Psychosomatic Medicine</i> , 2005 , 67, 72-7	3.7	26
37	Lung cancer mortality and serum levels of carotenoids, retinol, tocopherols, and folic acid in men and women: a case-control study nested in the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S140-9	3.4	37
36	Updated information on risk factors for lung cancer: findings from the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S134-9	3.4	19
35	Colorectal cancer and serum C-reactive protein levels: a case-control study nested in the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S185-9	3.4	46
34	Association of serum phytoestrogen concentration and dietary habits in a sample set of the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S196-202	3.4	31
33	Association of serum carotenoid concentration and dietary habits among the JACC Study subjects. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S220-7	3.4	5
32	Medical history of circulatory diseases and colorectal cancer death in the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S168-72	3.4	6
31	Alcohol consumption and colorectal cancer risk: findings from the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S173-9	3.4	17
30	Glucose intolerance and colorectal cancer risk in a nested case-control study among Japanese People. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 2, S180-4	3.4	8
29	Serum levels of polyunsaturated fatty acids and risk of colorectal cancer: a prospective study. <i>American Journal of Epidemiology</i> , 2005 , 161, 462-71	3.8	62
28	Serum carotenoids, retinol, and tocopherols, and colorectal cancer risk in a Japanese cohort: effect modification by sex for carotenoids. <i>Nutrition and Cancer</i> , 2005 , 51, 13-24	2.8	24
27	Cancer mortality and serum levels of carotenoids, retinol, and tocopherol: a population-based follow-up study of inhabitants of a rural area of Japan. <i>Asian Pacific Journal of Cancer Prevention</i> , 2005 , 6, 10-5	1.7	29

26	Diet and colorectal cancer mortality: results from the Japan Collaborative Cohort Study. <i>Nutrition and Cancer</i> , 2004 , 50, 23-32	2.8	64
25	A prospective study on the possible association between having children and colon cancer risk: findings from the JACC Study. <i>Cancer Science</i> , 2004 , 95, 243-7	6.9	14
24	A prospective study of reproductive and menstrual factors and colon cancer risk in Japanese women: findings from the JACC study. <i>Cancer Science</i> , 2004 , 95, 602-7	6.9	16
23	Serum phytoestrogens and prostate cancer risk in a nested case-control study among Japanese men. <i>Cancer Science</i> , 2004 , 95, 65-71	6.9	126
22	Serum oxidized low-density lipoprotein levels and risk of colorectal cancer: a case-control study nested in the Japan Collaborative Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2004 , 13, 1781-7	4	29
21	The relationship between smoking habits and serum levels of 8-OHdG, oxidized LDL antibodies, Mn-SOD and carotenoids in rural Japanese residents. <i>Journal of Epidemiology</i> , 2003 , 13, 29-37	3.4	35
20	Smoking and colorectal cancer in a non-Western population: a prospective cohort study in Japan. <i>Journal of Epidemiology</i> , 2003 , 13, 323-32	3.4	25
19	Attributable and absolute risk of lung cancer death by smoking status: findings from the Japan Collaborative Cohort Study. <i>International Journal of Cancer</i> , 2003 , 105, 249-54	7.5	52
18	Serum carotenoids and mortality from lung cancer: a case-control study nested in the Japan Collaborative Cohort (JACC) study. <i>Cancer Science</i> , 2003 , 94, 57-63	6.9	48
17	Relationship between obesity and serum markers of oxidative stress and inflammation in Japanese. <i>Asian Pacific Journal of Cancer Prevention</i> , 2003 , 4, 259-66	1.7	59
16	Serum antioxidants and subsequent mortality rates of all causes or cancer among rural Japanese inhabitants. <i>International Journal for Vitamin and Nutrition Research</i> , 2002 , 72, 237-50	1.7	11
15	Serum insulin-like growth factors, insulin-like growth factor-binding protein-3, and risk of lung cancer death: a case-control study nested in the Japan Collaborative Cohort (JACC) Study. <i>Japanese Journal of Cancer Research</i> , 2002 , 93, 1279-86		38
14	Mortality rates from cancer or all causes and SOD activity level and Zn/Cu ratio in peripheral blood: population-based follow-up study. <i>Journal of Epidemiology</i> , 2002 , 12, 14-21	3.4	12
13	Relationship between serum carotenoids and hyperglycemia: a population-based cross-sectional study. <i>Journal of Epidemiology</i> , 2002 , 12, 357-66	3.4	41
12	Validity and reliability of single-item questions about physical activity. <i>Journal of Epidemiology</i> , 2001 , 11, 211-8	3.4	51
11	The effect of serum carotenoids on atrophic gastritis among the inhabitants of a rural area in Hokkaido, Japan. <i>Environmental Health and Preventive Medicine</i> , 2001 , 6, 184-8	4.2	4
10	TT virus genotype changes frequently in multiply transfused patients with hemophilia but rarely in patients with chronic hepatitis C and in healthy subjects. <i>Transfusion</i> , 2001 , 41, 1130-5	2.9	8
9	Decrease in risk of lung cancer death in males after smoking cessation by age at quitting: findings from the JACC study. <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 821-8		24

8	Dietary habits and risk of lung cancer death in a large-scale cohort study (JACC Study) in Japan by sex and smoking habit. <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 1259-69		53
7	Effects of polyunsaturated fatty acids on atrophic gastritis in a Japanese population. <i>Cancer Letters</i> , 2001 , 163, 171-8	9.9	4
6	The Effect of Serum Carotenoids on Atrophic Gastritis Among the Inhabitants of a Rural Area in Hokkaido, Japan. <i>Environmental Health and Preventive Medicine</i> , 2001 , 6, 184-188	4.2	
5	The Risk of Helicobacter Pylori Infection and Atrophic Gastritis from Food and Drink Intake: a Cross-sectional Study in Hokkaido, Japan. <i>Asian Pacific Journal of Cancer Prevention</i> , 2000 , 1, 147-156	1.7	10
4	A study on serum carotenoid levels in breast cancer patients of Indian women in Chennai (Madras), India. <i>Journal of Epidemiology</i> , 1999 , 9, 306-14	3.4	14
3	Relationship between serum carotenoid levels and cancer death rates in the residents, living in a rural area of Hokkaido, Japan. <i>Journal of Epidemiology</i> , 1997 , 7, 1-8	3.4	19
2	Detection of non-heme iron proteins following polyacrylamide gel electrophoresis. <i>Electrophoresis</i> , 1985 , 6, 351-352	3.6	1
1	STUDIES ON UROLITHIASIS. <i>Japanese Journal of Urology</i> , 1981 , 72, 842-855		