

Thereza Christina M de Lima

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3953431/thereza-christina-m-de-lima-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

97
papers

2,563
citations

30
h-index

45
g-index

98
ext. papers

2,781
ext. citations

4.1
avg, IF

4.58
L-index

#	Paper	IF	Citations
97	Enduring effects of muscarinic receptor activation on adult hippocampal neurogenesis, microRNA expression and behaviour. <i>Behavioural Brain Research</i> , 2019 , 362, 188-198	3.4	2
96	Hippocampus-dependent fear conditioning is not sensitized by muscarinic receptor activation following systemic injection of pilocarpine. <i>Neurology Psychiatry and Brain Research</i> , 2019 , 34, 44-49	2.1	
95	Role of dorsal hippocampus μ opioid receptors in contextual aversive memory consolidation in rats. <i>Neuropharmacology</i> , 2018 , 135, 253-267	5.5	8
94	The sedative activity of flavonoids from <i>Passiflora quadrangularis</i> is mediated through the GABAergic pathway. <i>Biomedicine and Pharmacotherapy</i> , 2018 , 100, 388-393	7.5	14
93	Distinctive stress sensitivity and anxiety-like behavior in female mice: Strain differences matter. <i>Neurobiology of Stress</i> , 2018 , 9, 55-63	7.6	9
92	The role of Neuropeptide Y in fear conditioning and extinction. <i>Neuropeptides</i> , 2016 , 55, 111-26	3.3	66
91	Evidence for involvement of NK β receptors in the anxiogenic-like effect of SP6-11(C-terminal), a metabolite of substance P, in rats evaluated in the elevated plus-maze. <i>Behavioural Brain Research</i> , 2016 , 303, 168-75	3.4	6
90	Short-term enriched environment exposure facilitates fear extinction in adult rats: The NPY-Y1 receptor modulation. <i>Neuropeptides</i> , 2016 , 55, 73-8	3.3	10
89	The Role of Hippocampal NMDA Receptors in Long-Term Emotional Responses following Muscarinic Receptor Activation. <i>PLoS ONE</i> , 2016 , 11, e0147293	3.7	9
88	Gene and stress history interplay in emergence of PTSD-like features. <i>Behavioural Brain Research</i> , 2015 , 292, 266-77	3.4	8
87	Involvement of GABAergic pathway in the sedative activity of apigenin, the main flavonoid from <i>Passiflora quadrangularis</i> pericarp. <i>Revista Brasileira De Farmacognosia</i> , 2015 , 25, 158-163	2	23
86	Statins enhance cognitive performance in object location test in albino Swiss mice: involvement of beta-adrenoceptors. <i>Physiology and Behavior</i> , 2015 , 143, 27-34	3.5	6
85	Involvement of monoaminergic systems in the antidepressant-like properties of <i>Lafoensia pacari</i> A. St. Hil. <i>Journal of Ethnopharmacology</i> , 2015 , 170, 218-25	5	5
84	Involvement of the monoamine system in antidepressant-like properties of 4-(1-phenyl-1h-pyrazol-4-ylmethyl)-piperazine-1-carboxylic acid ethyl ester. <i>Life Sciences</i> , 2015 , 143, 187-93	6.8	9
83	Anxiolytic-like, stimulant and neuroprotective effects of <i>Ilex paraguariensis</i> extracts in mice. <i>Neuroscience</i> , 2015 , 292, 13-21	3.9	22
82	Nociceptin/orphanin FQ induces simultaneously anxiolytic and amnesic effects in the mouse elevated T-maze task. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2015 , 388, 33-41	3.4	7
81	Assessment of sedative effects of <i>Passiflora edulis</i> f. <i>flavicarpa</i> and <i>Passiflora alata</i> extracts in mice, measured by telemetry. <i>Phytotherapy Research</i> , 2014 , 28, 706-13	6.7	20

80	NK1 receptors antagonism of dorsal hippocampus counteract the anxiogenic-like effects induced by pilocarpine in non-convulsive Wistar rats. <i>Behavioural Brain Research</i> , 2014 , 265, 53-60	3.4	8
79	Anxiogenic-like profile of Wistar adult rats based on the pilocarpine model: an animal model for trait anxiety?. <i>Psychopharmacology</i> , 2013 , 227, 209-19	4.7	15
78	NMDA preconditioning attenuates cortical and hippocampal seizures induced by intracerebroventricular quinolinic acid infusion. <i>Neurotoxicity Research</i> , 2013 , 24, 55-62	4.3	8
77	GABA-A receptor modulators alter emotionality and hippocampal theta rhythm in an animal model of long-lasting anxiety. <i>Brain Research</i> , 2013 , 1532, 21-31	3.7	13
76	Neuropeptide Y (NPY) prevents depressive-like behavior, spatial memory deficits and oxidative stress following amyloid- β (1-40) administration in mice. <i>Behavioural Brain Research</i> , 2013 , 244, 107-15	3.4	62
75	Role of NPY Y1 receptor on acquisition, consolidation and extinction on contextual fear conditioning: dissociation between anxiety, locomotion and non-emotional memory behavior. <i>Neurobiology of Learning and Memory</i> , 2013 , 103, 26-33	3.1	50
74	A proposal for refining the forced swim test in Swiss mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013 , 45, 150-5	5.5	43
73	Evaluation of analgesic and anti-inflammatory activities of <i>Hydrocotyle umbellata</i> L., Araliaceae (acari β ba) in mice. <i>Anais Da Academia Brasileira De Ciencias</i> , 2013 , 85, 987-97	1.4	14
72	Phytochemical profile, toxicity and antioxidant activity of <i>Aloysia gratissima</i> (Verbenaceae). <i>Quimica Nova</i> , 2013 , 36, 69-73	1.6	15
71	Ghrelin as a neuroprotective and palliative agent in Alzheimer β and Parkinson β disease. <i>Current Pharmaceutical Design</i> , 2013 , 19, 6773-90	3.3	39
70	The anxiolytic-like effect of an essential oil derived from <i>Spiranthera odoratissima</i> A. St. Hil. leaves and its major component, E Caryophyllene, in male mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012 , 38, 276-84	5.5	77
69	Central pharmacological activity of a new piperazine derivative: 4-(1-phenyl-1h-pyrazol-4-ylmethyl)-piperazine-1-carboxylic acid ethyl ester. <i>Life Sciences</i> , 2012 , 90, 910-6	6.8	13
68	Murine model of repeated exposures to conspecific trained aggressors simulates features of post-traumatic stress disorder. <i>Behavioural Brain Research</i> , 2012 , 235, 55-66	3.4	42
67	Quantitative changes of nicotinic receptors in the hippocampus of dystrophin-deficient mice. <i>Brain Research</i> , 2012 , 1483, 96-104	3.7	14
66	Anti-inflammatory and opioid-like activities in methanol extract of <i>Mikania lindleyana</i> , sucuri β ju. <i>Revista Brasileira De Farmacognosia</i> , 2012 , 22, 150-156	2	2
65	Involvement of GABAergic non-benzodiazepine sites in the anxiolytic-like and sedative effects of the flavonoid baicalein in mice. <i>Behavioural Brain Research</i> , 2011 , 221, 75-82	3.4	70
64	Anxiolytic-like effect of central administration of NOP receptor antagonist UFP-101 in rats submitted to the elevated T-maze. <i>Behavioural Brain Research</i> , 2011 , 222, 206-11	3.4	23
63	Effects of social isolation and enriched environment on behavior of adult Swiss mice do not require hippocampal neurogenesis. <i>Behavioural Brain Research</i> , 2011 , 225, 85-90	3.4	15

62	Role of ventral hippocampal nitric oxide/cGMP pathway in anxiety-related behaviors in rats submitted to the elevated T-maze. <i>Behavioural Brain Research</i> , 2010 , 207, 112-7	3.4	27
61	Short- and long-term anxiogenic effects induced by a single injection of subconvulsant doses of pilocarpine in rats: investigation of the putative role of hippocampal pathways. <i>Psychopharmacology</i> , 2010 , 212, 653-61	4.7	15
60	Allosteric interaction of the anticholinergic drug [N-(4-phenyl)-phenacyl-l-hyoscyamine] (Phenthonium) with nicotinic receptors of post-ganglionic sympathetic neurons of the rat vas deferens. <i>European Journal of Pharmacology</i> , 2009 , 616, 229-35	5.3	1
59	Neuropharmacological activity of the pericarp of <i>Passiflora edulis flavicarpa</i> degener: putative involvement of C-glycosylflavonoids. <i>Experimental Biology and Medicine</i> , 2009 , 234, 967-75	3.7	49
58	The antidepressant-like effects of <i>Aloysia polystachya</i> (Griseb.) Moldenke (Verbenaceae) in mice. <i>Phytomedicine</i> , 2008 , 15, 478-83	6.5	19
57	Frequency of climbing behavior as a predictor of altered motor activity in rat forced swimming test. <i>Neuroscience Letters</i> , 2008 , 445, 170-3	3.3	28
56	Systemic administration of a nitric oxide synthase inhibitor impairs fear sensitization in the plus-maze. <i>Neurobiology of Learning and Memory</i> , 2008 , 90, 455-9	3.1	2
55	GABA(A) signalling is involved in N/OFQ anxiolytic-like effects but not in nocistatin anxiogenic-like action as evaluated in the mouse elevated plus maze. <i>Peptides</i> , 2008 , 29, 1404-12	3.8	14
54	Evidence for the involvement of the monoaminergic system in the antidepressant-like action of two 4-amine derivatives of 10,11-dihydro-5H-dibenzo [a,d] cycloheptane in mice evaluated in the tail suspension test. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008 , 32, 368-74	5.5	10
53	Modulation of anxiety in rats evaluated in the elevated T-maze: evidence of the relationship between substance P and diazepam. <i>Behavioural Brain Research</i> , 2008 , 187, 140-5	3.4	9
52	The microinjection of AMPA receptor antagonist into the accumbens shell, but not into the accumbens core, induces anxiolysis in an animal model of anxiety. <i>Behavioural Brain Research</i> , 2008 , 188, 91-9	3.4	26
51	Assessment of luteolin (3R,4R,5,7-tetrahydroxyflavone) neuropharmacological activity. <i>Behavioural Brain Research</i> , 2008 , 189, 75-82	3.4	75
50	Nitric oxide involvement and neural substrates of the conditioned and innate fear as evaluated in the T-maze test in rats. <i>Behavioural Brain Research</i> , 2008 , 189, 341-9	3.4	12
49	Anticonvulsant and anxiolytic-like effects of compounds isolated from <i>Polygala sabulosa</i> (Polygalaceae) in rodents: in vitro and in vivo interactions with benzodiazepine binding sites. <i>Psychopharmacology</i> , 2008 , 197, 351-60	4.7	26
48	Antidepressant-like profile of action of two 4-amine derivatives of 10,11-dihydro-5H-dibenzo [a,d] cycloheptane in mice evaluated in the forced swimming test. <i>Bioorganic and Medicinal Chemistry</i> , 2007 , 15, 1645-50	3.4	5
47	Participation of dihydrostyryl-2-pyrones and styryl-2-pyrones in the central effects of <i>Polygala sabulosa</i> (Polygalaceae), a folk medicine topical anesthetic. <i>Pharmacology Biochemistry and Behavior</i> , 2007 , 86, 150-61	3.9	24
46	Chemical standardization of the aqueous extract of <i>Cecropia glaziovii</i> Sneth endowed with antihypertensive, bronchodilator, antiacid secretion and antidepressant-like activities. <i>Phytomedicine</i> , 2007 , 14, 309-13	6.5	38
45	Antihypertensive effect of a standardized aqueous extract of <i>Cecropia glaziovii</i> Sneth in rats: an in vivo approach to the hypotensive mechanism. <i>Phytomedicine</i> , 2007 , 14, 314-20	6.5	37

44	Antidepressant-like effect of <i>Cecropia glazioui</i> Sneth and its constituents - in vivo and in vitro characterization of the underlying mechanism. <i>Phytomedicine</i> , 2007 , 14, 396-402	6.5	45
43	An approach to evaluate the ability of rats to discriminate different levels of illumination in the plus maze test: effects of scopolamine. <i>Behavioural Brain Research</i> , 2007 , 180, 86-94	3.4	9
42	Involvement of NK1 receptors in metabolic stress markers after the central administration of substance P. <i>Behavioural Brain Research</i> , 2007 , 181, 232-8	3.4	14
41	GABAA and GABAB agonist microinjections into medial accumbens shell increase feeding and induce anxiolysis in an animal model of anxiety. <i>Behavioural Brain Research</i> , 2007 , 184, 142-9	3.4	28
40	Behavioral effects of a neurotoxic compound isolated from <i>Clibadium surinamense</i> L (Asteraceae). <i>Neurotoxicology and Teratology</i> , 2006 , 28, 349-53	3.9	5
39	Anxiolytic and antidepressant-like activity of a standardized extract from <i>Galphimia glauca</i> . <i>Phytomedicine</i> , 2006 , 13, 23-8	6.5	45
38	P.1.d.017 Participation of the ventral hippocampus in the modulation of anxiety: role of nitric oxide, Rizzi A., et al., in the 2002, Br. behaviour of J. Pharmacol, 136, rats evaluated in the 303-311. elevated T-maze. <i>European Neuropsychopharmacology</i> , 2006 , 16, S258-S259	1.2	
37	Antidepressant treatment reduces Fos-like immunoreactivity induced by swim stress in different columns of the periaqueductal gray matter. <i>Brain Research Bulletin</i> , 2006 , 70, 414-21	3.9	22
36	The anxiolytic-like effects of <i>Aloysia polystachya</i> (Griseb.) Moldenke (Verbenaceae) in mice. <i>Journal of Ethnopharmacology</i> , 2006 , 105, 400-8	5	45
35	Neuropharmacological evaluation of the putative anxiolytic effects of <i>Passiflora edulis</i> Sims, its sub-fractions and flavonoid constituents. <i>Phytotherapy Research</i> , 2006 , 20, 1067-73	6.7	70
34	Synthesis and antidepressant-like action of stereoisomers of imidobenzenesulfonylaziridines in mice evaluated in the forced swimming test. <i>Bioorganic and Medicinal Chemistry</i> , 2006 , 14, 5397-401	3.4	13
33	Central nervous system activity of the hydroalcoholic extract of <i>Casimiroa edulis</i> in rats and mice. <i>Journal of Ethnopharmacology</i> , 2005 , 97, 191-7	5	33
32	Structure of the rat behaviour in the forced swimming test. <i>Behavioural Brain Research</i> , 2005 , 158, 243-50	9.4	66
31	Behavioral and immunological effects of substance P in female and male mice. <i>Pharmacology Biochemistry and Behavior</i> , 2004 , 79, 1-9	3.9	10
30	Antidepressant-like effects of the nociceptin/orphanin FQ receptor antagonist UFP-101: new evidence from rats and mice. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2004 , 369, 547-53	3.4	84
29	Further evidence on the anxiogenic-like effect of substance P evaluated in the elevated plus-maze in rats. <i>Behavioural Brain Research</i> , 2004 , 154, 501-10	3.4	23
28	Involvement of tachykinin NK1 receptor in the behavioral and immunological responses to swimming stress in mice. <i>Neuropeptides</i> , 2003 , 37, 307-15	3.3	12
27	Antidepressant-like effect of Ro5-4864, a peripheral-type benzodiazepine receptor ligand, in forced swimming test. <i>European Journal of Pharmacology</i> , 2003 , 471, 21-6	5.3	24

26	Blockade of nociceptin/orphanin FQ-NOP receptor signalling produces antidepressant-like effects: pharmacological and genetic evidences from the mouse forced swimming test. <i>European Journal of Neuroscience</i> , 2003 , 17, 1987-90	3.5	103
25	Evaluation of the anxiolytic-like effects of <i>Cecropia glazioui</i> Sneth in mice. <i>Pharmacology Biochemistry and Behavior</i> , 2002 , 71, 183-90	3.9	37
24	Central injections of nocistatin or its C-terminal hexapeptide exert anxiogenic-like effect on behaviour of mice in the plus-maze test. <i>British Journal of Pharmacology</i> , 2002 , 136, 764-72	8.6	54
23	Dorsal periaqueductal gray matter inhibits passive coping strategy elicited by forced swimming stress in rats. <i>Neuroscience Letters</i> , 2002 , 335, 87-90	3.3	16
22	The role of lateral septal NK1 receptors in mediating anxiogenic effects induced by intracerebroventricular injection of substance P. <i>Behavioural Brain Research</i> , 2002 , 134, 411-5	3.4	29
21	CNS activities of liquid and spray-dried extracts from <i>Lippia alba</i> -Verbenaceae (Brazilian false melissa). <i>Journal of Ethnopharmacology</i> , 2002 , 82, 207-15	5	40
20	Participation of GABAA receptors in the modulation of experimental anxiety by tachykinin agonists and antagonists in mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2002 , 26, 861-9	5.5	15
19	Nitric oxide involvement in the anxiogenic-like effect of substance P. <i>Behavioural Brain Research</i> , 2001 , 121, 199-205	3.4	15
18	Tachykinin NK(3)receptor involvement in anxiety. <i>Neuropeptides</i> , 1999 , 33, 181-8	3.3	52
17	Behavioral effects of essential oils from <i>Lippia alba</i> (Mill.) N.E. Brown chemotypes. <i>Journal of Ethnopharmacology</i> , 1999 , 67, 127-33	5	54
16	Anxiogenic-like effect induced by substance P injected into the lateral septal nucleus. <i>NeuroReport</i> , 1999 , 10, 3399-403	1.7	40
15	Naloxone-induced changes in tachykinin NK3 receptor modulation of experimental anxiety in mice. <i>Neuroscience Letters</i> , 1998 , 258, 155-8	3.3	33
14	Opposing roles of the amygdala and dorsolateral periaqueductal gray in fear-potentiated startle. <i>Neuroscience and Biobehavioral Reviews</i> , 1997 , 21, 743-53	9	84
13	Effects of central administration of tachykinin receptor agonists and antagonists on plus-maze behavior in mice. <i>European Journal of Pharmacology</i> , 1996 , 311, 7-14	5.3	133
12	Antinociceptive effects of clebopride in the mouse. <i>General Pharmacology</i> , 1995 , 26, 1083-7		2
11	Involvement of cyclic AMP at the level of the nucleus reticularis pontis caudalis in the acoustic startle response. <i>Brain Research</i> , 1995 , 700, 59-69	3.7	12
10	Evaluation of the central properties of <i>Artemisia verlotorum</i> . <i>Planta Medica</i> , 1993 , 59, 326-9	3.1	17
9	Effects of cold-restraint and swim stress on convulsions induced by pentylenetetrazol and electroshock: influence of naloxone pretreatment. <i>Pharmacology Biochemistry and Behavior</i> , 1991 , 40, 297-300	3.9	51

8	Chemical and pharmacological analysis of the crude aqueous/alcoholic extract from <i>Cordyline dracaenoides</i> . <i>Phytotherapy Research</i> , 1990 , 4, 167-171	6.7	6
7	The influence of stress on convulsive parameters in the mouse. <i>Neuroscience and Biobehavioral Reviews</i> , 1990 , 14, 491-4	9	16
6	Influence of long-term treatment of the rat with clebopride on the morphology of the mammary gland. <i>Pharmacology</i> , 1990 , 40, 54-9	2.3	1
5	Chemical and pharmacological studies on <i>Talauma ovata</i> St. Hil. (Magnoliaceae). <i>Journal of Ethnopharmacology</i> , 1989 , 26, 277-86	5	10
4	Pharmacological actions of tannic acid; II. Evaluation of CNS activity in animals. <i>Planta Medica</i> , 1986 , 52, 272-5	3.1	18
3	Effects of withdrawal from long-term barbital treatment on open-field behaviour and seizure susceptibility of rats. <i>Neuropharmacology</i> , 1982 , 21, 277-81	5.5	3
2	Effects of withdrawal from long-term diphenylhydantoin treatment on audiogenic and maximal electroshock-induced seizures in rats. <i>Acta Neurologica Scandinavica</i> , 1981 , 63, 189-96	3.8	2
1	The effects of diphenylhydantoin on rat behavior. <i>Psychopharmacology</i> , 1980 , 69, 183-5	4.7	8