

# Abdelhamid Khaldi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3955887/publications.pdf>

Version: 2024-02-01

63  
papers

1,445  
citations

331670

21  
h-index

361022

35  
g-index

65  
all docs

65  
docs citations

65  
times ranked

1833  
citing authors

#	ARTICLE	IF	CITATIONS
1	Metabolite profiling and potential antioxidant activity of sixteen fennel ( <i>Foeniculum vulgare</i> Mill.) populations growing wild in Tunisia. <i>South African Journal of Botany</i> , 2022, 148, 407-414.	2.5	20
2	Effect of Growing Area on Total Polyphenols, Flavonoids, Tannins and Antimicrobial Activity in <i>Quercus suber</i> L. Acorn Oil. <i>Journal of Food Chemistry and Nanotechnology</i> , 2021, 7, 30-33.	0.3	1
3	Mineral Composition of Bluish-Black and Yellowish- White <i>Myrtus communis</i> L. Berries and <i>Arbutus unedo</i> L. Fruits. <i>Journal of Food Chemistry and Nanotechnology</i> , 2021, 7, 1-3.	0.3	0
4	Morphological and Chemical Differentiation between Tunisian Populations of <i>Pinus halepensis</i> , <i>Pinus brutia</i> , and <i>Pinus pinaster</i> . <i>Chemistry and Biodiversity</i> , 2021, 18, e2100071.	2.1	3
5	Chemotaxonomic Study of Four Subspecies of <i>Pinus nigra</i> Arn. Grown in Common Garden Based on Essential Oil Composition. <i>Journal of Food Quality</i> , 2021, 2021, 1-7.	2.6	1
6	Tree growth and leaf gas exchange variability of three Mediterranean <i>Pinus</i> spp. growing in a common garden in Northeastern Tunisia. <i>Euro-Mediterranean Journal for Environmental Integration</i> , 2020, 5, 1.	1.3	2
7	Variation of Essential Oil Composition, Antioxidant and Anticholinesterase Activities between <i>Pinus halepensis</i> Mill. Plant Organs. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2020, 23, 1450-1462.	1.9	6
8	Relationship between climate and growth of two North African varieties of <i>Pinus pinaster</i> Arn.. <i>African Journal of Ecology</i> , 2019, 57, 327-334.	0.9	3
9	Protective effects of phytochemicals of <i>Capparis spinosa</i> seeds with cisplatin and CCl <sub>4</sub> toxicity in mice. <i>Food Bioscience</i> , 2019, 28, 42-48.	4.4	25
10	Fatty acids and triacylglycerols composition from Tunisian <i>Acacia</i> species seed oil. <i>Arabian Journal of Chemistry</i> , 2019, 12, 3302-3308.	4.9	15
11	Chemical composition and biological activities essential oil from the needles African of <i>Pinus pinaster</i> Var.. <i>Revue Roumaine De Chimie</i> , 2019, 64, 511-518.	0.2	7
12	Variation in essential oil composition and biological activities of <i>Foeniculum vulgare</i> Mill. populations growing widely in Tunisia. <i>Journal of Food Biochemistry</i> , 2018, 42, e12532.	2.9	32
13	Phenolic profile and effect of growing area on <i>Pistacia lentiscus</i> seed oil. <i>Food Chemistry</i> , 2018, 257, 206-210.	8.2	14
14	Chemical composition and antioxidant activity of the volatile fraction extracted from air-dried fruits of Tunisian <i>Eryngium maritimum</i> L. ecotypes. <i>Journal of the Science of Food and Agriculture</i> , 2018, 98, 635-643.	3.5	16
15	Tree-rings to climate relationships in nineteen provenances of four black pines sub-species ( <i>Pinus</i> ) Tj ETQq1 1 0.784314 rgBT /Overlo	2.2	11
16	<i>Schinus terebinthifolius</i> vs <i>Schinus molle</i> : A comparative study of the effect of species and location on the phytochemical content of fruits. <i>Industrial Crops and Products</i> , 2018, 122, 559-565.	5.2	28
17	Lipid characterization of <i>Eryngium maritimum</i> seeds grown in Tunisia. <i>Industrial Crops and Products</i> , 2017, 105, 47-52.	5.2	14
18	Carbon stocks distribution in shrub species of a North African cork oak forest. <i>African Journal of Ecology</i> , 2017, 55, 693-696.	0.9	0

#	ARTICLE	IF	CITATIONS
19	Capparis spinosa leaves extract: Source of bioantioxidants with nephroprotective and hepatoprotective effects. Biomedicine and Pharmacotherapy, 2017, 87, 171-179.	5.6	61
20	Dendroecological study of Pinus halepensis and Pinus pinea in northeast coastal dunes in Tunisia according to distance from the shoreline and dieback intensity. Dendrochronologia, 2017, 45, 62-72.	2.2	20
21	Adjustment of photosynthetic carbon assimilation to higher growth irradiance in three-year-old seedlings of two Tunisian provenances of Cork Oak (Quercus suber L.). IForest, 2017, 10, 618-624.	1.4	7
22	Diversity of Sterol Composition in Tunisian <i>Pistacia lentiscus</i> Seed Oil. Chemistry and Biodiversity, 2016, 13, 544-548.	2.1	8
23	Plant diversity in different bioclimatic zones in Tunisia. Journal of Asia-Pacific Biodiversity, 2016, 9, 56-62.	0.4	40
24	Effects of Rhus tripartitum fruit extract on CCl4-induced hepatotoxicity and cisplatin-induced nephrotoxicity in rats. Canadian Journal of Physiology and Pharmacology, 2016, 94, 801-807.	1.4	17
25	Evolution-based approach needed for the conservation and silviculture of peripheral forest tree populations. Forest Ecology and Management, 2016, 375, 66-75.	3.2	97
26	Evaluation of <i>Pistacia lentiscus</i> seed oil and phenolic compounds for <i>in vitro</i> antiproliferative effects against BHK21 cells. Pharmaceutical Biology, 2016, 54, 747-751.	2.9	18
27	Estimate of biomass and carbon pools in disturbed and undisturbed oak forests in Tunisia. Forest Systems, 2016, 25, e060.	0.3	11
28	Unexploited <i>Thapsia garganica</i> , <i>Orlaya maritima</i> , and <i>Retama raetam</i> Seeds: Potential Sources of Unsaturated Fatty Acid and Natural Antioxidants. JAOCS, Journal of the American Oil Chemists' Society, 2015, 92, 1175-1181.	1.9	3
29	Phenolic profile and antioxidant activity of Capparis spinosa seeds harvested from different wild habitats. Industrial Crops and Products, 2015, 76, 930-935.	5.2	54
30	<i>In vitro</i> antimicrobial activity of <i>Pistacia lentiscus</i> L. edible oil and phenolic extract. Natural Product Research, 2015, 29, 565-570.	1.8	23
31	IMPACTS OF LOCATION AND FORESTRY CONDITIONS ON SOME PHYSICAL AND MECHANICAL PROPERTIES OF NORTHERN TUNISIAN PINUS PINEA L. WOOD.. Bois Et Forets Des Tropiques, 2015, 324, 65.	0.2	5
32	Transcriptome profiling the basal region of poplar stems during the early gravitropic response. Biologia Plantarum, 2014, 58, 55-63.	1.9	13
33	Effect of growing area on tocopherols, carotenoids and fatty acid composition of <i>Pistacia lentiscus</i> edible oil. Natural Product Research, 2014, 28, 1225-1230.	1.8	21
34	Phytochemicals and antioxidant activities of Rhus tripartitum (Ucra) fruits depending on locality and different stages of maturity. Food Chemistry, 2014, 160, 98-103.	8.2	64
35	Variation in protein and oil content and fatty acid composition of Rhus tripartitum fruits collected at different maturity stages in different locations. Industrial Crops and Products, 2014, 59, 197-201.	5.2	23
36	Contents of Carotenoids, Tocopherols and Sterols in <i>Acacia cyanophylla</i> Seed Oils. JAOCS, Journal of the American Oil Chemists' Society, 2013, 90, 429-436.	1.9	9

#	ARTICLE	IF	CITATIONS
37	Screening of Natural Antioxidants from Selected Medicinal Plants. International Journal of Food Properties, 2013, 16, 1117-1126.	3.0	61
38	Evolution of growth-climate relationships of three pine species in Kroumirie (North-West Tunisia). SÅcheresse, 2013, 24, 138-146.	0.1	3
39	Minor lipid components of some Acacia species: potential dietary health benefits of the unexploited seeds. Lipids in Health and Disease, 2012, 11, 49.	3.0	12
40	Unexploited Acacia cyanophylla seeds: potential food sources of %6 fatty acids and antioxidants?. Journal of the Science of Food and Agriculture, 2012, 92, 1526-1532.	3.5	12
41	Assessment of land-cover change using GIS and remotely-sensed data: A case study in Ain Snoussi area of northern Tunisia. Forest Science and Technology, 2011, 7, 75-81.	0.8	7
42	Volatile Constituents of Pinus pinea L. Needles. Journal of Essential Oil Research, 2011, 23, 15-19.	2.7	11
43	Chemical compounds from Phoenician juniper berries (<i>Juniperus phoenicea</i>). Natural Product Research, 2011, 25, 1733-1742.	1.8	27
44	Chemical Polymorphism of Essential Oils from Populations of <i>Laurus nobilis</i> Grown on Tunisia, Algeria and France. Natural Product Communications, 2011, 6, 1934578X1100601.	0.5	9
45	PHENOLIC COMPOUNDS, TOCOPHEROLS, CAROTENOIDS AND VITAMIN C OF COMMERCIAL CAPER. Journal of Food Biochemistry, 2011, 35, 472-483.	2.9	28
46	The caper (Capparis L.): Ethnopharmacology, phytochemical and pharmacological properties. FÅ-toterapÅ-Å¢, 2011, 82, 93-101.	2.2	116
47	Salinity tolerance of hydroponically grown Pinus pinea L. seedlings. Acta Physiologiae Plantarum, 2011, 33, 765-775.	2.1	9
48	Protein, Lipid, Aliphatic and Triterpenic Alcohol Content of Caper Seeds â€œ<i>Capparis spinosa</i>â€œ. JAOCS, Journal of the American Oil Chemists' Society, 2011, 88, 265-270.	1.9	29
49	Triacylglycerols and Phospholipids Composition of Caper Seeds (Capparis spinosa). JAOCS, Journal of the American Oil Chemists' Society, 2011, 88, 1787-1793.	1.9	8
50	Phenolic Compounds and Vitamin Antioxidants of Caper (Capparis spinosa). Plant Foods for Human Nutrition, 2010, 65, 260-265.	3.2	97
51	Essential Oils of Daucus carota subsp. carota of Tunisia Obtained by Supercritical Carbon Dioxide Extraction. Natural Product Communications, 2010, 5, 1934578X1000501.	0.5	9
52	Fatty Acid Composition, Essential Oil and Antibacterial Activity of Berries of Laurus nobilisL.. Journal of Essential Oil-bearing Plants: JEOP, 2009, 12, 422-434.	1.9	3
53	Biological activity evaluation of the oils from <i>Laurus nobilis</i> of Tunisia and Algeria extracted by supercritical carbon dioxide. Natural Product Research, 2009, 23, 230-237.	1.8	28
54	High tocopherol and triacylglycerol contents in <i>Pinus</i> <i>pinea</i> L. seeds. International Journal of Food Sciences and Nutrition, 2009, 60, 161-169.	2.8	12

#	ARTICLE	IF	CITATIONS
55	FATTY ACIDS, TOCOPHEROLS AND CAROTENOIDS FROM SEEDS OF TUNISIAN CAPER <i>CAPPARIS SPINOSA</i> . Journal of Food Lipids, 2009, 16, 452-464.	1.0	34
56	Intraspecific Variation of <i>Capparis spinosa</i> L. in Tunisia. Journal of Herbs, Spices and Medicinal Plants, 2009, 15, 9-15.	1.1	12
57	Carotenoid and Tocopherol Composition of Leaves, Buds, and Flowers of <i>Capparis spinosa</i> Grown Wild in Tunisia. Journal of Agricultural and Food Chemistry, 2009, 57, 5381-5385.	5.2	45
58	Population Genetic Structure of <i>Laurus nobilis</i> L. Inferred From Transferred Nuclear Microsatellites. <i>Silvae Genetica</i> , 2009, 58, 270-276.	0.8	5
59	Fatty Acid Composition of Two Tunisian Pine Seed Oils. <i>Biotechnology Progress</i> , 2008, 21, 998-1001.	2.6	20
60	The analysis of crude and purified locust bean gum: A comparison of samples from different carob tree populations in Tunisia. <i>Food Chemistry</i> , 2007, 101, 1508-1515.	8.2	69
61	Fatty acids from seeds of <i>Pinus pinea</i> L.: Composition and population profiling. <i>Phytochemistry</i> , 2005, 66, 1729-1735.	2.9	80
62	Breeding Improvement of <i>Laurus nobilis</i> L. by Conventional and <i>In Vitro</i> Propagation Techniques. <i>Journal of Herbs, Spices and Medicinal Plants</i> , 2002, 9, 101-105.	1.1	5
63	Towards optimizing acorn use as animal feed in Tunisia: evaluation and impact on natural regeneration. <i>Bois Et Forets Des Tropiques</i> , 0, 348, 17-26.	0.2	2