

Daniel T Monaghan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3957038/publications.pdf>

Version: 2024-02-01

41
papers

2,318
citations

236833

25
h-index

315616

38
g-index

42
all docs

42
docs citations

42
times ranked

2024
citing authors

#	ARTICLE	IF	CITATIONS
1	The continually evolving role of NMDA receptors in neurobiology and disease. <i>Neuropharmacology</i> , 2022, 210, 109042.	2.0	3
2	Pharmacological characterization of a novel negative allosteric modulator of NMDA receptors, UBP792. <i>Neuropharmacology</i> , 2021, 201, 108818.	2.0	0
3	NMDA receptors containing GluN2C and GluN2D subunits have opposing roles in modulating neuronal oscillations; potential mechanism for bidirectional feedback. <i>Brain Research</i> , 2020, 1727, 146571.	1.1	12
4	Structural basis of subtype-selective competitive antagonism for GluN2C/2D-containing NMDA receptors. <i>Nature Communications</i> , 2020, 11, 423.	5.8	19
5	Investigation of the structural requirements for N-methyl-D-aspartate receptor positive and negative allosteric modulators based on 2-naphthoic acid. <i>European Journal of Medicinal Chemistry</i> , 2019, 164, 471-498.	2.6	10
6	In the Telencephalon, GluN2C NMDA Receptor Subunit mRNA is Predominately Expressed in Glial Cells and GluN2D mRNA in Interneurons. <i>Neurochemical Research</i> , 2019, 44, 61-77.	1.6	33
7	The NMDA receptor intracellular C-terminal domains reciprocally interact with allosteric modulators. <i>Biochemical Pharmacology</i> , 2019, 159, 140-153.	2.0	13
8	Positive and Negative Allosteric Modulators of <i>N</i> -Methyl-D-aspartate (NMDA) Receptors: Structure-Activity Relationships and Mechanisms of Action. <i>Journal of Medicinal Chemistry</i> , 2019, 62, 3-23.	2.9	44
9	Mechanism and properties of positive allosteric modulation of N-methyl-d-aspartate receptors by 6-alkyl 2-naphthoic acid derivatives. <i>Neuropharmacology</i> , 2017, 125, 64-79.	2.0	15
10	A single-channel mechanism for pharmacological potentiation of GluN1/GluN2A NMDA receptors. <i>Scientific Reports</i> , 2017, 7, 6933.	1.6	7
11	Multiple roles of GluN2B-containing NMDA receptors in synaptic plasticity in juvenile hippocampus. <i>Neuropharmacology</i> , 2017, 112, 76-83.	2.0	33
12	The NMDA receptor GluN2C subunit controls cortical excitatory-inhibitory balance, neuronal oscillations and cognitive function. <i>Scientific Reports</i> , 2016, 6, 38321.	1.6	50
13	GluN2D N-Methyl-D-Aspartate Receptor Subunit Contribution to the Stimulation of Brain Activity and Gamma Oscillations by Ketamine: Implications for Schizophrenia. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2016, 356, 702-711.	1.3	56
14	Bidirectional Effect of Pregnenolone Sulfate on GluN1/GluN2A <i>N</i> -Methyl-d-Aspartate Receptor Gating Depending on Extracellular Calcium and Intracellular Milieu. <i>Molecular Pharmacology</i> , 2015, 88, 650-659.	1.0	11
15	Synthesis of a Series of Novel 3,9-Disubstituted Phenanthrenes as Analogues of Known N-Methyl-d-aspartate Receptor Allosteric Modulators. <i>Synthesis</i> , 2015, 47, 1593-1610.	1.2	9
16	Different NMDA receptor subtypes mediate induction of long-term potentiation and two forms of short-term potentiation at CA1 synapses in rat hippocampus <i>in vitro</i> . <i>Journal of Physiology</i> , 2013, 591, 955-972.	1.3	83
17	The NMDA receptor as a target for cognitive enhancement. <i>Neuropharmacology</i> , 2013, 64, 13-26.	2.0	206
18	Piperazine-2,3-dicarboxylic Acid Derivatives as Dual Antagonists of NMDA and GluK1-Containing Kainate Receptors. <i>Journal of Medicinal Chemistry</i> , 2012, 55, 327-341.	2.9	19

#	ARTICLE	IF	CITATIONS
19	Structure-activity relationships for allosteric NMDA receptor inhibitors based on 2-naphthoic acid. <i>Neuropharmacology</i> , 2012, 62, 1730-1736.	2.0	33
20	Coumarin-3-carboxylic acid derivatives as potentiators and inhibitors of recombinant and native N-methyl-d-aspartate receptors. <i>Neurochemistry International</i> , 2012, 61, 593-600.	1.9	37
21	Pharmacological modulation of NMDA receptor activity and the advent of negative and positive allosteric modulators. <i>Neurochemistry International</i> , 2012, 61, 581-592.	1.9	77
22	Selective inhibition of GluN2D-containing N-methyl-D-aspartate receptors prevents tissue plasminogen activator-promoted neurotoxicity both in vitro and in vivo. <i>Molecular Neurodegeneration</i> , 2011, 6, 68.	4.4	33
23	A Novel Family of Negative and Positive Allosteric Modulators of NMDA Receptors. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2010, 335, 614-621.	1.3	80
24	<i>N</i> -Methyl-d-aspartate (NMDA) Receptor NR2 Subunit Selectivity of a Series of Novel Piperazine-2,3-dicarboxylate Derivatives: Preferential Blockade of Extrasynaptic NMDA Receptors in the Rat Hippocampal CA3-CA1 Synapse. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2009, 331, 618-626.	1.3	46
25	Pharmacology of NMDA Receptors. <i>Frontiers in Neuroscience</i> , 2008, , 257-281.	0.0	3
26	CPP. , 2007, , 1-6.		0
27	Regional variations in NMDA receptor downregulation in streptozotocin-diabetic rat brain. <i>Brain Research</i> , 2006, 1115, 217-222.	1.1	9
28	Identification of Subunit- and Antagonist-Specific Amino Acid Residues in the N-Methyl-d-aspartate Receptor Glutamate-Binding Pocket. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005, 313, 1066-1074.	1.3	37
29	Synthesis and Pharmacology of <i>N</i> -1-Substituted Piperazine-2,3-dicarboxylic Acid Derivatives Acting as NMDA Receptor Antagonists. <i>Journal of Medicinal Chemistry</i> , 2005, 48, 2627-2637.	2.9	56
30	The effect of competitive antagonist chain length on NMDA receptor subunit selectivity. <i>Neuropharmacology</i> , 2005, 48, 354-359.	2.0	52
31	Structure-activity analysis of a novel NR2C/NR2D-preferring NMDA receptor antagonist: 1-(phenanthrene-2-carbonyl) piperazine-2,3-dicarboxylic acid. <i>British Journal of Pharmacology</i> , 2004, 141, 508-516.	2.7	122
32	Extrasynaptic NR2B and NR2D subunits of NMDA receptors shape "superslow" afterburst EPSC in rat hippocampus. <i>Journal of Physiology</i> , 2004, 558, 451-463.	1.3	142
33	Distinct NMDA Receptor Subpopulations Contribute to Long-Term Potentiation and Long-Term Depression Induction. <i>Journal of Neuroscience</i> , 2000, 20, RC81-RC81.	1.7	116
34	Organotypic Brain Slice Cultures for Functional Analysis of Alcohol-Related Disorders: Novel Versus Conventional Preparations. <i>Alcoholism: Clinical and Experimental Research</i> , 1998, 22, 51-59.	1.4	13
35	Chapter 12 Molecular determinants of NMDA receptor pharmacological diversity. <i>Progress in Brain Research</i> , 1998, 116, 171-190.	0.9	31
36	Dextromethorphan and Other N-Methyl-D-Aspartate Receptor Antagonists Are Teratogenic in the Avian Embryo Model. <i>Pediatric Research</i> , 1998, 43, 1-7.	1.1	65

#	ARTICLE	IF	CITATIONS
37	Pharmacological heterogeneity of NMDA receptors: characterization of NR1a/NR2D heteromers expressed in <i>Xenopus</i> oocytes. <i>European Journal of Pharmacology</i> , 1997, 320, 87-94.	1.7	71
38	Insulin potentiates N-methyl-d-aspartate receptor activity in <i>Xenopus</i> oocytes and rat hippocampus. <i>Neuroscience Letters</i> , 1995, 192, 5-8.	1.0	109
39	The distribution of [3H]kainate binding sites in primate hippocampus is similar to the distribution of both Ca ²⁺ -sensitive and Ca ²⁺ -insensitive [3H]kainate binding sites in rat hippocampus. <i>Neurochemical Research</i> , 1986, 11, 1073-1082.	1.6	31
40	Anatomical distributions of four pharmacologically distinct 3H-L-glutamate binding sites. <i>Nature</i> , 1983, 306, 176-179.	13.7	528
41	The Excitatory Amino Acid System. , 0, , 67-84.		2