

Armin W Lorenz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3957782/publications.pdf>

Version: 2024-02-01

55
papers

2,073
citations

159585

30
h-index

243625

44
g-index

55
all docs

55
docs citations

55
times ranked

1941
citing authors

#	ARTICLE	IF	CITATIONS
1	Decline in niche specialization and trait $\hat{\rho}^2$ -diversity in benthic invertebrate communities of Central European low-mountain streams over 25 years. <i>Science of the Total Environment</i> , 2022, 810, 151770.	8.0	3
2	Generalist parasites persist in degraded environments: a lesson learned from microsporidian diversity in amphipods. <i>Parasitology</i> , 2022, 149, 973-982.	1.5	7
3	Woody buffer effects on water temperature: The role of spatial configuration and daily temperature fluctuations. <i>Hydrological Processes</i> , 2021, 35, e14008.	2.6	13
4	Continuous riverine biodiversity changes in a 10 years post-restoration study: Impacts and pitfalls. <i>River Research and Applications</i> , 2021, 37, 270-282.	1.7	12
5	The interplay of nutrients, dissolved inorganic carbon and algae in determining macrophyte occurrences in rivers. <i>Science of the Total Environment</i> , 2021, 781, 146728.	8.0	13
6	Active reintroduction of benthic invertebrates to increase stream biodiversity. <i>Limnologica</i> , 2020, 80, 125726.	1.5	7
7	The effect of lateral connectedness on the taxonomic and functional structure of fish communities in a lowland river floodplain. <i>Science of the Total Environment</i> , 2020, 719, 137169.	8.0	7
8	Diverging response patterns of terrestrial and aquatic species to hydromorphological restoration. <i>Conservation Biology</i> , 2019, 33, 132-141.	4.7	18
9	Reintroduction of freshwater macroinvertebrates: challenges and opportunities. <i>Biological Reviews</i> , 2019, 94, 368-387.	10.4	43
10	Effect of river restoration on life-history strategies in fish communities. <i>Science of the Total Environment</i> , 2019, 663, 486-495.	8.0	14
11	Development and validation of a macroinvertebrate-based biomonitoring tool to assess fine sediment impact in small mountain streams. <i>Science of the Total Environment</i> , 2019, 652, 1290-1301.	8.0	43
12	Moderate warming over the past 25 years has already reorganized stream invertebrate communities. <i>Science of the Total Environment</i> , 2019, 658, 1531-1538.	8.0	53
13	Revisiting restored river reaches – Assessing change of aquatic and riparian communities after five years. <i>Science of the Total Environment</i> , 2018, 613-614, 1185-1195.	8.0	45
14	How to facilitate freshwater macroinvertebrate reintroduction?. <i>Limnologica</i> , 2018, 69, 24-27.	1.5	7
15	Riparian plant species preferences indicate diversification of site conditions after river restoration. <i>Ecohydrology</i> , 2017, 10, e1852.	2.4	14
16	Mechanistic modelling for predicting the effects of restoration, invasion and pollution on benthic macroinvertebrate communities in rivers. <i>Freshwater Biology</i> , 2017, 62, 1083-1093.	2.4	5
17	Insights into species diversity of the genus <i>Hydropsyche</i> Pictet, 1834 (Hydropsychidae, Trichoptera) from the Lake Kinneret catchment (Israel). <i>Aquatic Insects</i> , 2017, 38, 125-140.	0.9	1
18	Integrating and extending ecological river assessment: Concept and test with two restoration projects. <i>Ecological Indicators</i> , 2017, 72, 131-141.	6.3	35

#	ARTICLE	IF	CITATIONS
19	Hydromorphological restoration stimulates river ecosystem metabolism. <i>Biogeosciences</i> , 2017, 14, 1989-2002.	3.3	22
20	Time is no healer: increasing restoration age does not lead to improved benthic invertebrate communities in restored river reaches. <i>Science of the Total Environment</i> , 2016, 557-558, 722-732.	8.0	52
21	Start at zero: succession of benthic invertebrate assemblages in restored former sewage channels. <i>Aquatic Sciences</i> , 2016, 78, 683-694.	1.5	16
22	Response of fish assemblages to hydromorphological restoration in central and northern European rivers. <i>Hydrobiologia</i> , 2016, 769, 67-78.	2.0	44
23	River restoration and the trophic structure of benthic invertebrate communities across 16 European restoration projects. <i>Hydrobiologia</i> , 2016, 769, 105-120.	2.0	26
24	The response of hydrophyte growth forms and plant strategies to river restoration. <i>Hydrobiologia</i> , 2016, 769, 41-54.	2.0	19
25	Fish community responses and the temporal dynamics of recovery following river habitat restorations in Europe. <i>Freshwater Science</i> , 2015, 34, 975-990.	1.8	28
26	Contrasting the roles of section length and instream habitat enhancement for river restoration success: a field study of 20 European restoration projects. <i>Journal of Applied Ecology</i> , 2015, 52, 1518-1527.	4.0	64
27	Bundles of stream restoration measures and their effects on fish communities. <i>Limnologica</i> , 2015, 55, 1-8.	1.5	10
28	A multi-trait approach for the identification and protection of European freshwater species that are potentially vulnerable to the impacts of climate change. <i>Ecological Indicators</i> , 2015, 50, 150-160.	6.3	37
29	Recolonisation patterns of benthic invertebrates: a field investigation of restored former sewage channels. <i>Freshwater Biology</i> , 2014, 59, 1932-1944.	2.4	32
30	Mountain river restoration measures and their success(ion): Effects on river morphology, local species pool, and functional composition of three organism groups. <i>Ecological Indicators</i> , 2014, 38, 243-255.	6.3	42
31	The Importance of the Regional Species Pool, Ecological Species Traits and Local Habitat Conditions for the Colonization of Restored River Reaches by Fish. <i>PLoS ONE</i> , 2014, 9, e84741.	2.5	65
32	Do adult and YOY fish benefit from river restoration measures?. <i>Ecological Engineering</i> , 2013, 61, 174-181.	3.6	56
33	Substratum associations of benthic invertebrates in lowland and mountain streams. <i>Ecological Indicators</i> , 2013, 30, 178-189.	6.3	42
34	Upstream river morphology and riparian land use overrule local restoration effects on ecological status assessment. <i>Hydrobiologia</i> , 2013, 704, 489-501.	2.0	102
35	Small and impoverished regional species pools constrain colonisation of restored river reaches by fishes. <i>Freshwater Biology</i> , 2013, 58, 664-674.	2.4	49
36	A comparison of habitat diversity and interannual habitat dynamics in actively and passively restored mountain rivers of Germany. <i>Hydrobiologia</i> , 2013, 712, 89-104.	2.0	12

#	ARTICLE	IF	CITATIONS
37	Macrophytes respond to reach-scale river restorations. <i>Journal of Applied Ecology</i> , 2012, 49, 202-212.	4.0	67
38	Dispersal as a limiting factor in the colonization of restored mountain streams by plants and macroinvertebrates. <i>Journal of Applied Ecology</i> , 2011, 48, 1241-1250.	4.0	100
39	Hydromorphological restoration of running waters: effects on benthic invertebrate assemblages. <i>Freshwater Biology</i> , 2011, 56, 1689-1702.	2.4	67
40	A comparative analysis of restoration measures and their effects on hydromorphology and benthic invertebrates in 26 central and southern European rivers. <i>Journal of Applied Ecology</i> , 2010, 47, 671-680.	4.0	128
41	Restoration effort, habitat mosaics, and macroinvertebrates – does channel form determine community composition?. <i>Aquatic Conservation: Marine and Freshwater Ecosystems</i> , 2009, 19, 157-169.	2.0	39
42	Re-Meandering German Lowland Streams: Qualitative and Quantitative Effects of Restoration Measures on Hydromorphology and Macroinvertebrates. <i>Environmental Management</i> , 2009, 44, 745-754.	2.7	76
43	Effects of rebraiding measures on hydromorphology, floodplain vegetation, ground beetles and benthic invertebrates in mountain rivers. <i>Journal of Applied Ecology</i> , 2009, 46, 406-416.	4.0	87
44	Substrate-specific macroinvertebrate diversity patterns following stream restoration. <i>Aquatic Sciences</i> , 2008, 70, 292-303.	1.5	52
45	Hydromorphological parameters indicating differences between single- and multiple-channel mountain rivers in Germany, in relation to their modification and recovery. <i>Aquatic Conservation: Marine and Freshwater Ecosystems</i> , 2008, 18, 1200-1216.	2.0	39
46	Sample coherence – a field study approach to assess similarity of macroinvertebrate samples. <i>Hydrobiologia</i> , 2006, 566, 461-476.	2.0	14
47	Effects of sampling and sub-sampling variation using the STAR-AQEM sampling protocol on the precision of macroinvertebrate metrics. <i>Hydrobiologia</i> , 2006, 566, 441-459.	2.0	45
48	The AQEM/STAR taxalist – a pan-European macro-invertebrate ecological database and taxa inventory. <i>Hydrobiologia</i> , 2006, 566, 325-342.	2.0	49
49	The AQEM/STAR taxalist – a pan-European macro-invertebrate ecological database and taxa inventory. , 2006, , 325-342.		6
50	Sample coherence – a field study approach to assess similarity of macroinvertebrate samples. , 2006, , 461-476.		1
51	Effects of sampling and sub-sampling variation using the STAR-AQEM sampling protocol on the precision of macroinvertebrate metrics. , 2006, , 441-459.		7
52	‘Electronic Subsampling’™ of Macrobenthic Samples: How Many Individuals are Needed for a Valid Assessment Result?. , 2004, , 299-312.		7
53	A new method for assessing the impact of hydromorphological degradation on the macroinvertebrate fauna of five German stream types. <i>Hydrobiologia</i> , 2004, 516, 107-127.	2.0	149
54	‘Electronic subsampling’ of macrobenthic samples: how many individuals are needed for a valid assessment result?. <i>Hydrobiologia</i> , 2004, 516, 299-312.	2.0	36

#	ARTICLE	IF	CITATIONS
55	Typology of streams in Germany based on benthic invertebrates: Ecoregions, zonation, geology and substrate. <i>Limnologica</i> , 2004, 34, 379-389.	1.5	46