

Petra Innerhofer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3959004/publications.pdf>

Version: 2024-02-01

13
papers

756
citations

1307543

7
h-index

1125717

13
g-index

13
all docs

13
docs citations

13
times ranked

532
citing authors

#	ARTICLE	IF	CITATIONS
1	First-Line Administration of Fibrinogen Concentrate in the Bleeding Trauma Patient: Searching for Effective Dosages and Optimal Post-Treatment Levels Limiting Massive Transfusion—Further Results of the RETIC Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 3930.	2.4	5
2	Efficacy of prehospital administration of fibrinogen concentrate in trauma patients bleeding or presumed to bleed (FlinTIC). <i>European Journal of Anaesthesiology</i> , 2021, 38, 348-357.	1.7	43
3	Dynamics of Platelet Counts in Major Trauma: The Impact of Haemostatic Resuscitation and Effects of Platelet Transfusion—A Sub-Study of the Randomized Controlled RETIC Trial. <i>Journal of Clinical Medicine</i> , 2020, 9, 2420.	2.4	5
4	Response patterns of routinely measured inflammatory and coagulatory parameters in sepsis. <i>PeerJ</i> , 2019, 7, e7147.	2.0	4
5	Inflammatory and coagulatory parameters linked to survival in critically ill children with sepsis. <i>Annals of Intensive Care</i> , 2018, 8, 111.	4.6	18
6	Reversal of trauma-induced coagulopathy using first-line coagulation factor concentrates or fresh frozen plasma (RETIC): a single-centre, parallel-group, open-label, randomised trial. <i>Lancet Haematology</i> , 2017, 4, e258-e271.	4.6	236
7	Early fibrinogen-concentrate administration in management of trauma-induced coagulopathy — Authors' reply. <i>Lancet Haematology</i> , 2017, 4, e348-e349.	4.6	4
8	FXIII Deficiency Occurs Frequently in Major Trauma and Influences Transfusion Requirements While Levels of Von Willebrand Factor are well above Thresholds. <i>Emergency Medicine Investigations</i> , 2017, 4, .	0.0	1
9	The exclusive use of coagulation factor concentrates enables reversal of coagulopathy and decreases transfusion rates in patients with major blunt trauma. <i>Injury</i> , 2013, 44, 209-216.	1.7	133
10	The impact of fresh frozen plasma vs coagulation factor concentrates on morbidity and mortality in trauma-associated haemorrhage and massive transfusion. <i>Injury</i> , 2011, 42, 697-701.	1.7	154
11	Principles of perioperative coagulopathy. <i>Bailliere's Best Practice and Research in Clinical Anaesthesiology</i> , 2010, 24, 1-14.	4.0	46
12	Fibrinogen and prothrombin complex concentrate for the management of massive bleeding. <i>Wiener Klinische Wochenschrift</i> , 2010, 122 Suppl 5, S16-7.	1.9	1
13	Time for changing coagulation management in trauma-related massive bleeding. <i>Current Opinion in Anaesthesiology</i> , 2009, 22, 267-274.	2.0	106