
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3967005/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Real-World Effectiveness of Antipsychotic Treatments in a Nationwide Cohort of 29†823 Patients With Schizophrenia. JAMA Psychiatry, 2017, 74, 686.	6.0	460
2	20â€year followâ€up study of physical morbidity and mortality in relationship to antipsychotic treatment in a nationwide cohort of 62,250 patients with schizophrenia (FIN20). World Psychiatry, 2020, 19, 61-68.	4.8	232
3	Association of Antipsychotic Polypharmacy vs Monotherapy With Psychiatric Rehospitalization Among Adults With Schizophrenia. JAMA Psychiatry, 2019, 76, 499.	6.0	191
4	Antipsychotics and mortality in a nationwide cohort of 29,823 patients with schizophrenia. Schizophrenia Research, 2018, 197, 274-280.	1.1	179
5	Mortality in people with schizophrenia: a systematic review and metaâ€analysis of relative risk and aggravating or attenuating factors. World Psychiatry, 2022, 21, 248-271.	4.8	156
6	From prescription drug purchases to drug use periods – a second generation method (PRE2DUP). BMC Medical Informatics and Decision Making, 2015, 15, 21.	1.5	137
7	20-Year Nationwide Follow-Up Study on Discontinuation of Antipsychotic Treatment in First-Episode Schizophrenia. American Journal of Psychiatry, 2018, 175, 765-773.	4.0	134
8	Comparative Effectiveness of Antipsychotic Drugs for Rehospitalization in Schizophrenia—A Nationwide Study With 20-Year Follow-up. Schizophrenia Bulletin, 2018, 44, 1381-1387.	2.3	117
9	Real-world Effectiveness of Pharmacologic Treatments for the Prevention of Rehospitalization in a Finnish Nationwide Cohort of Patients With Bipolar Disorder. JAMA Psychiatry, 2018, 75, 347.	6.0	116
10	The clinical course of schizophrenia in women and men—a nation-wide cohort study. NPJ Schizophrenia, 2020, 6, 12.	2.0	93
11	Mortality in schizophrenia: 30â€year nationwide followâ€up study. Acta Psychiatrica Scandinavica, 2018, 138, 492-499.	2.2	87
12	Impact of High Risk Drug Use on Hospitalization and Mortality in Older People with and without Alzheimer's Disease: A National Population Cohort Study. PLoS ONE, 2014, 9, e83224.	1.1	78
13	Cohort profile: the Finnish Medication and Alzheimer's disease (MEDALZ) study. BMJ Open, 2016, 6, e012100.	0.8	75
14	No Association Between Proton Pump Inhibitor Use and Risk of Alzheimer's Disease. American Journal of Gastroenterology, 2017, 112, 1802-1808.	0.2	69
15	Half a century of research on antipsychotics and schizophrenia: A scientometric study of hotspots, nodes, bursts, and trends. Neuroscience and Biobehavioral Reviews, 2022, 136, 104608.	2.9	67
16	Psychosis relapse during treatment with long-acting injectable antipsychotics in individuals with schizophrenia-spectrum disorders: an individual participant data meta-analysis. Lancet Psychiatry,the, 2020, 7, 749-761.	3.7	66
17	High prevalence of psychotropic drug use among persons with and without Alzheimer× ³ s disease in Finnish nationwide cohort. European Neuropsychopharmacology, 2014, 24, 1729-1737.	0.3	64
18	Agreement between PRE2DUP register data modeling method and comprehensive drug use interview among older persons. Clinical Epidemiology, 2016, Volume 8, 363-371.	1.5	61

#	Article	IF	CITATIONS
19	Antipsychotic use and risk of breast cancer in women with schizophrenia: a nationwide nested case-control study in Finland. Lancet Psychiatry,the, 2021, 8, 883-891.	3.7	50
20	Risk of Alzheimer's disease among users of postmenopausal hormone therapy: A nationwide case-control study. Maturitas, 2017, 98, 7-13.	1.0	47
21	Representation and Outcomes of Individuals With Schizophrenia Seen in Everyday Practice Who Are Ineligible for Randomized Clinical Trials. JAMA Psychiatry, 2022, 79, 210.	6.0	47
22	Risk of pneumonia associated with incident benzodiazepine use among community-dwelling adults with Alzheimer disease. Cmaj, 2017, 189, E519-E529.	0.9	46
23	Antidementia drug use among community-dwelling individuals with Alzheimer's disease in Finland. International Clinical Psychopharmacology, 2014, 29, 216-223.	0.9	44
24	Use of existing data sources in clinical epidemiology: Finnish health care registers in Alzheimer's disease research – the Medication use among persons with Alzheimer's disease (MEDALZ-2005) study. Clinical Epidemiology, 2013, 5, 277.	1.5	43
25	Comparison of predictors of hip fracture and mortality after hip fracture in community-dwellers with and without Alzheimer's disease – exposure-matched cohort study. BMC Geriatrics, 2016, 16, 204.	1.1	43
26	Hospital-treated mental and behavioral disorders and risk of Alzheimer's disease: A nationwide nested case-control study. European Psychiatry, 2017, 43, 92-98.	0.1	42
27	Use of Antiepileptic Drugs and Dementia Risk—an Analysis of Finnish Health Register and German Health Insurance Data. Journal of the American Geriatrics Society, 2018, 66, 1123-1129.	1.3	41
28	Adenosine modulators and calcium channel blockers as add-on treatment for schizophrenia. NPJ Schizophrenia, 2021, 7, 1.	2.0	40
29	Employment among people with schizophrenia or bipolar disorder: A populationâ€based study using nationwide registers. Acta Psychiatrica Scandinavica, 2021, 143, 61-71.	2.2	39
30	Pharmacological treatments and risk of readmission to hospital for unipolar depression in Finland: a nationwide cohort study. Lancet Psychiatry,the, 2017, 4, 547-553.	3.7	39
31	Research Priorities for Optimizing Geriatric Pharmacotherapy: An International Consensus. Journal of the American Medical Directors Association, 2018, 19, 193-199.	1.2	35
32	Comparative Effectiveness of Antipsychotics for Risk of Attempted or Completed Suicide Among Persons With Schizophrenia. Schizophrenia Bulletin, 2021, 47, 23-30.	2.3	35
33	Long-term use of opioids for nonmalignant pain among community-dwelling persons with and without Alzheimer disease in Finland: a nationwide register-based study. Pain, 2017, 158, 252-260.	2.0	34
34	Antipsychotic Use and Risk of Hospitalization or Death Due to Pneumonia in Persons With and Those Without Alzheimer Disease. Chest, 2016, 150, 1233-1241.	0.4	33
35	The risk of Alzheimer's disease associated with benzodiazepines and related drugs: a nested case–control study. Acta Psychiatrica Scandinavica, 2018, 138, 91-100.	2.2	33
36	Long-term Continuity of Antipsychotic Treatment for Schizophrenia: A Nationwide Study. Schizophrenia Bulletin, 2021, 47, 1611-1620.	2.3	33

#	Article	IF	CITATIONS
37	Long-term use of benzodiazepines and related drugs among community-dwelling individuals with and without Alzheimer's disease. International Clinical Psychopharmacology, 2015, 30, 202-208.	0.9	32
38	Drug exposure in register-based research—An expert-opinion based evaluation of methods. PLoS ONE, 2017, 12, e0184070.	1.1	31
39	Hospital admissions, outpatient visits and healthcare costs of communityâ€dwellers with Alzheimer's disease. Alzheimer's and Dementia, 2015, 11, 955-963.	0.4	30
40	Head or brain injuries and Alzheimer's disease: A nested caseâ€control register study. Alzheimer's and Dementia, 2017, 13, 1371-1379.	0.4	30
41	In the aftermath of clozapine discontinuation: comparative effectiveness and safety of antipsychotics in patients with schizophrenia who discontinue clozapine. British Journal of Psychiatry, 2020, 217, 498-505.	1.7	30
42	The Incidence of Benzodiazepine andÂRelated Drug Use in Persons withÂandÂwithout Alzheimer's Disease. Journal of Alzheimer's Disease, 2015, 49, 809-818.	1.2	27
43	Antidepressant use and risk of hip fractures among communityâ€dwelling persons with and without Alzheimer's disease. International Journal of Geriatric Psychiatry, 2017, 32, e107-e115.	1.3	27
44	Risk of Hip Fracture in Benzodiazepine Users With and Without Alzheimer Disease. Journal of the American Medical Directors Association, 2017, 18, 87.e15-87.e21.	1.2	26
45	Psychotropic drugs use and psychotropic polypharmacy among persons with Alzheimer's disease. European Neuropsychopharmacology, 2018, 28, 1260-1269.	0.3	26
46	Psychosis breakthrough on antipsychotic maintenance: results from a nationwide study. Psychological Medicine, 2020, 50, 1356-1367.	2.7	26
47	Antipsychotics Use Is Associated With Greater Adherence to Cardiometabolic Medications in Patients With Schizophrenia: Results From a Nationwide, Within-subject Design Study. Schizophrenia Bulletin, 2022, 48, 166-175.	2.3	26
48	Incidence of antidepressant use in community-dwelling persons with and without Alzheimer's disease: 13-year follow-up. International Journal of Geriatric Psychiatry, 2017, 32, 94-101.	1.3	25
49	Morbidity and mortality in schizophrenia with comorbid substance use disorders. Acta Psychiatrica Scandinavica, 2021, 144, 42-49.	2.2	25
50	Real-world effectiveness of antipsychotic doses for relapse prevention in patients with first-episode schizophrenia in Finland: a nationwide, register-based cohort study. Lancet Psychiatry,the, 2022, 9, 271-279.	3.7	25
51	Long-term use of antipsychotics among community-dwelling persons with Alzheimer׳s disease: A nationwide register-based study. European Neuropsychopharmacology, 2015, 25, 1706-1713.	0.3	24
52	Risk of Mortality Associated with Antipsychotic Monotherapy and Polypharmacy Among Community-Dwelling Persons with Alzheimer's Disease. Journal of Alzheimer's Disease, 2017, 56, 107-118.	1.2	24
53	Proton pump inhibitor use and risk of hip fractures among communityâ€dwelling persons with Alzheimer's disease—a nested caseâ€control study. Alimentary Pharmacology and Therapeutics, 2018, 47, 1135-1142.	1.9	24
54	Comparison of dispensed medications and forensicâ€ŧoxicological findings to assess pharmacotherapy in the Swedish population 2006 to 2013. Pharmacoepidemiology and Drug Safety, 2018, 27, 1112-1122.	0.9	24

#	Article	IF	CITATIONS
55	Risk of death associated with new benzodiazepine use among persons with Alzheimer disease: A matched cohort study. International Journal of Geriatric Psychiatry, 2018, 33, 583-590.	1.3	24
56	Burden for Parents of Patients With Schizophrenia—A Nationwide Comparative Study of Parents of Offspring With Rheumatoid Arthritis, Multiple Sclerosis, Epilepsy, and Healthy Controls. Schizophrenia Bulletin, 2019, 45, 794-803.	2.3	24
57	Incidence of and Characteristics Associated With Long-term Benzodiazepine Use in Finland. JAMA Network Open, 2020, 3, e2019029.	2.8	24
58	Risk Factors, Incidence, and Outcomes of Neuroleptic Malignant Syndrome on Long-Acting Injectable vs Oral Antipsychotics in a Nationwide Schizophrenia Cohort. Schizophrenia Bulletin, 2021, 47, 1621-1630.	2.3	24
59	Differences in analgesic use in communityâ€dwelling persons with and without Alzheimer's disease. European Journal of Pain, 2017, 21, 658-667.	1.4	23
60	Antipsychotic Use and the Risk of Hip Fracture Among Community-Dwelling Persons With Alzheimer's Disease. Journal of Clinical Psychiatry, 2017, 78, e257-e263.	1.1	23
61	Incidence of antipsychotic use in relation to diagnosis of Alzheimer's disease among community-dwelling persons. British Journal of Psychiatry, 2015, 207, 444-449.	1.7	22
62	lnitiation and longâ€ŧerm use of benzodiazepines and Zâ€drugs in bipolar disorder. Bipolar Disorders, 2018, 20, 634-646.	1.1	22
63	Adherence to psychotropic medication in completed suicide in Sweden 2006–2013: a forensic-toxicological matched case-control study. European Journal of Clinical Pharmacology, 2019, 75, 1421-1430.	0.8	22
64	Hospital care and drug costs from five years before until two years after the diagnosis of Alzheimer's disease in a Finnish nationwide cohort. Scandinavian Journal of Public Health, 2016, 44, 150-158.	1.2	21
65	Incident opioid use and risk of hip fracture among persons with Alzheimer disease: a nationwide matched cohort study. Pain, 2019, 160, 417-423.	2.0	20
66	Suicide mortality and use of psychotropic drugs in patients hospitalized due to bipolar disorder: A Finnish nationwide cohort study. Journal of Affective Disorders, 2020, 277, 885-892.	2.0	20
67	Antipsychotic Polypharmacy among a Nationwide Sample of Community-Dwelling Persons with Alzheimer's Disease. Journal of Alzheimer's Disease, 2014, 41, 1223-1228.	1.2	19
68	Primary Nonadherence to Antipsychotic Treatment Among Persons with Schizophrenia. Schizophrenia Bulletin, 2022, 48, 655-663.	2.3	19
69	Antipsychotic Doses Among Community-Dwelling Persons With Alzheimer Disease in Finland. Journal of Clinical Psychopharmacology, 2014, 34, 435-440.	0.7	18
70	The Risk of Head Injuries Associated With Antipsychotic Use Among Persons With Alzheimer's disease. Journal of the American Geriatrics Society, 2020, 68, 595-602.	1.3	16
71	Realâ€world effectiveness of pharmacological treatments of alcohol use disorders in a Swedish nationâ€wide cohort of 125 556 patients. Addiction, 2021, 116, 1990-1998.	1.7	16
72	Registry-based studies: What they can tell us, and what they cannot. European Neuropsychopharmacology, 2021, 45, 35-37.	0.3	16

#	Article	IF	CITATIONS
73	Antipsychotic use among persons with schizophrenia in Sweden and Finland, trends and differences. Nordic Journal of Psychiatry, 2021, 75, 315-322.	0.7	15
74	Long-term treatment with clozapine and other antipsychotic drugs and the risk of haematological malignancies in people with schizophrenia: a nationwide case-control and cohort study in Finland. Lancet Psychiatry,the, 2022, 9, 353-362.	3.7	15
75	Drug use in persons with and without Alzheimer's disease aged 90 years or more. Age and Ageing, 2016, 45, 900-904.	0.7	14
76	A Comparison of Sex Differences in Psychotropic Medication Use in Older People with Alzheimer's Disease in the US and Finland. Drugs and Aging, 2017, 34, 55-65.	1.3	14
77	Molecular signaling pathways underlying schizophrenia. Schizophrenia Research, 2021, 232, 33-41.	1.1	14
78	Use of benzodiazepines and related drugs is associated with a risk of stroke among persons with Alzheimer's disease. International Clinical Psychopharmacology, 2017, 32, 135-141.	0.9	13
79	Use of antidepressants and mood stabilizers in persons with first-episode schizophrenia. European Journal of Clinical Pharmacology, 2020, 76, 711-718.	0.8	13
80	Use of antidementia drugs and risk of pneumonia in older persons with Alzheimer's disease. Annals of Medicine, 2017, 49, 230-239.	1.5	12
81	Incidence and Prevalence of Antiepileptic Medication Use in Community-Dwelling Persons with and without Alzheimer's Disease. Journal of Alzheimer's Disease, 2018, 66, 387-395.	1.2	11
82	Risk and risk factors for disability pension among patients with treatment resistant depression– a matched cohort study. BMC Psychiatry, 2020, 20, 232.	1.1	11
83	Incidence of antidepressant use among community dwellers with and without Parkinson's disease – a nationwide cohort study. BMC Geriatrics, 2021, 21, 202.	1.1	11
84	Incidence and Duration of Cumulative Bisphosphonate Use among Community-Dwelling Persons with or without Alzheimer's Disease. Journal of Alzheimer's Disease, 2016, 52, 127-132.	1.2	10
85	Sedative load and salivary secretion and xerostomia in communityâ€dwelling older people. Gerodontology, 2016, 33, 177-184.	0.8	10
86	Risk Factors for Initiation of Potentially Inappropriate Medications in Community-Dwelling Older Adults with and without Alzheimer's Disease. Drugs and Aging, 2017, 34, 67-77.	1.3	10
87	Is Alzheimer's Disease Associated with Previous Opioid Use?. Pain Medicine, 2018, 19, 2115-2121.	0.9	10
88	Predictors of high hospital care and medication costs and cost trajectories in community-dwellers with Alzheimer's disease. Annals of Medicine, 2019, 51, 294-305.	1.5	10
89	Antiepileptic drug use and mortality among community-dwelling persons with Alzheimer disease. Neurology, 2020, 94, e2099-e2108.	1.5	10
90	Long-Term Real-World Effectiveness of Pharmacotherapies for Schizoaffective Disorder. Schizophrenia Bulletin, 2021, 47, 1099-1107.	2.3	10

#	Article	IF	CITATIONS
91	Effectiveness of pharmacotherapies for delusional disorder in a Swedish national cohort of 9076 patients. Schizophrenia Research, 2021, 228, 367-372.	1.1	10
92	Concomitant Use of Acetylcholine Esterase Inhibitors and Urinary Antispasmodics Among Finnish Community-Dwelling Persons With Alzheimer Disease. Journal of Clinical Psychopharmacology, 2014, 34, 722-727.	0.7	9
93	Anticholinergic Exposure and Risk of Pneumonia in Persons with Alzheimer's Disease: A Nested Case-Control Study. Journal of Alzheimer's Disease, 2017, 56, 119-128.	1.2	9
94	Length of Hospital Stay for Hip Fracture and 30-Day Mortality in People With Alzheimer's Disease: A Cohort Study in Finland. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2020, 75, 2184-2192.	1.7	9
95	Month and Season of Birth as a Risk Factor for Alzheimer's Disease: A Nationwide Nested Case-control Study. Journal of Preventive Medicine and Public Health, 2016, 49, 134-138.	0.7	9
96	Optimal Doses of Specific Antipsychotics for Relapse Prevention in a Nationwide Cohort of Patients with Schizophrenia. Schizophrenia Bulletin, 2022, 48, 774-784.	2.3	9
97	Use of Cholinesterase Inhibitors Increases Initiation of Urinary Anticholinergics in Persons with Alzheimer's Disease. Journal of the American Geriatrics Society, 2016, 64, 1510-1512.	1.3	8
98	Incidence of head injury and traumatic brain injury among people with Alzheimer's disease. Journal of Epidemiology and Community Health, 2019, 73, 451-454.	2.0	8
99	Duration of new antidepressant use and factors associated with discontinuation among community-dwelling persons with Alzheimer's disease. European Journal of Clinical Pharmacology, 2019, 75, 417-425.	0.8	8
100	Medication Use and Health Care Utilization After a Cost-sharing Increase in Schizophrenia. Medical Care, 2020, 58, 763-769.	1.1	8
101	Courses of treatment and risk factors for treatment-resistant depression in Finnish primary and special healthcare: A nationwide cohort study. Journal of Affective Disorders, 2022, 308, 236-242.	2.0	8
102	Associations across spatial patterns of disease incidences, socio-demographics, and land use in Finland 1991–2010. Scandinavian Journal of Public Health, 2015, 43, 356-363.	1.2	7
103	Risk of head and traumatic brain injuries associated with antidepressant use among community-dwelling persons with Alzheimer's disease: a nationwide matched cohort study. Alzheimer's Research and Therapy, 2017, 9, 59.	3.0	7
104	Causes of death in a nationwide cohort of community-dwellers with Alzheimer's disease. BMC Geriatrics, 2020, 20, 441.	1.1	7
105	Initiation of antidepressant use among refugee and Swedish-born youth after diagnosis of a common mental disorder: findings from the REMAIN study. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 463-474.	1.6	7
106	Concomitant use of benzodiazepines and opioids in communityâ€dwelling older people with or without Alzheimer's disease—A nationwide registerâ€based study in <scp>Finland</scp> . International Journal of Geriatric Psychiatry, 2019, 34, 280-288.	1.3	6
107	Asthma and Chronic Obstructive Pulmonary Disease as a Comorbidity and Association with the Choice of Antidementia Medication Among Persons with Alzheimer's Disease. Journal of Alzheimer's Disease, 2020, 73, 1243-1251.	1.2	6
108	Risk Factors for Disability Pension among Young Adults Diagnosed with Attention-deficit Hyperactivity Disorder (ADHD) in Adulthood. Journal of Attention Disorders, 2022, 26, 723-734.	1.5	6

#	Article	IF	CITATIONS
109	Labour market marginalisation in young adults diagnosed with attention-deficit hyperactivity disorder (ADHD): a population-based longitudinal cohort study in Sweden. Psychological Medicine, 2023, 53, 1224-1232.	2.7	6
110	Persistence of Antipsychotic Use After Clozapine Discontinuation: A Realâ€World Study Across Antipsychotics. Clinical and Translational Science, 2020, 13, 1170-1177.	1.5	6
111	The effect of comprehensive geriatric assessment on anticholinergic exposure assessed by four ranked anticholinergic lists. Archives of Gerontology and Geriatrics, 2017, 68, 195-201.	1.4	5
112	Accumulation of Hospital Days Among Antipsychotic Initiators With Alzheimer's Disease. Journal of the American Medical Directors Association, 2019, 20, 1488-1494.e3.	1.2	5
113	Long-term thiazide use and risk of low-energy fractures among persons with Alzheimer's disease—nested case-control study. Osteoporosis International, 2019, 30, 1481-1489.	1.3	5
114	Incidence of schizophrenia and influence of prenatal and infant exposure to viral infectious diseases. Acta Psychiatrica Scandinavica, 2021, 143, 487-494.	2.2	5
115	Does Recent Hospitalization Increase Antipsychotic Initiation Among Community Dwellers With Alzheimer's Disease?. Journal of the American Medical Directors Association, 2021, 22, 1543-1547.e3.	1.2	5
116	Regional changes in psychotropic use among Finnish persons with newly diagnosed Alzheimer's disease in 2005-2011. PLoS ONE, 2017, 12, e0173450.	1.1	5
117	Antipsychotic Use and Risk of Stroke Among Community-Dwelling People With Alzheimer's Disease. Journal of the American Medical Directors Association, 2022, 23, 1059-1065.e4.	1.2	5
118	Realâ€world effectiveness of pharmacological treatments of opioid use disorder in a national cohort. Addiction, 2022, 117, 1683-1691.	1.7	5
119	Antiepileptic Drugs and Accumulation of Hospital Days Among Persons With Alzheimer's Disease. Journal of the American Medical Directors Association, 2019, 20, 751-758.	1.2	4
120	Prevalence of cardiovascular drug use before and after diagnosis of Alzheimer's disease. International Journal of Cardiology, 2020, 300, 221-225.	0.8	4
121	Coronary Revascularization and Postoperative Outcomes in People With and Without Alzheimer's Disease. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2021, 76, 1524-1530.	1.7	4
122	Association of recent hospitalisation with antidepressant initiation among community dwellers with Alzheimer's disease. International Journal of Geriatric Psychiatry, 2021, 36, 1075-1084.	1.3	4
123	Persistence of antidepressant use among refugee youth with common mental disorder. Journal of Affective Disorders, 2021, 294, 831-837.	2.0	4
124	Antidepressant use among immigrants with depressive disorder living in Finland: A register-based study. Journal of Affective Disorders, 2022, 299, 528-535.	2.0	4
125	Prescription opioids among older adults: ten years of data across five countries. BMC Geriatrics, 2022, 22, 429.	1.1	4
126	Prevalence of Drug Combinations Increasing Bleeding Risk Among Warfarin Users With and Without Alzheimer's Disease. Drugs and Aging, 2015, 32, 937-945.	1.3	3

#	Article	IF	CITATIONS
127	Sedative load, carious teeth and infection in the periodontium among communityâ€dwelling older people. Gerodontology, 2017, 34, 13-23.	0.8	3
128	Screening approach for identifying candidate drugs and drugâ€drug interactions related to hip fracture risk in persons with Alzheimer disease. Pharmacoepidemiology and Drug Safety, 2017, 26, 875-889.	0.9	3
129	Systemic Estrogen Use and Discontinuation After Alzheimer's Disease Diagnosis in Finland 2005–2012: A Nationwide Exposure-Matched Cohort Study. Drugs and Aging, 2018, 35, 985-992.	1.3	3
130	Commentary on Robert Whitaker's viewpoint. Psychological Medicine, 2020, 50, 2653-2654.	2.7	3
131	Trajectories of antidepressant use and characteristics associated with trajectory groups among young refugees and their Swedish-born peers with diagnosed common mental disorders—findings from the REMAIN study. Social Psychiatry and Psychiatric Epidemiology, 2022, 57, 305-317.	1.6	3
132	Switches and early discontinuations of antidepressant medication in young adults with depression. Journal of Affective Disorders, 2021, 295, 1474-1481.	2.0	3
133	Trajectories of labour market marginalisation among young adults with newly diagnosed attention-deficit/hyperactivity disorder (ADHD). Epidemiology and Psychiatric Sciences, 2021, 30, e67.	1.8	3
134	Allopurinol, dipyridamole and calcium channel blockers in the treatment of bipolar disorder – A nationwide cohort study. Journal of Affective Disorders, 2022, 313, 43-48.	2.0	3
135	Detection of medications associated with Alzheimer's disease using ensemble methods and cooperative game theory. International Journal of Medical Informatics, 2020, 141, 104142.	1.6	2
136	Health and work disability outcomes in parents of patients with schizophrenia associated with antipsychotic exposure by the offspring. Scientific Reports, 2020, 10, 1219.	1.6	2
137	Hip Fracture Risk in Antiepileptic Drug Initiators and Non-Initiators with Alzheimer's Disease. Clinical Epidemiology, 2021, Volume 13, 295-307.	1.5	2
138	Changes in the use of diabetes drugs among community-dwelling people with Alzheimer's disease. BMC Geriatrics, 2021, 21, 701.	1.1	2
139	Occupational Branch and Labor Market Marginalization among Young Employees with Adult Onset of Attention Deficit Hyperactivity Disorder—A Population-Based Matched Cohort Study. International Journal of Environmental Research and Public Health, 2022, 19, 7254.	1.2	2
140	Antipsychotic Use and the Risk of Initiating Medication for Benign Prostate Hyperplasia in Persons With Alzheimer Disease. Journal of Clinical Psychopharmacology, 2018, 38, 494-497.	0.7	1
141	Risk of Treatment Failure: Response to Ghosh and Noble. American Journal of Psychiatry, 2018, 175, 1267-1267.	4.0	1
142	Hospitalization after Oral Antibiotic Initiation in Finnish Community Dwellers with and without Alzheimer's Disease: Retrospective Register-Based Cohort Study. Journal of Alzheimer's Disease, 2018, 64, 437-445.	1.2	1
143	Opioid Use After Hospital Care due to Hip Fracture Among Community-Dwelling Persons With and Without Alzheimer's Disease. Drugs and Aging, 2020, 37, 193-203.	1.3	1
144	Continuity of antipsychotic medication use among migrant and Finnish-born populations with a psychotic disorder: a register-based study. Psychological Medicine, 2023, 53, 833-843.	2.7	1

#	Article	IF	CITATIONS
145	Incidence of Bisphosphonate Use in Relation to Diagnosis of Alzheimer's Disease in Communityâ€Đwelling Persons. Journal of the American Geriatrics Society, 2016, 64, e48-9.	1.3	0
146	Reply to: Use of Antiepileptic Drugs and Dementia Risk—an Analysis of Finnish Health Register and German Health Insurance Data. Journal of the American Geriatrics Society, 2018, 66, 1853-1854.	1.3	0
147	Risk of Treatment Failure: Response to Czobor and Volavka. American Journal of Psychiatry, 2018, 175, 909-909.	4.0	Ο
148	S210. COMPARATIVE EFFECTIVENESS OF ANTIPSYCHOTICS FOR RISK OF ATTEMPTED OR COMPLETED SUICIDE AMONG PERSONS WITH SCHIZOPHRENIA. Schizophrenia Bulletin, 2020, 46, S118-S119.	2.3	0
149	T208. EFFECTIVENESS OF PHARMACOLOGICAL THERAPIES FOR DELUSIONAL DISORDER. Schizophrenia Bulletin, 2020, 46, S311-S312.	2.3	0
150	Rejoinder to rejoinder to commentary. Psychological Medicine, 2020, 50, 2812-2813.	2.7	0
151	Automated multi-dose dispensing in persons with and without Alzheimer's disease—impacts on pharmacotherapy. European Journal of Clinical Pharmacology, 2022, 78, 513-521.	0.8	0
152	Effectiveness of antipsychotics in schizophrenia with comorbid substance use disorder. European Psychiatry, 2021, 64, S161-S161.	0.1	0
153	Use of pharmacotherapies for treatment resistant depression in finland: A nationwide cohort study. European Psychiatry, 2021, 64, S108-S108.	0.1	0
154	Morbidity and mortality in schizophrenia with comorbid substance use disorders in Finland and Sweden. European Psychiatry, 2021, 64, S237-S238.	0.1	0
155	Real-world effectiveness studies of low doses of antipsychotics – Authors' reply. Lancet Psychiatry the 2022 9 536-537	3.7	0