

Senthilkumar S Karuppagounder

List of Publications by Year in Descending Order

Source:
<https://exaly.com/author-pdf/3967031/senthilkumar-s-karuppagounder-publications-by-year.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

52 papers	3,680 citations	29 h-index	56 g-index
56 ext. papers	4,681 ext. citations	10.9 avg, IF	4.82 L-index

#	Paper	IF	Citations
52	The Stroke Preclinical Assessment Network: Rationale, Design, Feasibility, and Stage 1 Results.. <i>Stroke</i> , 2022 , 101161STROKEAHA121038047	6.7	2
51	STING mediates neurodegeneration and neuroinflammation in nigrostriatal β -synucleinopathy.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022 , 119, e2118819119 ^{11.5}	11.5	5
50	A high-affinity cocaine binding site associated with the brain acid soluble protein 1.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022 , 119, e2200545119	11.5	0
49	TRIP12 ubiquitination of glucocerebrosidase contributes to neurodegeneration in Parkinson's disease. <i>Neuron</i> , 2021 , 109, 3758-3774.e11	13.9	1
48	PARIS farnesylation prevents neurodegeneration in models of Parkinson's disease. <i>Science Translational Medicine</i> , 2021 , 13,	17.5	6
47	Caffeine, a natural methylxanthine nutraceutical, exerts dopaminergic neuroprotection. <i>Neurochemistry International</i> , 2021 , 148, 105066	4.4	2
46	PARIS induced defects in mitochondrial biogenesis drive dopamine neuron loss under conditions of parkin or PINK1 deficiency. <i>Molecular Neurodegeneration</i> , 2020 , 15, 17	19	31
45	NLRP3 inflammasome activation in dopamine neurons contributes to neurodegeneration in Parkinson's Disease. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	4
44	Defects in Mitochondrial Biogenesis Drive Mitochondrial Alterations in PARKIN-Deficient Human Dopamine Neurons. <i>Stem Cell Reports</i> , 2020 , 15, 629-645	8	21
43	Transneuronal Propagation of Pathologic β -Synuclein from the Gut to the Brain Models Parkinson's Disease. <i>Neuron</i> , 2019 , 103, 627-641.e7	13.9	453
42	Parkin interacting substrate zinc finger protein 746 is a pathological mediator in Parkinson's disease. <i>Brain</i> , 2019 , 142, 2380-2401	11.2	21
41	FAM19A1, a brain-enriched and metabolically responsive neurokinin, regulates food intake patterns and mouse behaviors. <i>FASEB Journal</i> , 2019 , 33, 14734-14747	0.9	10
40	Synthetic mRNAs Drive Highly Efficient iPS Cell Differentiation to Dopaminergic Neurons. <i>Stem Cells Translational Medicine</i> , 2019 , 8, 112-123	6.9	28
39	Robust kinase- and age-dependent dopaminergic and norepinephrine neurodegeneration in LRRK2 G2019S transgenic mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 1635-1640	11.5	45
38	GBA1 deficiency negatively affects physiological β -synuclein tetramers and related multimers. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 798-803	11.5	106
37	β -Synuclein accumulation and GBA deficiency due to L444P GBA mutation contributes to MPTP-induced parkinsonism. <i>Molecular Neurodegeneration</i> , 2018 , 13, 1	19	63
36	Block of A1 astrocyte conversion by microglia is neuroprotective in models of Parkinson's disease. <i>Nature Medicine</i> , 2018 , 24, 931-938	50.5	413

35	Poly(ADP-ribose) drives pathologic β -synuclein neurodegeneration in Parkinson's disease. <i>Science</i> , 2018 , 362,	33.3	196
34	Synaptic Plasticity onto Dopamine Neurons Shapes Fear Learning. <i>Neuron</i> , 2017 , 93, 425-440	13.9	31
33	c-Abl and Parkinson's Disease: Mechanisms and Therapeutic Potential. <i>Journal of Parkinson's Disease</i> , 2017 , 7, 589-601	5.3	41
32	Overexpression of Parkinson's Disease-Associated Mutation LRRK2 G2019S in Mouse Forebrain Induces Behavioral Deficits and β -Synuclein Pathology. <i>ENeuro</i> , 2017 , 4,	3.9	19
31	Activation of tyrosine kinase c-Abl contributes to β -Synuclein-induced neurodegeneration. <i>Journal of Clinical Investigation</i> , 2016 , 126, 2970-88	15.9	88
30	Adult Conditional Knockout of PGC-1 α Leads to Loss of Dopamine Neurons. <i>ENeuro</i> , 2016 , 3,	3.9	61
29	Methamphetamine-induced dopaminergic toxicity prevented owing to the neuroprotective effects of salicylic acid. <i>Life Sciences</i> , 2016 , 154, 24-9	6.8	26
28	LRRK2 G2019S transgenic mice display increased susceptibility to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-mediated neurotoxicity. <i>Journal of Chemical Neuroanatomy</i> , 2016 , 76, 90-97	3.2	29
27	Pathological β -Synuclein transmission initiated by binding lymphocyte-activation gene 3. <i>Science</i> , 2016 , 353,	33.3	364
26	The c-Abl inhibitor, nilotinib, protects dopaminergic neurons in a preclinical animal model of Parkinson's disease. <i>Scientific Reports</i> , 2014 , 4, 4874	4.9	145
25	MicroRNA-132 dysregulation in <i>Toxoplasma gondii</i> infection has implications for dopamine signaling pathway. <i>Neuroscience</i> , 2014 , 268, 128-38	3.9	77
24	Poly(ADP-ribose) polymerase-dependent energy depletion occurs through inhibition of glycolysis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 10209-14	11.5	203
23	Genetic deficiency of the mitochondrial protein PGAM5 causes a Parkinson's-like movement disorder. <i>Nature Communications</i> , 2014 , 5, 4930	17.4	87
22	Elucidating the neurotoxic effects of MDMA and its analogs. <i>Life Sciences</i> , 2014 , 101, 37-42	6.8	18
21	Proneural transcription factor Atoh1 drives highly efficient differentiation of human pluripotent stem cells into dopaminergic neurons. <i>Stem Cells Translational Medicine</i> , 2014 , 3, 888-98	6.9	30
20	Assessment of therapeutic potential of amantadine in methamphetamine induced neurotoxicity. <i>Neurochemical Research</i> , 2013 , 38, 2084-94	4.6	13
19	Parthanatos mediates AIMP2-activated age-dependent dopaminergic neuronal loss. <i>Nature Neuroscience</i> , 2013 , 16, 1392-400	25.5	142
18	Selective inhibition of phosphodiesterase 5 enhances glutamatergic synaptic plasticity and memory in mice. <i>Synapse</i> , 2013 , 67, 741-7	2.4	9

17	Sulphydration mediates neuroprotective actions of parkin. <i>Nature Communications</i> , 2013 , 4, 1626	17.4	201
16	Developmental nicotine exposure induced alterations in behavior and glutamate receptor function in hippocampus. <i>Cellular and Molecular Life Sciences</i> , 2012 , 69, 829-41	10.3	49
15	Central insulin resistance and synaptic dysfunction in intracerebroventricular-streptozotocin injected rodents. <i>Neurobiology of Aging</i> , 2012 , 33, 430.e5-18	5.6	58
14	Investigate the chronic neurotoxic effects of diquat. <i>Neurochemical Research</i> , 2012 , 37, 1102-11	4.6	17
13	Phosphorylation by the c-Abl protein tyrosine kinase inhibits parkin's ubiquitination and protective function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 16691-6	11.5	199
12	Neurotoxic effects of methamphetamine. <i>Neurochemical Research</i> , 2010 , 35, 171-9	4.6	45
11	Effect of dopaminergic neurotoxin MPTP/MPP+ on coenzyme Q content. <i>Life Sciences</i> , 2008 , 83, 92-5	6.8	4
10	Role of lipoamide dehydrogenase and metallothionein on 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced neurotoxicity. <i>Neurochemical Research</i> , 2008 , 33, 980-4	4.6	13
9	Unilateral implantation of dopamine-loaded biodegradable hydrogel in the striatum attenuates motor abnormalities in the 6-hydroxydopamine model of hemi-parkinsonism. <i>Behavioural Brain Research</i> , 2007 , 184, 11-8	3.4	34
8	Evaluation of neuroprotective and anti-fatigue effects of sildenafil. <i>Life Sciences</i> , 2007 , 81, 988-92	6.8	30
7	Versatile effects of sildenafil: recent pharmacological applications. <i>Pharmacological Reports</i> , 2007 , 59, 150-63	3.9	55
6	Paraquat and maneb induced neurotoxicity. <i>Proceedings of the Western Pharmacology Society</i> , 2007 , 50, 31-42		20
5	Acute intranigral homocysteine administration produces stereotypic behavioral changes and striatal dopamine depletion in Sprague-Dawley rats. <i>Brain Research</i> , 2006 , 1075, 81-92	3.7	20
4	Ebselen effects on MPTP-induced neurotoxicity. <i>Brain Research</i> , 2006 , 1118, 251-4	3.7	17
3	Rats with unilateral median forebrain bundle, but not striatal or nigral, lesions by the neurotoxins MPP+ or rotenone display differential sensitivity to amphetamine and apomorphine. <i>Pharmacology Biochemistry and Behavior</i> , 2006 , 84, 321-9	3.9	46
2	L-deprenyl protects against rotenone-induced, oxidative stress-mediated dopaminergic neurodegeneration in rats. <i>Neurochemistry International</i> , 2006 , 49, 28-40	4.4	66
1	Aplp1 and the Aplp1-Lag3 Complex facilitates transmission of pathologic β -synuclein		3