

Takayoshi Ohkubo

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3967961/takayoshi-ohkubo-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

427
papers

31,860
citations

70
h-index

172
g-index

453
ext. papers

37,532
ext. citations

4.6
avg, IF

6.09
L-index

#	Paper	IF	Citations
427	Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. <i>Lancet, The</i> , 2014 , 384, 766-81	40	7175
426	Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. <i>Nature</i> , 2011 , 478, 103-9	50.4	1564
425	Global Burden of Hypertension and Systolic Blood Pressure of at Least 110 to 115 mm Hg, 1990-2015. <i>JAMA - Journal of the American Medical Association</i> , 2017 , 317, 165-182	27.4	957
424	Global, regional, and national levels and causes of maternal mortality during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. <i>Lancet, The</i> , 2014 , 384, 980-1004	40	950
423	European Society of Hypertension position paper on ambulatory blood pressure monitoring. <i>Journal of Hypertension</i> , 2013 , 31, 1731-68	1.9	898
422	Prognostic significance of the nocturnal decline in blood pressure in individuals with and without high 24-h blood pressure: the Ohasama study. <i>Journal of Hypertension</i> , 2002 , 20, 2183-9	1.9	767
421	Prognostic accuracy of day versus night ambulatory blood pressure: a cohort study. <i>Lancet, The</i> , 2007 , 370, 1219-29	40	655
420	Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. <i>Lancet, The</i> , 2014 , 384, 1005-70	40	653
419	Decline in estimated glomerular filtration rate and subsequent risk of end-stage renal disease and mortality. <i>JAMA - Journal of the American Medical Association</i> , 2014 , 311, 2518-2531	27.4	580
418	European Society of Hypertension guidelines for blood pressure monitoring at home: a summary report of the Second International Consensus Conference on Home Blood Pressure Monitoring. <i>Journal of Hypertension</i> , 2008 , 26, 1505-26	1.9	578
417	European Society of Hypertension practice guidelines for ambulatory blood pressure monitoring. <i>Journal of Hypertension</i> , 2014 , 32, 1359-66	1.9	547
416	Home blood pressure measurement has a stronger predictive power for mortality than does screening blood pressure measurement: a population-based observation in Ohasama, Japan. <i>Journal of Hypertension</i> , 1998 , 16, 971-5	1.9	521
415	Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. <i>Lancet, The</i> , 2014 , 384, 957-79	40	497
414	The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2019). <i>Hypertension Research</i> , 2019 , 42, 1235-1481	4.7	468
413	Prognosis of "masked" hypertension and "white-coat" hypertension detected by 24-h ambulatory blood pressure monitoring 10-year follow-up from the Ohasama study. <i>Journal of the American College of Cardiology</i> , 2005 , 46, 508-15	15.1	459
412	Prognostic value of reading-to-reading blood pressure variability over 24 hours in 8938 subjects from 11 populations. <i>Hypertension</i> , 2010 , 55, 1049-57	8.5	332
411	Prognostic significance for stroke of a morning pressor surge and a nocturnal blood pressure decline: the Ohasama study. <i>Hypertension</i> , 2006 , 47, 149-54	8.5	327

410	Ambulatory blood pressure and 10-year risk of cardiovascular and noncardiovascular mortality: the Ohasama study. <i>Hypertension</i> , 2005 , 45, 240-5	8.5	320
409	Day-by-day variability of blood pressure and heart rate at home as a novel predictor of prognosis: the Ohasama study. <i>Hypertension</i> , 2008 , 52, 1045-50	8.5	313
408	Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without hypertension: a meta-analysis. <i>Lancet, The</i> , 2012 , 380, 1649-61	4.0	301
407	Prognostic superiority of daytime ambulatory over conventional blood pressure in four populations: a meta-analysis of 7,030 individuals. <i>Journal of Hypertension</i> , 2007 , 25, 1554-64	1.9	266
406	Prognostic value of isolated nocturnal hypertension on ambulatory measurement in 8711 individuals from 10 populations. <i>Journal of Hypertension</i> , 2010 , 28, 2036-45	1.9	263
405	Short- and long-term incidence of stroke in white-coat hypertension. <i>Hypertension</i> , 2005 , 45, 203-8	8.5	230
404	Diagnostic thresholds for ambulatory blood pressure monitoring based on 10-year cardiovascular risk. <i>Circulation</i> , 2007 , 115, 2145-52	16.7	229
403	Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. <i>Nature Genetics</i> , 2015 , 47, 1282-1293	36.3	223
402	Brachial-Ankle Pulse Wave Velocity and the Risk Prediction of Cardiovascular Disease: An Individual Participant Data Meta-Analysis. <i>Hypertension</i> , 2017 , 69, 1045-1052	8.5	220
401	Prognostic value of the morning blood pressure surge in 5645 subjects from 8 populations. <i>Hypertension</i> , 2010 , 55, 1040-8	8.5	205
400	Prognosis of white-coat and masked hypertension: International Database of HOme blood pressure in relation to Cardiovascular Outcome. <i>Hypertension</i> , 2014 , 63, 675-82	8.5	204
399	How many times should blood pressure be measured at home for better prediction of stroke risk? Ten-year follow-up results from the Ohasama study. <i>Journal of Hypertension</i> , 2004 , 22, 1099-104	1.9	201
398	Prediction of mortality by ambulatory blood pressure monitoring versus screening blood pressure measurements: a pilot study in Ohasama. <i>Journal of Hypertension</i> , 1997 , 15, 357-64	1.9	186
397	Prediction of stroke by ambulatory blood pressure monitoring versus screening blood pressure measurements in a general population: the Ohasama study. <i>Journal of Hypertension</i> , 2000 , 18, 847-54	1.9	175
396	Kidney dysfunction as a risk factor for first symptomatic stroke events in a general Japanese population--the Ohasama study. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 1910-5	4.3	155
395	Significance of white-coat hypertension in older persons with isolated systolic hypertension: a meta-analysis using the International Database on Ambulatory Blood Pressure Monitoring in Relation to Cardiovascular Outcomes population. <i>Hypertension</i> , 2012 , 59, 564-71	8.5	146
394	Reference values for 24-hour ambulatory blood pressure monitoring based on a prognostic criterion: the Ohasama Study. <i>Hypertension</i> , 1998 , 32, 255-9	8.5	145
393	Meta-analysis of genome-wide association studies in East Asian-ancestry populations identifies four new loci for body mass index. <i>Human Molecular Genetics</i> , 2014 , 23, 5492-504	5.6	141

392	Association of Office and Ambulatory Blood Pressure With Mortality and Cardiovascular Outcomes. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 322, 409-420	27.4	136
391	The Japanese Society of Hypertension Guidelines for Self-monitoring of Blood Pressure at Home (Second Edition). <i>Hypertension Research</i> , 2012 , 35, 777-95	4.7	135
390	Epidemiology of hypertension in Japan: where are we now?. <i>Circulation Journal</i> , 2013 , 77, 2226-31	2.9	130
389	Global Cardiovascular and Renal Outcomes of Reduced GFR. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 2167-2179	12.7	127
388	Cardiovascular outcomes in the first trial of antihypertensive therapy guided by self-measured home blood pressure. <i>Hypertension Research</i> , 2012 , 35, 1102-10	4.7	126
387	Prediction of stroke by home "morning" versus "evening" blood pressure values: the Ohasama study. <i>Hypertension</i> , 2006 , 48, 737-43	8.5	124
386	Genome-wide association study of coronary artery disease in the Japanese. <i>European Journal of Human Genetics</i> , 2012 , 20, 333-40	5.3	123
385	Added predictive value of night-time blood pressure variability for cardiovascular events and mortality: the Ambulatory Blood Pressure-International Study. <i>Hypertension</i> , 2014 , 64, 487-93	8.5	121
384	Validation of a food-frequency questionnaire for cohort studies in rural Japan. <i>Public Health Nutrition</i> , 2003 , 6, 147-57	3.3	120
383	Ambulatory arterial stiffness index and 24-hour ambulatory pulse pressure as predictors of mortality in Ohasama, Japan. <i>Stroke</i> , 2007 , 38, 1161-6	6.7	115
382	Masked hypertension in diabetes mellitus: treatment implications for clinical practice. <i>Hypertension</i> , 2013 , 61, 964-71	8.5	114
381	White-coat hypertension as a risk factor for the development of home hypertension: the Ohasama study. <i>Archives of Internal Medicine</i> , 2005 , 165, 1541-6		114
380	The International Database of Ambulatory Blood Pressure in relation to Cardiovascular Outcome (IDACO): protocol and research perspectives. <i>Blood Pressure Monitoring</i> , 2007 , 12, 255-62	1.3	111
379	Setting thresholds to varying blood pressure monitoring intervals differentially affects risk estimates associated with white-coat and masked hypertension in the population. <i>Hypertension</i> , 2014 , 64, 935-42	8.5	108
378	Prognosis of isolated systolic and isolated diastolic hypertension as assessed by self-measurement of blood pressure at home: the Ohasama study. <i>Archives of Internal Medicine</i> , 2000 , 160, 3301-6		108
377	Prediction of stroke by self-measurement of blood pressure at home versus casual screening blood pressure measurement in relation to the Joint National Committee 7 classification: the Ohasama study. <i>Stroke</i> , 2004 , 35, 2356-61	6.7	102
376	Control of blood pressure as measured at home and office, and comparison with physicians' assessment of control among treated hypertensive patients in Japan: First Report of the Japan Home versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Hypertension Research</i> , 2004 , 27, 755-63	4.7	100
375	The Cardiovascular Risk of White-Coat Hypertension. <i>Journal of the American College of Cardiology</i> , 2016 , 68, 2033-2043	15.1	96

374	Home blood pressure variability as cardiovascular risk factor in the population of Ohasama. <i>Hypertension</i> , 2013 , 61, 61-9	8.5	95
373	Methodology and technology for peripheral and central blood pressure and blood pressure variability measurement: current status and future directions - Position statement of the European Society of Hypertension Working Group on blood pressure monitoring and cardiovascular variability. <i>Journal of Hypertension</i> , 2016 , 34, 1665-77	1.9	89
372	Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. <i>Hypertension Research</i> , 2012 , 35, 947-53	4.7	87
371	Isolated uncontrolled hypertension at home and in the office among treated hypertensive patients from the J-HOME study. <i>Journal of Hypertension</i> , 2005 , 23, 1653-60	1.9	84
370	Characteristics of blood pressure measured at home in the morning and in the evening: the Ohasama study. <i>Journal of Hypertension</i> , 1999 , 17, 889-98	1.9	84
369	Ambulatory blood pressure monitoring in 9357 subjects from 11 populations highlights missed opportunities for cardiovascular prevention in women. <i>Hypertension</i> , 2011 , 57, 397-405	8.5	80
368	Device for the self-measurement of blood pressure that can monitor blood pressure during sleep. <i>Blood Pressure Monitoring</i> , 2001 , 6, 203-5	1.3	80
367	A prospective cohort study on National Health Insurance beneficiaries in Ohsaki, Miyagi Prefecture, Japan: study design, profiles of the subjects and medical cost during the first year. <i>Journal of Epidemiology</i> , 1998 , 8, 258-63	3.4	78
366	Stool color card screening for early detection of biliary atresia and long-term native liver survival: a 19-year cohort study in Japan. <i>Journal of Pediatrics</i> , 2015 , 166, 897-902.e1	3.6	76
365	Blood pressure variability assessed by home measurements: a systematic review. <i>Hypertension Research</i> , 2014 , 37, 565-72	4.7	76
364	Outcome-driven thresholds for home blood pressure measurement: international database of home blood pressure in relation to cardiovascular outcome. <i>Hypertension</i> , 2013 , 61, 27-34	8.5	75
363	Detection of carotid atherosclerosis in individuals with masked hypertension and white-coat hypertension by self-measured blood pressure at home: the Ohasama study. <i>Journal of Hypertension</i> , 2007 , 25, 321-7	1.9	75
362	Usefulness of home blood pressure measurements in assessing the effect of treatment in a single-blind placebo-controlled open trial. <i>Journal of Hypertension</i> , 2001 , 19, 179-85	1.9	74
361	Seasonal variation in blood pressure in normotensive women studied by home measurements. <i>Clinical Science</i> , 1996 , 90, 55-60	6.5	74
360	Prognostic value of ambulatory heart rate revisited in 6928 subjects from 6 populations. <i>Hypertension</i> , 2008 , 52, 229-35	8.5	73
359	Conformational change in transfer RNA is an early indicator of acute cellular damage. <i>Journal of the American Society of Nephrology: JASN</i> , 2014 , 25, 2316-26	12.7	70
358	Cigarette smoking, systolic blood pressure, and cardiovascular diseases in the Asia-Pacific region. <i>Stroke</i> , 2008 , 39, 1694-702	6.7	70
357	Prognostic value of home heart rate for cardiovascular mortality in the general populationThe Ohasama study. <i>American Journal of Hypertension</i> , 2004 , 17, 1005-1010	2.3	69

356	Blood pressure control assessed by home, ambulatory and conventional blood pressure measurements in the Japanese general population: the Ohasama study. <i>Hypertension Research</i> , 2002 , 25, 57-63	4.7	66
355	Hypertension and related diseases in the era of COVID-19: a report from the Japanese Society of Hypertension Task Force on COVID-19. <i>Hypertension Research</i> , 2020 , 43, 1028-1046	4.7	65
354	The reason why home blood pressure measurements are preferred over clinic or ambulatory blood pressure in Japan. <i>Hypertension Research</i> , 2013 , 36, 661-72	4.7	64
353	Prognostic value of home heart rate for cardiovascular mortality in the general population: the Ohasama study. <i>American Journal of Hypertension</i> , 2004 , 17, 1005-10	2.3	64
352	Hypertension types defined by clinic and ambulatory blood pressure in 14 143 patients referred to hypertension clinics worldwide. Data from the ARTEMIS study. <i>Journal of Hypertension</i> , 2016 , 34, 2187-98	1.9	64
351	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCH-JAPAN study. <i>Journal of Clinical Lipidology</i> , 2018 , 12, 674-684.e5	4.9	61
350	Ambulatory blood pressure, blood pressure variability and the prevalence of carotid artery alteration: the Ohasama study. <i>Journal of Hypertension</i> , 2007 , 25, 1704-10	1.9	61
349	Characteristics of resistant hypertension determined by self-measured blood pressure at home and office blood pressure measurements: the J-HOME study. <i>Journal of Hypertension</i> , 2006 , 24, 1737-43	1.9	61
348	Age-specific differences between conventional and ambulatory daytime blood pressure values. <i>Hypertension</i> , 2014 , 64, 1073-9	8.5	60
347	Predictive value of night-time heart rate for cardiovascular events in hypertension. The ABP-International study. <i>International Journal of Cardiology</i> , 2013 , 168, 1490-5	3.2	60
346	Impact of metabolic syndrome on the risk of cardiovascular disease mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014 , 113, 84-9	3	60
345	Genome-wide association study meta-analysis reveals transethnic replication of mean arterial and pulse pressure loci. <i>Hypertension</i> , 2013 , 62, 853-9	8.5	60
344	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population: NIPPON DATA2010. <i>Journal of Hypertension</i> , 2017 , 35, 401-408	1.9	59
343	Long-term risk of BP values above normal for cardiovascular mortality: a 24-year observation of Japanese aged 30 to 92 years. <i>Journal of Hypertension</i> , 2012 , 30, 2299-306	1.9	59
342	Predictive value of ambulatory heart rate in the Japanese general population: the Ohasama study. <i>Journal of Hypertension</i> , 2008 , 26, 1571-6	1.9	59
341	Ambulatory hypertension subtypes and 24-hour systolic and diastolic blood pressure as distinct outcome predictors in 8341 untreated people recruited from 12 populations. <i>Circulation</i> , 2014 , 130, 466-74	16.7	58
340	Risk stratification by self-measured home blood pressure across categories of conventional blood pressure: a participant-level meta-analysis. <i>PLoS Medicine</i> , 2014 , 11, e1001591	11.6	57
339	Long-Term Stroke Risk Due to Partial White-Coat or Masked Hypertension Based on Home and Ambulatory Blood Pressure Measurements: The Ohasama Study. <i>Hypertension</i> , 2016 , 67, 48-55	8.5	55

338	Ambulatory versus home versus clinic blood pressure: the association with subclinical cerebrovascular diseases: the Ohasama Study. <i>Hypertension</i> , 2012 , 59, 22-8	8.5	54
337	Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA80 cohort study. <i>BMJ Open</i> , 2016 , 6, e011632	3	53
336	Insufficient duration of action of antihypertensive drugs mediates high blood pressure in the morning in hypertensive population: the Ohasama study. <i>Clinical and Experimental Hypertension</i> , 2002 , 24, 261-75	2.2	53
335	Development of Risk Prediction Equations for Incident Chronic Kidney Disease. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 322, 2104-2114	27.4	52
334	Stroke risk and antihypertensive drug treatment in the general population: the Japan arteriosclerosis longitudinal study. <i>Journal of Hypertension</i> , 2009 , 27, 357-64	1.9	51
333	Prevalence of masked uncontrolled and treated white-coat hypertension defined according to the average of morning and evening home blood pressure value: from the Japan Home versus Office Measurement Evaluation Study. <i>Blood Pressure Monitoring</i> , 2005 , 10, 311-6	1.3	51
332	Day-to-day variability in home blood pressure is associated with cognitive decline: the Ohasama study. <i>Hypertension</i> , 2014 , 63, 1333-8	8.5	50
331	Prevalence, Treatment, and Control Rates of Conventional and Ambulatory Hypertension Across 10 Populations in 3 Continents. <i>Hypertension</i> , 2017 , 70, 50-58	8.5	49
330	Combined effects of maternal age and parity on successful initiation of exclusive breastfeeding. <i>Preventive Medicine Reports</i> , 2016 , 3, 121-6	2.6	49
329	Factors associated with day-by-day variability of self-measured blood pressure at home: the Ohasama study. <i>American Journal of Hypertension</i> , 2010 , 23, 980-6	2.3	49
328	Use of 2003 European Society of Hypertension-European Society of Cardiology guidelines for predicting stroke using self-measured blood pressure at home: the Ohasama study. <i>European Heart Journal</i> , 2005 , 26, 2026-31	9.5	49
327	Ambulatory blood pressure monitoring in evaluating the prevalence of hypertension in adults in Ohasama, a rural Japanese community. <i>Hypertension Research</i> , 1996 , 19, 207-12	4.7	49
326	Cost-effectiveness of the introduction of home blood pressure measurement in patients with office hypertension. <i>Journal of Hypertension</i> , 2008 , 26, 685-90	1.9	48
325	Association of arterial stiffness with silent cerebrovascular lesions: the Ohasama study. <i>Cerebrovascular Diseases</i> , 2011 , 31, 329-37	3.2	47
324	Seasonal trends of blood pressure during pregnancy in Japan: the babies and their parentsQ longitudinal observation in Suzuki Memorial Hospital in Intrauterine Period study. <i>Journal of Hypertension</i> , 2008 , 26, 2406-13	1.9	47
323	Fruit and vegetable consumption and the risk of hypertension determined by self measurement of blood pressure at home: the Ohasama study. <i>Hypertension Research</i> , 2008 , 31, 1435-43	4.7	45
322	Prediction of ischaemic and haemorrhagic stroke by self-measured blood pressure at home: the Ohasama study. <i>Blood Pressure Monitoring</i> , 2004 , 9, 315-20	1.3	45
321	Nocturnal blood pressure measured by home devices: evidence and perspective for clinical application. <i>Journal of Hypertension</i> , 2019 , 37, 905-916	1.9	45

320	Outcome-Driven Thresholds for Increased Home Blood Pressure Variability. <i>Hypertension</i> , 2017 , 69, 599-607	4.7	44
319	Hypertension with diabetes mellitus: significance from an epidemiological perspective for Japanese. <i>Hypertension Research</i> , 2017 , 40, 795-806	4.7	44
318	Reproducibility of nocturnal blood pressure assessed by self-measurement of blood pressure at home. <i>Hypertension Research</i> , 2007 , 30, 707-12	4.7	44
317	Genome-wide association meta-analysis identifies novel variants associated with fasting plasma glucose in East Asians. <i>Diabetes</i> , 2015 , 64, 291-8	0.9	43
316	Six random specimens of daytime casual urine on different days are sufficient to estimate daily sodium/potassium ratio in comparison to 7-day 24-h urine collections. <i>Hypertension Research</i> , 2014 , 37, 765-71	4.7	43
315	The economic impact of the introduction of home blood pressure measurement for the diagnosis and treatment of hypertension. <i>Blood Pressure Monitoring</i> , 2006 , 11, 257-67	1.3	43
314	Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: a 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , 2014 , 232, 384-9	3.1	42
313	HbA1c and the risks for all-cause and cardiovascular mortality in the general Japanese population: NIPPON DATA90. <i>Diabetes Care</i> , 2013 , 36, 3759-65	14.6	42
312	Prognostic significance of variability in ambulatory and home blood pressure from the Ohasama study. <i>Journal of Epidemiology</i> , 2007 , 17, 109-13	3.4	42
311	Thirty years of research on diagnostic and therapeutic thresholds for the self-measured blood pressure at home. <i>Blood Pressure Monitoring</i> , 2008 , 13, 352-65	1.3	41
310	Glomerular hyperfiltration is a predictor of adverse cardiovascular outcomes. <i>Kidney International</i> , 2018 , 93, 195-203	9.9	40
309	Cardiovascular risk with and without antihypertensive drug treatment in the Japanese general population: participant-level meta-analysis. <i>Hypertension</i> , 2014 , 63, 1189-97	8.5	40
308	Stroke risk in treated hypertension based on home blood pressure: the Ohasama study. <i>American Journal of Hypertension</i> , 2010 , 23, 508-14	2.3	40
307	Plasma fibrinogen, ambulatory blood pressure, and silent cerebrovascular lesions: the Ohasama study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 963-8	9.4	39
306	Medical cost for disability: a longitudinal observation of national health insurance beneficiaries in Japan. <i>Journal of the American Geriatrics Society</i> , 1999 , 47, 470-6	5.6	39
305	Risk stratification by ambulatory blood pressure monitoring across JNC classes of conventional blood pressure. <i>American Journal of Hypertension</i> , 2014 , 27, 956-65	2.3	37
304	Association of environmental tobacco smoke exposure with elevated home blood pressure in Japanese women: the Ohasama study. <i>Journal of Hypertension</i> , 2010 , 28, 1814-20	1.9	37
303	Predicting stroke using 4 ambulatory blood pressure monitoring-derived blood pressure indices: the Ohasama Study. <i>Hypertension</i> , 2006 , 48, 877-82	8.5	37

302	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016 , 246, 141-7	3.1	36
301	Home blood pressure monitoring in the 21st century. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 1116-1121	3	36
300	Emergence of Home Blood Pressure-Guided Management of Hypertension Based on Global Evidence. <i>Hypertension</i> , 2019 , HYPERTENSIONAHA11912630	8.5	36
299	Masked hypertension determined by self-measured blood pressure at home and chronic kidney disease in the Japanese general population: the Ohasama study. <i>Hypertension Research</i> , 2008 , 31, 2129-35	4.7	36
298	Therapeutic effects of evening administration of guanabenz and clonidine on morning hypertension: evaluation using home-based blood pressure measurements. <i>Journal of Hypertension</i> , 2003 , 21, 805-11	1.9	36
297	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019 , 24, 1	4.2	36
296	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014 , 112, 916-24	3.6	35
295	Carotid intima-media thickness and plaque in apparently healthy Japanese individuals with an estimated 10-year absolute risk of CAD death according to the Japan Atherosclerosis Society (JAS) guidelines 2012: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 755-66	4	35
294	Pre-hypertension as a significant predictor of chronic kidney disease in a general population: the Ohasama Study. <i>Nephrology Dialysis Transplantation</i> , 2012 , 27, 3218-23	4.3	35
293	Efficacy and duration of action of the four selective angiotensin II subtype 1 receptor blockers, losartan, candesartan, valsartan and telmisartan, in patients with essential hypertension determined by home blood pressure measurements. <i>Clinical and Experimental Hypertension</i> , 2005 , 27, 477-89	2.2	35
292	Predictive power of home blood pressure and clinic blood pressure in hypertensive patients with impaired glucose metabolism and diabetes. <i>Journal of Hypertension</i> , 2013 , 31, 1593-602	1.9	34
291	Usefulness of assessing masked and white-coat hypertension by ambulatory blood pressure monitoring for determining prevalent risk of chronic kidney disease: the Ohasama study. <i>Hypertension Research</i> , 2010 , 33, 1192-8	4.7	34
290	Factors affecting the difference between screening and home blood pressure measurements: the Ohasama Study. <i>Journal of Hypertension</i> , 2001 , 19, 13-9	1.9	34
289	Association Between Blood Pressure Variability and Cerebral Small-Vessel Disease: A Systematic Review and Meta-Analysis. <i>Journal of the American Heart Association</i> , 2020 , 9, e013841	6	34
288	Association between tooth loss and cognitive impairment in community-dwelling older Japanese adults: a 4-year prospective cohort study from the Ohasama study. <i>BMC Oral Health</i> , 2018 , 18, 142	3.7	33
287	Predictive values of automated blood pressure measurement: what can we learn from the Japanese population - the Ohasama study. <i>Blood Pressure Monitoring</i> , 2001 , 6, 335-9	1.3	33
286	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 800-9	4	33
285	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980-2010. <i>Hypertension Research</i> , 2015 , 38, 790-5	4.7	32

284	Relationship between health practices and education level in the rural Japanese population. <i>Journal of Epidemiology</i> , 2000 , 10, 149-56	3.4	32
283	Combined effect of blood pressure and total cholesterol levels on long-term risks of subtypes of cardiovascular death: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan. <i>Hypertension</i> , 2015 , 65, 517-24	8.5	31
282	Blood pressure measured in the clinic and at home during pregnancy among nulliparous and multiparous women: the BOSHI study. <i>American Journal of Hypertension</i> , 2013 , 26, 141-8	2.3	31
281	Participation in health check-ups and mortality using propensity score matched cohort analyses. <i>Preventive Medicine</i> , 2010 , 51, 397-402	4.3	31
280	The association between masked hypertension and waist circumference as an obesity-related anthropometric index for metabolic syndrome: the Ohasama study. <i>Hypertension Research</i> , 2009 , 32, 438-43	4.7	31
279	Effects of exercise training on home blood pressure values in older adults: a randomized controlled trial. <i>Journal of Hypertension</i> , 2001 , 19, 1045-52	1.9	31
278	Association Between Amplitude of Seasonal Variation in Self-Measured Home Blood Pressure and Cardiovascular Outcomes: HOMED-BP (Hypertension Objective Treatment Based on Measurement By Electrical Devices of Blood Pressure) Study. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	30
277	How many measurements are needed to estimate blood pressure variability without loss of prognostic information?. <i>American Journal of Hypertension</i> , 2014 , 27, 46-55	2.3	30
276	Outcome-driven thresholds for ambulatory pulse pressure in 9938 participants recruited from 11 populations. <i>Hypertension</i> , 2014 , 63, 229-37	8.5	30
275	Blood pressure load does not add to ambulatory blood pressure level for cardiovascular risk stratification. <i>Hypertension</i> , 2014 , 63, 925-33	8.5	30
274	Association of kidney dysfunction with silent lacunar infarcts and white matter hyperintensity in the general population: the Ohasama study. <i>Cerebrovascular Diseases</i> , 2010 , 30, 43-50	3.2	30
273	Practice and awareness of physicians regarding home blood pressure measurement in Japan. <i>Hypertension Research</i> , 2010 , 33, 428-34	4.7	30
272	Accumulation of common polymorphisms is associated with development of hypertension: a 12-year follow-up from the Ohasama study. <i>Hypertension Research</i> , 2010 , 33, 129-34	4.7	30
271	Optimal number of days for home blood pressure measurement. <i>American Journal of Hypertension</i> , 2015 , 28, 595-603	2.3	29
270	Cross-sectional comparison of coronary artery calcium scores between Caucasian men in the United States and Japanese men in Japan: the multi-ethnic study of atherosclerosis and the Shiga epidemiological study of subclinical atherosclerosis. <i>American Journal of Epidemiology</i> , 2014 , 180, 590-8	3.8	29
269	Night-time blood pressure is associated with the development of chronic kidney disease in a general population: the Ohasama Study. <i>Journal of Hypertension</i> , 2013 , 31, 2410-7	1.9	29
268	Associations between day-by-day variability in blood pressure measured at home and antihypertensive drugs: the J-HOME-Morning study. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 297-304	2.2	29
267	Genome-wide response to antihypertensive medication using home blood pressure measurements: a pilot study nested within the HOMED-BP study. <i>Pharmacogenomics</i> , 2013 , 14, 1709-21	2.6	28

266	Seasonal variation in self-measured home blood pressure among patients on antihypertensive medications: HOMED-BP study. <i>Hypertension Research</i> , 2017 , 40, 284-290	4.7	28
265	Association between J-point elevation and death from coronary artery disease--15-year follow up of the NIPPON DATA90. <i>Circulation Journal</i> , 2013 , 77, 1260-6	2.9	28
264	Practical aspect of monitoring hypertension based on self-measured blood pressure at home. <i>Internal Medicine</i> , 2004 , 43, 771-8	1.1	28
263	Animal protein intake is associated with higher-level functional capacity in elderly adults: the Ohasama study. <i>Journal of the American Geriatrics Society</i> , 2014 , 62, 426-34	5.6	27
262	Breastfeeding leads to lower blood pressure in 7-year-old Japanese children: Tohoku Study of Child Development. <i>Hypertension Research</i> , 2013 , 36, 117-22	4.7	27
261	The relation between health literacy, hypertension knowledge, and blood pressure among middle-aged Japanese adults. <i>Blood Pressure Monitoring</i> , 2011 , 16, 224-30	1.3	27
260	Home blood pressure level, blood pressure variability, smoking, and stroke risk in Japanese men: the Ohasama study. <i>American Journal of Hypertension</i> , 2012 , 25, 883-91	2.3	27
259	Seasonal variation in blood pressure: Evidence, consensus and recommendations for clinical practice. Consensus statement by the European Society of Hypertension Working Group on Blood Pressure Monitoring and Cardiovascular Variability. <i>Journal of Hypertension</i> , 2020 , 38, 1235-1243	1.9	26
258	The International Database of HOme blood pressure in relation to Cardiovascular Outcome (IDHOCO): moving from baseline characteristics to research perspectives. <i>Hypertension Research</i> , 2012 , 35, 1072-9	4.7	26
257	Health behaviors as predictors for declines in higher-level functional capacity in older adults: the Ohasama study. <i>Journal of the American Geriatrics Society</i> , 2011 , 59, 1993-2000	5.6	26
256	Practice and awareness of physicians regarding casual-clinic blood pressure measurement in Japan. <i>Hypertension Research</i> , 2010 , 33, 960-4	4.7	26
255	Stroke risk of blood pressure indices determined by home blood pressure measurement: the Ohasama study. <i>Stroke</i> , 2009 , 40, 2859-61	6.7	26
254	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016 , 5,	6	26
253	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016 , 4, 225-32	2.6	26
252	Defining thresholds for home blood pressure monitoring in octogenarians. <i>Hypertension</i> , 2015 , 66, 865-73	7.5	25
251	Risk of hypercholesterolemia for cardiovascular disease and the population attributable fraction in a 24-year Japanese cohort study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 95-107	4	25
250	Double product reflects the predictive power of systolic pressure in the general population: evidence from 9,937 participants. <i>American Journal of Hypertension</i> , 2013 , 26, 665-72	2.3	25
249	Relationship of Estimated GFR and Albuminuria to Concurrent Laboratory Abnormalities: An Individual Participant Data Meta-analysis in a Global Consortium. <i>American Journal of Kidney Diseases</i> , 2019 , 73, 206-217	7.4	25

248	Seasonal variation in home blood pressure: findings from nationwide web-based monitoring in Japan. <i>BMJ Open</i> , 2018 , 8, e017351	3	24
247	Does Antihypertensive Drug Class Affect Day-to-Day Variability of Self-Measured Home Blood Pressure? The HOMED-BP Study. <i>Journal of the American Heart Association</i> , 2016 , 5, e002995	6	24
246	Plasma renin activity and the aldosterone-to-renin ratio are associated with the development of chronic kidney disease: the Ohasama Study. <i>Journal of Hypertension</i> , 2012 , 30, 1632-8	1.9	24
245	Relationships among blood pressures obtained using different measurement methods in the general population of Ohasama, Japan. <i>Hypertension Research</i> , 1999 , 22, 261-72	4.7	24
244	Risk for metabolic diseases in normal weight individuals with visceral fat accumulation: a cross-sectional study in Japan. <i>BMJ Open</i> , 2017 , 7, e013831	3	23
243	Predictive value for mortality of the double product at rest obtained by home blood pressure measurement: the Ohasama study. <i>American Journal of Hypertension</i> , 2012 , 25, 568-75	2.3	23
242	Long-term outcome of healthy participants with atrial premature complex: a 15-year follow-up of the NIPPON DATA 90 cohort. <i>PLoS ONE</i> , 2013 , 8, e80853	3.7	23
241	Daily serial hemodynamic data during pregnancy and seasonal variation: the BOSHI study. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 290-6	2.2	22
240	Prevalence of hypertension and rate of blood pressure control as assessed by home blood pressure measurements in a rural Japanese community, Ohasama. <i>Clinical and Experimental Hypertension</i> , 1996 , 18, 713-28	2.2	22
239	The fine-scale genetic structure and evolution of the Japanese population. <i>PLoS ONE</i> , 2017 , 12, e0185487	3.7	21
238	Effectiveness of a Self-monitoring Device for Urinary Sodium-to-Potassium Ratio on Dietary Improvement in Free-Living Adults: a Randomized Controlled Trial. <i>Journal of Epidemiology</i> , 2018 , 28, 41-47	3.4	21
237	A comparison of risk factors for mortality from heart failure in Asian and non-Asian populations: an overview of individual participant data from 32 prospective cohorts from the Asia-Pacific Region. <i>BMC Cardiovascular Disorders</i> , 2014 , 14, 61	2.3	21
236	Are blood pressure and diabetes additive or synergistic risk factors? Outcome in 8494 subjects randomly recruited from 10 populations. <i>Hypertension Research</i> , 2011 , 34, 714-21	4.7	21
235	Prognostic values of clockwise and counterclockwise rotation for cardiovascular mortality in Japanese subjects: a 24-year follow-up of the National Integrated Project for Prospective Observation of Noncommunicable Disease and Its Trends in the Aged, 1980-2004 (NIPPON DATA80). <i>Circulation</i> , 2012 , 125, 1226-33	16.7	21
234	Income-related inequalities in preventive and curative dental care use among working-age Japanese adults in urban areas: a cross-sectional study. <i>BMC Oral Health</i> , 2014 , 14, 117	3.7	20
233	The National Integrated Project for Prospective Observation of Non-communicable Disease and its Trends in the Aged 2010 (NIPPON DATA2010): Objectives, Design, and Population Characteristics. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S2-S9	3.4	20
232	A prospective study of the impact of diabetes mellitus on restrictive and obstructive lung function impairment: The Saku study. <i>Metabolism: Clinical and Experimental</i> , 2018 , 82, 58-64	12.7	19
231	Influence of alcohol intake on circadian blood pressure variation in Japanese men: the Ohasama study. <i>American Journal of Hypertension</i> , 2009 , 22, 1171-6	2.3	19

230	Aldosterone-to-renin ratio as a predictor of stroke under conditions of high sodium intake: the Ohasama study. <i>American Journal of Hypertension</i> , 2012 , 25, 777-83	2.3	19
229	How many times should we ask subjects to measure blood pressure at home on each occasion?. <i>Journal of Hypertension</i> , 2007 , 25, 1987-91	1.9	19
228	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level: EPOCH-JAPAN (Evidence for Cardiovascular Prevention From Observational Cohorts in Japan). <i>Hypertension</i> , 2019 , 73, 52-59	8.5	19
227	Relationship Between Socioeconomic Status and the Prevalence of Underweight, Overweight or Obesity in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S10-S16	3.4	19
226	Proposed Cutoff Value of Brachial-Ankle Pulse Wave Velocity for the Management of Hypertension. <i>Circulation Journal</i> , 2017 , 81, 1540-1542	2.9	18
225	Reference values and associated factors for Japanese newborns blood pressure and pulse rate: the babies and their parents longitudinal observation in Suzuki Memorial Hospital on intrauterine period (BOSHI) study. <i>Journal of Hypertension</i> , 2016 , 34, 1578-85	1.9	18
224	Patterns of ambulatory blood pressure: clinical relevance and application. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 1112-1115	2.3	18
223	Attributable risk fraction of prehypertension on cardiovascular disease mortality in the Japanese population: the Ohsaki Study. <i>American Journal of Hypertension</i> , 2009 , 22, 267-72	2.3	18
222	Does body mass index impact on the relationship between systolic blood pressure and cardiovascular disease?: meta-analysis of 419 488 individuals from the Asia pacific cohort studies collaboration. <i>Stroke</i> , 2012 , 43, 1478-83	6.7	18
221	Ankle-brachial index measured by oscillometry is predictive for cardiovascular disease and premature death in the Japanese population: An individual participant data meta-analysis. <i>Atherosclerosis</i> , 2018 , 275, 141-148	3.1	18
220	Masked tachycardia. A predictor of adverse outcome in hypertension. <i>Journal of Hypertension</i> , 2017 , 35, 487-492	1.9	17
219	Simultaneously Measured Interarm Blood Pressure Difference and Stroke: An Individual Participants Data Meta-Analysis. <i>Hypertension</i> , 2018 , 71, 1030-1038	8.5	17
218	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014 , 236, 237-43	3.1	17
217	Associations between Inflammatory Markers and Subclinical Atherosclerosis in Middle-aged White, Japanese-American and Japanese Men: The ERA-JUMP Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 590-8	4	17
216	Association between Pulse Wave Velocity and Coronary Artery Calcification in Japanese men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 1266-77	4	17
215	Risk stratification by 24-hour ambulatory blood pressure and estimated glomerular filtration rate in 5322 subjects from 11 populations. <i>Hypertension</i> , 2013 , 61, 18-26	8.5	17
214	Aldosterone-to-renin ratio and nocturnal blood pressure decline in a general population: the Ohasama study. <i>Journal of Hypertension</i> , 2011 , 29, 1940-7	1.9	17
213	Detection of silent cerebrovascular lesions in individuals with masked and white-coat hypertension by home blood pressure measurement: the Ohasama study. <i>Journal of Hypertension</i> , 2009 , 27, 1049-55	1.9	17

212	Factors affecting the nocturnal decrease in blood pressure: a community-based study in Ohasama. <i>Journal of Hypertension</i> , 1997 , 15, 827-38	1.9	17
211	The value of self-measured home blood pressure in predicting stroke. <i>Expert Review of Neurotherapeutics</i> , 2006 , 6, 163-73	4.3	17
210	The current status of home and office blood pressure control among hypertensive patients with diabetes mellitus: the Japan Home Versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Diabetes Research and Clinical Practice</i> , 2006 , 73, 276-83	7.4	17
209	Evidence-based proposal for the number of ambulatory readings required for assessing blood pressure level in research settings: an analysis of the IDACO database. <i>Blood Pressure</i> , 2018 , 27, 341-350	1.7	17
208	Are the cardiovascular outcomes of participants with white-coat hypertension poor compared to those of participants with normotension? A systemic review and meta-analysis. <i>Hypertension Research</i> , 2019 , 42, 825-833	4.7	16
207	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016 , 220, 262-7	3.2	16
206	Thresholds for conventional and home blood pressure by sex and age in 5018 participants from 5 populations. <i>Hypertension</i> , 2014 , 64, 695-701	8.5	16
205	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015 , 239, 444-50	3.1	16
204	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and U.S. whites. <i>European Heart Journal Cardiovascular Imaging</i> , 2013 , 14, 921-7	4.1	16
203	Uncontrolled hypertension based on morning and evening home blood pressure measurements from the J-HOME study. <i>Hypertension Research</i> , 2009 , 32, 1072-8	4.7	16
202	Aldosterone-to-renin ratio and home blood pressure in subjects with higher and lower sodium intake: the Ohasama study. <i>Hypertension Research</i> , 2011 , 34, 361-6	4.7	16
201	Repeated evening home blood pressure measurement improves prognostic significance for stroke: a 12-year follow-up of the Ohasama study. <i>Blood Pressure Monitoring</i> , 2009 , 14, 93-8	1.3	16
200	Association between alcohol consumption and incidence of impaired insulin secretion and insulin resistance in Japanese: The Saku study. <i>Diabetes Research and Clinical Practice</i> , 2018 , 135, 11-17	7.4	16
199	Characterising private and shared signatures of positive selection in 37 Asian populations. <i>European Journal of Human Genetics</i> , 2017 , 25, 499-508	5.3	15
198	Factors Associated with Pneumonia-caused Death in Older Adults with Autopsy-confirmed Dementia. <i>Internal Medicine</i> , 2017 , 56, 907-914	1.1	15
197	Distinct association between educational attainment and overweight/obesity in unmarried and married women: evidence from a population-based study in Japan. <i>BMC Public Health</i> , 2017 , 17, 903	4.1	15
196	Risk Factors for Stroke among Young-Old and Old-Old Community-Dwelling Adults in Japan: The Ohasama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017 , 24, 290-300	4	15
195	Maternal clinic and home blood pressure measurements during pregnancy and infant birth weight: the BOSHI study. <i>Hypertension Research</i> , 2016 , 39, 151-7	4.7	15

194	The velocity of antihypertensive effect of losartan/hydrochlorothiazide and angiotensin II receptor blocker. <i>Journal of Hypertension</i> , 2012 , 30, 1478-86	1.9	15
193	Prognostic significance of morning surge in blood pressure: which definition, which outcome?. <i>Blood Pressure Monitoring</i> , 2008 , 13, 161-2	1.3	15
192	Home blood pressure monitoring: methodology, clinical relevance and practical application: a 2021 position paper by the Working Group on Blood Pressure Monitoring and Cardiovascular Variability of the European Society of Hypertension. <i>Journal of Hypertension</i> , 2021 , 39, 1742-1767	1.9	15
191	Parity as a factor affecting the white-coat effect in pregnant women: the BOSHI study. <i>Hypertension Research</i> , 2015 , 38, 770-5	4.7	14
190	Aldosterone-to-renin ratio and nocturnal blood pressure decline assessed by self-measurement of blood pressure at home: the Ohasama Study. <i>Clinical and Experimental Hypertension</i> , 2014 , 36, 108-14	2.2	14
189	Proposal of a risk-stratification system for the Japanese population based on blood pressure levels: the Ohasama study. <i>Hypertension Research</i> , 2008 , 31, 1315-22	4.7	14
188	Relationship between office and home blood pressure with increasing age: The International Database of HOME blood pressure in relation to Cardiovascular Outcome (IDHOCO). <i>Hypertension Research</i> , 2016 , 39, 612-7	4.7	14
187	Epidemiology of hypertension in Japan: beyond the new 2019 Japanese guidelines. <i>Hypertension Research</i> , 2020 , 43, 1344-1351	4.7	13
186	Impaired Higher-Level Functional Capacity as a Predictor of Stroke in Community-Dwelling Older Adults: The Ohasama Study. <i>Stroke</i> , 2016 , 47, 323-8	6.7	13
185	Outcome-Driven Thresholds for Ambulatory Blood Pressure Based on the New American College of Cardiology/American Heart Association Classification of Hypertension. <i>Hypertension</i> , 2019 , 74, 776-783	8.5	13
184	Opposing Age-Related Trends in Absolute and Relative Risk of Adverse Health Outcomes Associated With Out-of-Office Blood Pressure. <i>Hypertension</i> , 2019 , 74, 1333-1342	8.5	13
183	Reference frame for home pulse pressure based on cardiovascular risk in 6470 subjects from 5 populations. <i>Hypertension Research</i> , 2014 , 37, 672-8	4.7	13
182	Introversion associated with large differences between screening blood pressure and home blood pressure measurement: The Ohasama study. <i>Journal of Hypertension</i> , 2006 , 24, 2183-9	1.9	13
181	Hyperuricemia predicts the risk for developing hypertension independent of alcohol drinking status in men and women: the Saku study. <i>Hypertension Research</i> , 2020 , 43, 442-449	4.7	13
180	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , 2016 , 213, 460-467	9.3	13
179	Home blood pressure variability and subclinical atherosclerosis in multiple vascular beds: a population-based study. <i>Journal of Hypertension</i> , 2018 , 36, 2193-2203	1.9	13
178	Comparison of blood pressure values-self-measured at home, measured at an unattended office, and measured at a conventional attended office. <i>Hypertension Research</i> , 2019 , 42, 1726-1737	4.7	12
177	Cumulative impact of axial, structural, and repolarization ECG findings on long-term cardiovascular mortality among healthy individuals in Japan: National Integrated Project for Prospective Observation of Non-Communicable Disease and its Trends in the Aged, 1980 and 1990. <i>European Journal of Preventive Cardiology</i> , 2014 , 21, 1501-8	3.9	12

176	Blood pressure, heart rate, and double product in a pooled cohort: the Japan Arteriosclerosis Longitudinal Study. <i>Journal of Hypertension</i> , 2017 , 35, 1808-1815	1.9	12
175	Association of Extreme Nocturnal Dipping With Cardiovascular Events Strongly Depends on Age. <i>Hypertension</i> , 2020 , 75, 324-330	8.5	12
174	Association of Total Energy Intake with 29-Year Mortality in the Japanese: NIPPON DATA80. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 339-54	4	12
173	Influence of pneumonia complications on the prognosis of patients with autopsy-confirmed Alzheimer's disease, dementia with Lewy bodies, and vascular dementia. <i>Psychogeriatrics</i> , 2016 , 16, 305-14	1.8	12
172	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 198-206	4	12
171	The effects of increasing calcium channel blocker dose vs. adding a diuretic to treatment regimens for patients with uncontrolled hypertension. <i>Hypertension Research</i> , 2017 , 40, 892-898	4.7	11
170	Validation of a wrist-type home nocturnal blood pressure monitor in the sitting and supine position according to the ANSI/AAMI/ISO81060-2:2013 guidelines: Omron HEM-9601T. <i>Journal of Clinical Hypertension</i> , 2020 , 22, 970-978	2.3	11
169	Association between N-terminal pro B-type natriuretic peptide and day-to-day blood pressure and heart rate variability in a general population: the Ohasama study. <i>Journal of Hypertension</i> , 2015 , 33, 1536-41	1.9	11
168	Relationship between maternal gestational hypertension and home blood pressure in 7-year-old children and their mothers: Tohoku Study of Child Development. <i>Hypertension Research</i> , 2015 , 38, 776-82	4.7	11
167	Difference between home and office blood pressures among treated hypertensive patients from the Japan Home versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Hypertension Research</i> , 2008 , 31, 1115-23	4.7	11
166	Day-to-day blood pressure variability is associated with lower cognitive performance among the Japanese community-dwelling oldest-old population: the SONIC study. <i>Hypertension Research</i> , 2020 , 43, 404-411	4.7	11
165	From mercury sphygmomanometer to electric device on blood pressure measurement: correspondence of Minamata Convention on Mercury. <i>Hypertension Research</i> , 2016 , 39, 179-82	4.7	11
164	Development and evaluation of a home nocturnal blood pressure monitoring system using a wrist-cuff device. <i>Blood Pressure Monitoring</i> , 2018 , 23, 318-326	1.3	11
163	Macronutrient Intake and Socioeconomic Status: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S17-S22	3.4	11
162	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S29-S34	3.4	11
161	Awareness of the Japanese Society of Hypertension Guidelines for the Management of Hypertension and their use in clinical practices: 2009 survey results. <i>Hypertension Research</i> , 2015 , 38, 400-4	4.7	10
160	The effect of fathers' and mothers' educational level on adult oral health in Japan. <i>Community Dentistry and Oral Epidemiology</i> , 2016 , 44, 283-91	2.8	10
159	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia LOX-1 ligand and IMT in Japanese. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 172-80.e1	4.9	10

158	High central blood pressure is associated with incident cardiovascular events in treated hypertensives: the ABC-J II Study. <i>Hypertension Research</i> , 2018 , 41, 947-956	4.7	10
157	Prognostic values of bundle branch blocks for cardiovascular mortality in Japanese (24year follow-up of NIPPON DATA80). <i>Journal of Electrocardiology</i> , 2013 , 46, 360-5	1.4	10
156	Mother-offspring aggregation in home versus conventional blood pressure in the Tohoku Study of Child Development (TSCD). <i>Acta Cardiologica</i> , 2012 , 67, 449-56	0.9	10
155	Influence of home blood pressure measuring conditions in the evening on the morning-evening home blood pressure difference in treated hypertensive patients: the J-HOME study. <i>Blood Pressure Monitoring</i> , 2009 , 14, 160-5	1.3	10
154	The velocity of antihypertensive effects of seven angiotensin II receptor blockers determined by home blood pressure measurements. <i>Journal of Hypertension</i> , 2016 , 34, 1218-23	1.9	10
153	Dietary Inflammatory Index Positively Associated With High-Sensitivity C-Reactive Protein Level in Japanese From NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2020 , 30, 98-107	3.4	10
152	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S59-S65	3.4	10
151	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 606-620	4	10
150	Socioeconomic and lifestyle factors associated with depressive tendencies in general Japanese men and women: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019 , 24, 37	4.2	9
149	Long-term outcomes associated with prolonged PR interval in the general Japanese population. <i>International Journal of Cardiology</i> , 2015 , 184, 291-293	3.2	9
148	Menstrual Factors and Stroke Incidence in Japanese Postmenopausal Women: The Ohasama Study. <i>Neuroepidemiology</i> , 2016 , 47, 109-116	5.4	9
147	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 1195-1202.e1	4.9	9
146	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: a 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014 , 64, 218-24	3	9
145	Self-monitoring of ambulatory blood pressure by the Microlife WatchBP O3--an application test. <i>Clinical and Experimental Hypertension</i> , 2011 , 33, 34-40	2.2	9
144	Blood pressure changes during pregnancy. <i>Hypertension Research</i> , 2012 , 35, 563-4; author reply 34	4.7	9
143	Efficacy of Combination Antihypertensive Therapy with Low-Dose Indapamide: Assessment by Blood Pressure Self-Monitoring at Home. <i>Clinical and Experimental Hypertension</i> , 2005 , 27, 331-341	2.2	9
142	Renal sinus fat volume on computed tomography in middle-aged patients at risk for cardiovascular disease and its association with coronary artery calcification. <i>Atherosclerosis</i> , 2016 , 246, 374-81	3.1	9
141	Prevalence and Determinants of Masked Hypertension Among Black Nigerians Compared With a Reference Population. <i>Hypertension</i> , 2016 , 67, 1249-55	8.5	9

140	Absolute risk score for stroke, myocardial infarction, and all cardiovascular disease: Japan Arteriosclerosis Longitudinal Study. <i>Hypertension Research</i> , 2019 , 42, 567-579	4.7	9
139	Nocturnal blood pressure decline based on different time intervals and long-term cardiovascular risk: the Ohasama Study. <i>Clinical and Experimental Hypertension</i> , 2018 , 40, 1-7	2.2	9
138	Relationships among Socioeconomic Factors and Self-rated Health in Japanese Adults: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S66-S72	3.4	9
137	Socioeconomic Status and Knowledge of Cardiovascular Risk Factors: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S46-S52	3.4	9
136	Is antihypertensive treatment based on home blood pressure recommended rather than that based on office blood pressure in adults with essential hypertension? (meta-analysis). <i>Hypertension Research</i> , 2019 , 42, 807-816	4.7	8
135	Association of aldosterone-to-renin ratio with hypertension differs by sodium intake: the Ohasama study. <i>American Journal of Hypertension</i> , 2015 , 28, 208-15	2.3	8
134	Home blood pressure level and decline in renal function among treated hypertensive patients: the J-HOME-Morning Study. <i>Hypertension Research</i> , 2016 , 39, 107-12	4.7	8
133	Electrocardiographic abnormalities and home blood pressure in treated elderly hypertensive patients: Japan home versus office blood pressure measurement evaluation in the elderly (J-HOME-Elderly) study. <i>Hypertension Research</i> , 2010 , 33, 670-7	4.7	8
132	Pharmacists' awareness and attitude toward blood pressure measurement at home and in the pharmacy in Japan. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 447-55	2.2	8
131	Study design of HOMED-BP: hypertension objective treatment based on measurement by electrical devices of blood pressure. <i>Clinical and Experimental Hypertension</i> , 2003 , 25, 143-4	2.2	8
130	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , 2016 , 39, 670-9	4.7	8
129	Association of maternal home blood pressure trajectory during pregnancy with infant birth weight: the BOSHI study. <i>Hypertension Research</i> , 2020 , 43, 550-559	4.7	7
128	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018 , 37, 182-188	5.9	7
127	Fasting Blood Glucose Predicts Incidence of Hypertension Independent of HbA1c Levels and Insulin Resistance in Middle-Aged Japanese: The Saku Study. <i>American Journal of Hypertension</i> , 2019 , 32, 1178-1185	2.3	7
126	Response to Masked hypertension in untreated and treated patients with diabetes mellitus: attractive but questionable interpretations and response to Is masked hypertension related to diabetes mellitus?. <i>Hypertension</i> , 2013 , 62, e23-5	8.5	7
125	Comparability in coronary artery calcium scores on CT scan between two community-based cohort studies. <i>International Journal of Cardiology</i> , 2011 , 149, 244-245	3.2	7
124	Prognostic significance of home arterial stiffness index derived from self-measurement of blood pressure: the Ohasama Study. <i>American Journal of Hypertension</i> , 2012 , 25, 67-73	2.3	7
123	Role of angiotensinogen and relative aldosterone excess in salt-sensitive hypertension. <i>Hypertension</i> , 2012 , 59, e57; author reply e58	8.5	7

122	Isolated systolic hypertension and 29-year cardiovascular mortality risk in Japanese adults aged 30–49 years. <i>Journal of Hypertension</i> , 2020 , 38, 2230-2236	1.9	7
121	Validation of an automated home blood pressure measurement device in oldest-old populations. <i>Hypertension Research</i> , 2020 , 43, 30-35	4.7	7
120	Association of Fatal and Nonfatal Cardiovascular Outcomes With 24-Hour Mean Arterial Pressure. <i>Hypertension</i> , 2021 , 77, 39-48	8.5	7
119	Added predictive value of high uric acid for cardiovascular events in the Ambulatory Blood Pressure International Study. <i>Journal of Clinical Hypertension</i> , 2019 , 21, 966-974	2.3	6
118	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 417-423	3.2	6
117	Identification of disaster-vulnerable communities by use of census data prior to the Great East Japan Earthquake. <i>Disaster Medicine and Public Health Preparedness</i> , 2015 , 9, 19-28	2.8	6
116	N-Terminal Pro-B-Type Natriuretic Peptide Is Not a Significant Predictor of Stroke Incidence After 5 Years - The Ohasama Study. <i>Circulation Journal</i> , 2018 , 82, 2055-2062	2.9	6
115	Mapping the genetic diversity of HLA haplotypes in the Japanese populations. <i>Scientific Reports</i> , 2015 , 5, 17855	4.9	6
114	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013 , 99, 1024-9	5.1	6
113	Detection of midpregnancy fall in blood pressure by out-of-office monitoring. <i>Hypertension</i> , 2009 , 53, e12-3; author reply e14	8.5	6
112	How many measurements are needed to provide reliable information in terms of the ambulatory arterial stiffness index? The Ohasama study. <i>Hypertension Research</i> , 2011 , 34, 314-8	4.7	6
111	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , 2016 , 252, 116-121	3.1	6
110	Steno-Stiffness Approach for Cardiovascular Disease Risk Assessment in Primary Prevention. <i>Hypertension</i> , 2019 , 73, 508-513	8.5	6
109	Socioeconomic Inequalities in Oral Health Among Unmarried and Married Women: Evidence From a Population-Based Study in Japan. <i>Journal of Epidemiology</i> , 2018 , 28, 341-346	3.4	6
108	Antihypertensives prescribed for pregnant women in Japan: Prevalence and timing determined from a database of health insurance claims. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 1325-1334	2.6	6
107	Identification of physical and psychosocial problems associated with diabetic nephropathy using the International Classification of Functioning, Disability and Health Core Set for Diabetes Mellitus. <i>Clinical and Experimental Nephrology</i> , 2016 , 20, 187-94	2.5	5
106	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , 2017 , 24, 505-513	3.9	5
105	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 995-1009	4	5

104	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 799-805	2.9	5
103	Cardiovascular risk and blood pressure lowering treatment among elderly individuals: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan. <i>Journal of Hypertension</i> , 2018 , 36, 410-418	1.9	5
102	Evaluating home blood pressure in treated hypertensives in comparison with the referential value of casual screening of blood pressure: the Ohasama study. <i>Blood Pressure Monitoring</i> , 2012 , 17, 89-95	1.3	5
101	Parental longevity and offspring home blood pressure: the Ohasama study. <i>Journal of Hypertension</i> , 2010 , 28, 272-7	1.9	5
100	Blood Pressure and Chronic Kidney Disease Stratified by Gender and the Use of Antihypertensive Drugs. <i>Journal of the American Heart Association</i> , 2020 , 9, e015592	6	5
99	Prognostic Value of Ambulatory Blood Pressure in the Obese: The Ambulatory Blood Pressure-International Study. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 111-8	2.3	5
98	Stroke risk due to partial white-coat or masked hypertension based on the ACC/AHA guideline blood pressure threshold: the Ohasama study. <i>Hypertension Research</i> , 2019 , 42, 120-122	4.7	5
97	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S23-S28	3.4	5
96	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men - A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , 2019 , 83, 1506-1513	2.9	4
95	Randomized trial comparing the velocities of the antihypertensive effects on home blood pressure of candesartan and candesartan with hydrochlorothiazide. <i>Hypertension Research</i> , 2015 , 38, 701-7	4.7	4
94	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018 , 82, 2542-2548	2.9	4
93	Lacunar Infarcts Rather than White Matter Hyperintensity as a Predictor of Future Higher Level Functional Decline: The Ohasama Study. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017 , 26, 376-384	2.8	4
92	Associations between visit-to-visit variability in blood pressure measured in the office and antihypertensive drugs: the J-HOME-Morning study. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 285-90	2.2	4
91	Association between white matter hyperintensity and lacunar infarction on MRI and subitem scores of the Japanese version of mini-mental state examination for testing cognitive decline: the Ohasama study. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 541-7	2.2	4
90	Association between low birth weight and high adult waist-to-height ratio in non-obese women: a cross-sectional study in a Japanese population. <i>Tohoku Journal of Experimental Medicine</i> , 2012 , 228, 205-214	2.4	4
89	Time-dependent effects of imidapril administration in patients with morning hypertension measured as home blood pressure. <i>Clinical and Experimental Hypertension</i> , 2008 , 30, 243-54	2.2	4
88	Effectiveness of a Self-Monitoring Quality of Life Intervention For Patients with Cancer Receiving Palliative Care: A Randomized Controlled Clinical Trial. <i>Asian Pacific Journal of Cancer Prevention</i> , 2019 , 20, 2795-2802	1.7	4
87	Effects of a Self-Monitoring Quality of Life Intervention for Patients with Cancer Receiving Palliative Care in Japan: Study Protocol for a Randomized Controlled Trial. <i>Asian Pacific Journal of Cancer Prevention</i> , 2018 , 19, 3027-3032	1.7	4

86	Identification of physical and psychosocial problems based on symptoms in patients with Behçet disease. <i>Rheumatology International</i> , 2020 , 40, 81-89	3.6	4
85	Relative and Absolute Risk to Guide the Management of Pulse Pressure, an Age-Related Cardiovascular Risk Factor. <i>American Journal of Hypertension</i> , 2021 , 34, 929-938	2.3	4
84	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019 , 37, 1676-1681	1.9	4
83	The association between glycemic control and lung function impairment in individuals with diabetes: the Saku study. <i>Diabetology International</i> , 2019 , 10, 213-218	2.3	4
82	The present situation of home blood pressure measurement among outpatients in Japan. <i>Clinical and Experimental Hypertension</i> , 2020 , 42, 67-74	2.2	4
81	A Combination of Blood Pressure and Total Cholesterol Increases the Lifetime Risk of Coronary Heart Disease Mortality: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 6-24	4	4
80	Predictive power of home blood pressure indices at baseline and during follow-up in hypertensive patients: HOMED-BP study. <i>Hypertension Research</i> , 2018 , 41, 622-628	4.7	4
79	Factors Related to Participation in Health Examinations for Japanese National Health Insurance: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S53-S58	3.4	4
78	Left ventricular hypertrophy by electrocardiogram as a predictor of success in home blood pressure control: HOMED-BP study. <i>Hypertension Research</i> , 2017 , 40, 504-510	4.7	3
77	Development and validation of a short-version checklist for patients undergoing hemodialysis based on the International Classification of Functioning, Disability and Health. <i>Clinical and Experimental Nephrology</i> , 2015 , 19, 953-60	2.5	3
76	Differential effects of angiotensin II receptor blocker and losartan/hydrochlorothiazide combination on central blood pressure and augmentation index. <i>Clinical and Experimental Hypertension</i> , 2015 , 37, 294-302	2.2	3
75	Comparison of nocturnal blood pressure based on home versus ambulatory blood pressure measurement: The Ohasama Study. <i>Clinical and Experimental Hypertension</i> , 2020 , 42, 685-691	2.2	3
74	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019 , 29, 133-138	3.4	3
73	Age-Related Trends in Home Blood Pressure, Home Pulse Rate, and Day-to-Day Blood Pressure and Pulse Rate Variability Based on Longitudinal Cohort Data: The Ohasama Study. <i>Journal of the American Heart Association</i> , 2019 , 8, e012121	6	3
72	Home blood pressure predicts stroke incidence among older adults with impaired physical function: the Ohasama study. <i>Journal of Hypertension</i> , 2017 , 35, 2395-2401	1.9	3
71	Differences between clinic and home blood pressure measurements during pregnancy. <i>Journal of Hypertension</i> , 2015 , 33, 1492-3	1.9	3
70	Compliance to the recommendations on blood pressure measurements in the clinical settings. <i>Journal of Hypertension</i> , 2013 , 31, 213	1.9	3
69	Blood pressure-lowering effect and duration of action of bedtime administration of doxazosin determined by home blood pressure measurement. <i>Clinical and Experimental Hypertension</i> , 2010 , 32, 311-7	2.2	3

68	Associated factors of home versus ambulatory heart rate variability in the general population: the Ohasama study. <i>Clinical and Experimental Hypertension</i> , 2011 , 33, 404-10	2.2	3
67	Role of relative aldosterone excess in salt-sensitive hypertension among African ancestry. <i>American Journal of Hypertension</i> , 2012 , 25, 398-9; author reply 400	2.3	3
66	Control of home heart rate and home blood pressure levels in treated patients with hypertension: the J-HOME study. <i>Blood Pressure Monitoring</i> , 2007 , 12, 289-95	1.3	3
65	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , 2016 , 11, e0157563	3.7	3
64	Blood pressure variability and arterial stiffness parameters derived from ambulatory blood pressure monitoring. <i>Kardiologia Polska</i> , 2019 , 77, 509-514	0.9	3
63	Isolated Diastolic Hypertension in the IDACO Study: An Age-Stratified Analysis Using 24-Hour Ambulatory Blood Pressure Measurements. <i>Hypertension</i> , 2021 , 78, 1222-1231	8.5	3
62	Recent status of self-measured home blood pressure in the Japanese general population: a modern database on self-measured home blood pressure (MDAS). <i>Hypertension Research</i> , 2020 , 43, 1403-1412	4.7	3
61	In-office and out-of-office blood pressure measurement. <i>Journal of Human Hypertension</i> , 2021 ,	2.6	3
60	Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2021 , 26, 57	4.2	3
59	Knowledge, Attitude, and Practices Toward Blood Pressure Measurement at Home Among Japanese Nurses. <i>Home Healthcare Now</i> , 2016 , 34, 210-7	0.5	3
58	Factors associated with intra-individual visit-to-visit variability of blood pressure in four countries: the INTERMAP study. <i>Journal of Human Hypertension</i> , 2019 , 33, 229-236	2.6	3
57	Risk of developing type 2 diabetes according to blood pressure levels and presence or absence of hypertensive treatment: the Saku study. <i>Hypertension Research</i> , 2019 , 42, 105-113	4.7	3
56	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020 , 30, 244-252	3.4	3
55	Insights on home blood pressure monitoring in Asia: Expert perspectives from 10 countries/regions. <i>Journal of Clinical Hypertension</i> , 2021 , 23, 3-11	2.3	3
54	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S40-S45	3.4	3
53	Response to Yatabe et al. <i>Hypertension Research</i> , 2017 , 40, 789-790	4.7	2
52	Personality traits as predictors of decline in higher-level functional capacity over a 7-year follow-up in older adults: the Ohasama study. <i>Tohoku Journal of Experimental Medicine</i> , 2014 , 234, 197-207	2.4	2
51	Contribution of the ABP-International study to the definition of night-time tachycardia. <i>Journal of Hypertension</i> , 2014 , 32, 2099-100	1.9	2

50	The velocity of home blood pressure reduction in response to low-dose eplerenone combined with other antihypertensive drugs determined by exponential decay function analysis. <i>Clinical and Experimental Hypertension</i> , 2014 , 36, 83-91	2.2	2
49	Change of the management of treated hypertensive patients with or without diabetes in Japan. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 79-86	2.2	2
48	Low birth weight is associated with high waist-to-height ratio in Japanese elementary school girls. <i>Tohoku Journal of Experimental Medicine</i> , 2013 , 231, 85-91	2.4	2
47	N-Terminal Pro-B-Type Natriuretic Peptide Is a Predictor of Chronic Kidney Disease in an Asian General Population - The Ohasama Study. <i>Circulation Reports</i> , 2019 , 2, 24-32	0.7	2
46	Diagnostic Value of Home Blood Pressure. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2020 , 45-54	0.1	2
45	Fatty Acid Intakes and Coronary Heart Disease Mortality in Japan: NIPPON DATA90, 1990-2005. <i>Current Nutrition and Food Science</i> , 2013 , 9, 26-32	0.7	2
44	Lifetime risk of stroke stratified by chronic kidney disease and hypertension in the general Asian population: the Ohasama study. <i>Hypertension Research</i> , 2021 , 44, 866-873	4.7	2
43	Data set for renal sinus fat volume and visceral adipose tissue volume on computed tomography. <i>Data in Brief</i> , 2016 , 7, 1658-64	1.2	2
42	Effects of measurement intervals on the values of repeated auscultatory blood pressure measurements. <i>Clinical and Experimental Hypertension</i> , 2020 , 42, 105-109	2.2	2
41	The Associations of Indices of Obesity with Lipoprotein Subfractions in Japanese American, African American and Korean Men. <i>Global Heart</i> , 2013 , 8, 273-280	2.9	2
40	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , 2021 , 44, 80-87	4.7	2
39	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , 2021 , 90, 57-65	2.7	2
38	Effect of amlodipine, efonidipine, and trichlormethiazide on home blood pressure and upper-normal microalbuminuria assessed by casual spot urine test in essential hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2018 , 40, 468-475	2.2	2
37	Genome-wide association study for white coat effect in Japanese middle-aged to elderly people: The HOMED-BP study. <i>Clinical and Experimental Hypertension</i> , 2018 , 40, 363-369	2.2	2
36	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S35-S39	3.4	2
35	Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. <i>PLoS ONE</i> , 2021 , 16, e0254706	3.7	2
34	Detailed association between serum uric acid levels and the incidence of chronic kidney disease stratified by sex in middle-aged adults. <i>Atherosclerosis</i> , 2021 , 330, 107-113	3.1	2
33	Antioxidant Capacities of Plant-Derived Foods Commonly Consumed in Japan. <i>Journal of Nutritional Science and Vitaminology</i> , 2020 , 66, 68-74	1.1	1

32	Do estimated 24-h pulse pressure components affect outcome? The Ohasama study. <i>Journal of Hypertension</i> , 2020 , 38, 1286-1292	1.9	1
31	Gender differences in the perception of difficulty of self-management in patients with diabetes mellitus: a mixed-methods approach. <i>Diabetology International</i> , 2016 , 7, 289-298	2.3	1
30	Changes in body weight and height in survivors of Menkes disease. <i>Journal of Trace Elements in Medicine and Biology</i> , 2014 , 28, 470-3	4.1	1
29	Is there a role for day-to-day home blood pressure variability in guiding management of hypertension?. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2014 , 41, 54-7	3	1
28	Children@ home blood pressure and growth environment. <i>Hypertension</i> , 2013 , 61, e33	8.5	1
27	Is more aggressive prevention of coronary artery disease required for patients with early repolarization syndrome?--Reply. <i>Circulation Journal</i> , 2013 , 77, 1643	2.9	1
26	Antihypertensive drug effects according to the pretreatment self-measured home blood pressure: the HOMED-BP study. <i>BMJ Open</i> , 2020 , 10, e040524	3	1
25	Moderate morning rise in blood pressure has lowest risk of stroke but only in women. <i>Journal of Hypertension</i> , 2019 , 37, 1437-1447	1.9	1
24	Altered immunogenicity of 23-valent pneumococcal polysaccharide vaccine in elderly patients with diabetes who revealed lower responses to concomitant administration of BIKEN varicella zoster vaccine: Results of post hoc analysis of a randomized double-blind trial. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 243-248	3.2	1
23	Electrocardiographic Left Atrial Abnormality and B-Type Natriuretic Peptide in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 34-43	4	1
22	Examining the trimester-specific effects of low gestational weight gain on birthweight: the BOSHI study. <i>Journal of Developmental Origins of Health and Disease</i> , 2021 , 12, 280-285	2.4	1
21	The Major Source of Antioxidants Intake From Typical Diet Among Rural Farmers in North-eastern Japan in the 1990s. <i>Journal of Epidemiology</i> , 2021 , 31, 101-108	3.4	1
20	Relationship Between Calcium Intake and Impaired Activities of Daily Living in a Japanese Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2021 , 31, 119-124	3.4	1
19	Impact of diastolic blood pressure threshold for the young population: the Japan Arteriosclerosis Longitudinal Study (JALS). <i>Journal of Hypertension</i> , 2019 , 37, 652-653	1.9	0
18	Relationship between stroke subtypes and morning surge or dipping pattern. <i>Hypertension</i> , 2013 , 61, e21	8.5	0
17	Prediction of Lifetime Risk of Cardiovascular Disease Deaths Stratified by Sex in the Japanese Population. <i>Journal of the American Heart Association</i> , 2021 , 10, e021753	6	0
16	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 252, 253-262	2.4	0
15	Association of self-measured home, ambulatory, and strictly measured office blood pressure and their variability with intracranial arterial stenosis. <i>Journal of Hypertension</i> , 2021 , 39, 2030-2039	1.9	0

14	Validity and Reliability of the Japanese Version of the 10-Item Perceived Efficacy in Patient-Physician Interactions (PEPPI-10) Scale in Breast Cancer Outpatients. <i>Tohoku Journal of Experimental Medicine</i> , 2019 , 249, 121-126	2.4	○
13	Cluster-randomized controlled trial for the early promotion of clinic visits for untreated hypertension. <i>Hypertension Research</i> , 2021 , 44, 355-362	4.7	○
12	The association of disproportionately enlarged subarachnoid space hydrocephalus with cognitive deficit in a general population: the Ohasama study. <i>Scientific Reports</i> , 2021 , 11, 17061	4.9	○
11	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 746	4	
10	Hypotensive and heart rate-lowering effects of low-dose bisoprolol determined based on self-measured blood pressure at home. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 284-9	2.2	
9	Is high prorenin levels related to relative aldosterone excess?. <i>American Journal of Hypertension</i> , 2013 , 26, 153	2.3	
8	Blood Pressure Phenotypes Defined by Ambulatory Blood Pressure Monitoring and Carotid Artery Changes in Community-Dwelling Older Japanese Adults: The Ohasama Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 252, 269-279	2.4	
7	Blood Pressure and Cardiovascular Disease in the Elderly. <i>Current Topics in Environmental Health and Preventive Medicine</i> , 2019 , 107-119	0.3	
6	Nocturnal Home Blood Pressure Monitoring. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2020 , 121-129	0.1	
5	Home Blood Pressure Variability. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2020 , 143-154	0.1	
4	Home Blood Pressure as Predictor of Adverse Health Outcomes. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2020 , 33-43	0.1	
3	Home Blood Pressure Monitoring Schedule. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2020 , 55-62	0.1	
2	Factors Associated with Lower Cognitive Performance Scores Among Older Japanese Men in Hawaii and Japan. <i>Journal of Alzheimer's Disease</i> , 2021 , 81, 403-412	4.3	
1	Validity and reliability of a checklist for patients with Behçet disease based on the International Classification of Functioning, Disability and Health. <i>Rheumatology International</i> , 2021 , 1	3.6	