
Riccardo Cristiani

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3969209/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Only one patient out of five achieves symmetrical knee function 6 months after primary anterior cruciate ligament reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2019, 27, 3461-3470.	4.2	59
2	Age, gender, quadriceps strength and hop test performance are the most important factors affecting the achievement of a patient-acceptable symptom state after ACL reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2020, 28, 369-380.	4.2	48
3	Increased knee laxity with hamstring tendon autograft compared to patellar tendon autograft: a cohort study of 5462 patients with primary anterior cruciate ligament reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2019, 27, 381-388.	4.2	46
4	Medial Meniscus Resection Increases and Medial Meniscus Repair Preserves Anterior Knee Laxity: A Cohort Study of 4497 Patients With Primary Anterior Cruciate Ligament Reconstruction. American Journal of Sports Medicine, 2018, 46, 357-362.	4.2	40
5	Revision anterior cruciate ligament reconstruction restores knee laxity but shows inferior functional knee outcome compared with primary reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2019, 27, 137-145.	4.2	36
6	Meniscal repair results in inferior short-term outcomes compared with meniscal resection: a cohort study of 6398 patients with primary anterior cruciate ligament reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2018, 26, 2251-2258.	4.2	33
7	Risk Factors for Abnormal Anteroposterior Knee Laxity After Primary Anterior Cruciate Ligament Reconstruction. Arthroscopy - Journal of Arthroscopic and Related Surgery, 2018, 34, 2478-2484.	2.7	26
8	Delayed Anterior Cruciate Ligament Reconstruction Increases the Risk of Abnormal Prereconstruction Laxity, Cartilage, and Medial Meniscus Injuries. Arthroscopy - Journal of Arthroscopic and Related Surgery, 2021, 37, 1214-1220.	2.7	25
9	Age, time from injury to surgery and quadriceps strength affect the risk of revision surgery after primary ACL reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2021, 29, 4154-4162.	4.2	24
10	Autograft type affects muscle strength and hop performance after ACL reconstruction. A randomised controlled trial comparing patellar tendon and hamstring tendon autografts with standard or accelerated rehabilitation. Knee Surgery, Sports Traumatology, Arthroscopy, 2021, 29, 3025-3036.	4.2	22
11	Meniscus Repair Does Not Result in an Inferior Short-term Outcome Compared With Meniscus Resection: An Analysis of 5,378 Patients With Primary Anterior Cruciate Ligament Reconstruction. Arthroscopy - Journal of Arthroscopic and Related Surgery, 2020, 36, 1145-1153.	2.7	19
12	One sixth of primary anterior cruciate ligament reconstructions may undergo reoperation due to complications or new injuries within 2Âyears. Knee Surgery, Sports Traumatology, Arthroscopy, 2020, 28, 2478-2485.	4.2	13
13	Suture tape reinforcement of hamstring tendon graft reduces postoperative knee laxity after primary ACL reconstruction. Journal of Experimental Orthopaedics, 2022, 9, 20.	1.8	10
14	Contralateral knee hyperextension is associated with increased anterior tibial translation and fewer meniscal injuries in the anterior cruciate ligament-injured knee. Knee Surgery, Sports Traumatology, Arthroscopy, 2018, 26, 3020-3028.	4.2	5
15	Age, time from injury to surgery and hop performance after primary ACLR affect the risk of contralateral ACLR. Knee Surgery, Sports Traumatology, Arthroscopy, 2022, 30, 1828-1835.	4.2	5
16	Subsequent surgery after primary ACLR results in a significantly inferior subjective outcome at a 2-year follow-up. Knee Surgery, Sports Traumatology, Arthroscopy, 2022, 30, 1927-1936.	4.2	5
17	Knee laxity and functional knee outcome after contralateral ACLR are comparable to those after primary ACLR. Knee Surgery, Sports Traumatology, Arthroscopy, 2021, 29, 3864-3870.	4.2	4
18	Comparison of Knee Function and Activity Level Between Bilateral and Unilateral ACL Reconstruction: A Matched-Group Analysis With Minimum 5-Year Follow-up. Orthopaedic Journal of Sports Medicine, 2022, 10, 232596712210835.	1.7	3

#	Article	IF	CITATIONS
19	Regarding "Editorial Commentary: Meniscal Repair—Why Bother?― Arthroscopy - Journal of Arthroscopic and Related Surgery, 2020, 36, 1794-1795.	2.7	0