

Riccardo Cristiani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3969209/publications.pdf>

Version: 2024-02-01

19
papers

423
citations

840776

11
h-index

839539

18
g-index

19
all docs

19
docs citations

19
times ranked

319
citing authors

#	ARTICLE	IF	CITATIONS
1	Age, time from injury to surgery and hop performance after primary ACLR affect the risk of contralateral ACLR. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022, 30, 1828-1835.	4.2	5
2	Suture tape reinforcement of hamstring tendon graft reduces postoperative knee laxity after primary ACL reconstruction. <i>Journal of Experimental Orthopaedics</i> , 2022, 9, 20.	1.8	10
3	Subsequent surgery after primary ACLR results in a significantly inferior subjective outcome at a 2-year follow-up. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022, 30, 1927-1936.	4.2	5
4	Comparison of Knee Function and Activity Level Between Bilateral and Unilateral ACL Reconstruction: A Matched-Group Analysis With Minimum 5-Year Follow-up. <i>Orthopaedic Journal of Sports Medicine</i> , 2022, 10, 232596712210835.	1.7	3
5	Autograft type affects muscle strength and hop performance after ACL reconstruction. A randomised controlled trial comparing patellar tendon and hamstring tendon autografts with standard or accelerated rehabilitation. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 3025-3036.	4.2	22
6	Delayed Anterior Cruciate Ligament Reconstruction Increases the Risk of Abnormal Prereconstruction Laxity, Cartilage, and Medial Meniscus Injuries. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2021, 37, 1214-1220.	2.7	25
7	Age, time from injury to surgery and quadriceps strength affect the risk of revision surgery after primary ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 4154-4162.	4.2	24
8	Knee laxity and functional knee outcome after contralateral ACLR are comparable to those after primary ACLR. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 3864-3870.	4.2	4
9	Age, gender, quadriceps strength and hop test performance are the most important factors affecting the achievement of a patient-acceptable symptom state after ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 369-380.	4.2	48
10	Meniscus Repair Does Not Result in an Inferior Short-term Outcome Compared With Meniscus Resection: An Analysis of 5,378 Patients With Primary Anterior Cruciate Ligament Reconstruction. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2020, 36, 1145-1153.	2.7	19
11	One sixth of primary anterior cruciate ligament reconstructions may undergo reoperation due to complications or new injuries within 2 years. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 2478-2485.	4.2	13
12	Regarding "Editorial Commentary: Meniscal Repair" Why Bother? <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2020, 36, 1794-1795.	2.7	0
13	Increased knee laxity with hamstring tendon autograft compared to patellar tendon autograft: a cohort study of 5462 patients with primary anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2019, 27, 381-388.	4.2	46
14	Revision anterior cruciate ligament reconstruction restores knee laxity but shows inferior functional knee outcome compared with primary reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2019, 27, 137-145.	4.2	36
15	Only one patient out of five achieves symmetrical knee function 6 months after primary anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2019, 27, 3461-3470.	4.2	59
16	Medial Meniscus Resection Increases and Medial Meniscus Repair Preserves Anterior Knee Laxity: A Cohort Study of 4497 Patients With Primary Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> , 2018, 46, 357-362.	4.2	40
17	Meniscal repair results in inferior short-term outcomes compared with meniscal resection: a cohort study of 6398 patients with primary anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018, 26, 2251-2258.	4.2	33
18	Contralateral knee hyperextension is associated with increased anterior tibial translation and fewer meniscal injuries in the anterior cruciate ligament-injured knee. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018, 26, 3020-3028.	4.2	5

#	ARTICLE	IF	CITATIONS
19	Risk Factors for Abnormal Anteroposterior Knee Laxity After Primary Anterior Cruciate Ligament Reconstruction. Arthroscopy - Journal of Arthroscopic and Related Surgery, 2018, 34, 2478-2484.	2.7	26