

Laurentius Huber

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3971479/publications.pdf>

Version: 2024-02-01

33
papers

1,730
citations

394286

19
h-index

395590

33
g-index

40
all docs

40
docs citations

40
times ranked

1344
citing authors

#	ARTICLE	IF	CITATIONS
1	Concurrent CBF and BOLD fMRI with dual-echo spiral simultaneous multi-slice acquisitions at 7T. <i>NeuroImage</i> , 2022, 247, 118820.	2.1	5
2	Layer-specific activation in human primary somatosensory cortex during tactile temporal prediction error processing. <i>NeuroImage</i> , 2022, 248, 118867.	2.1	11
3	Higher and deeper: Bringing layer fMRI to association cortex. <i>Progress in Neurobiology</i> , 2021, 207, 101930.	2.8	21
4	Layer-dependent functional connectivity methods. <i>Progress in Neurobiology</i> , 2021, 207, 101835.	2.8	67
5	Sub-millimetre resolution laminar fMRI using Arterial Spin Labelling in humans at 7 T. <i>PLoS ONE</i> , 2021, 16, e0250504.	1.1	27
6	LayNii: A software suite for layer-fMRI. <i>NeuroImage</i> , 2021, 237, 118091.	2.1	64
7	Challenges and opportunities of mesoscopic brain mapping with fMRI. <i>Current Opinion in Behavioral Sciences</i> , 2021, 40, 189-200.	2.0	15
8	Linking cortical circuit models to human cognition with laminar fMRI. <i>Neuroscience and Biobehavioral Reviews</i> , 2021, 128, 467-478.	2.9	17
9	Magnetization transfer weighted EPI facilitates cortical depth determination in native fMRI space. <i>NeuroImage</i> , 2021, 242, 118455.	2.1	4
10	Sub-millimeter fMRI reveals multiple topographical digit representations that form action maps in human motor cortex. <i>NeuroImage</i> , 2020, 208, 116463.	2.1	88
11	Layer-Specific Contributions to Imagined and Executed Hand Movements in Human Primary Motor Cortex. <i>Current Biology</i> , 2020, 30, 1721-1725.e3.	1.8	35
12	Comparison of BOLD and CBV using 3D EPI and 3D GRASE for cortical layer functional MRI at 7 T. <i>Magnetic Resonance in Medicine</i> , 2020, 84, 3128-3145.	1.9	33
13	Cortical laminar resting-state signal fluctuations scale with the hypercapnic blood oxygenation level-dependent response. <i>Human Brain Mapping</i> , 2020, 41, 2014-2027.	1.9	25
14	Layer-dependent activity in human prefrontal cortex during working memory. <i>Nature Neuroscience</i> , 2019, 22, 1687-1695.	7.1	130
15	Layer-specific activation of sensory input and predictive feedback in the human primary somatosensory cortex. <i>Science Advances</i> , 2019, 5, eaav9053.	4.7	62
16	Non-BOLD contrast for laminar fMRI in humans: CBF, CBV, and CMRO ₂ . <i>NeuroImage</i> , 2019, 197, 742-760.	2.1	96
17	Techniques for blood volume fMRI with VASO: From low-resolution mapping towards sub-millimeter layer-dependent applications. <i>NeuroImage</i> , 2018, 164, 131-143.	2.1	101
18	Ultra-high resolution blood volume fMRI and BOLD fMRI in humans at 9.4 T: Capabilities and challenges. <i>NeuroImage</i> , 2018, 178, 769-779.	2.1	44

#	ARTICLE	IF	CITATIONS
19	Physiological basis of vascular autocalibration (VasA): Comparison to hypercapnia calibration methods. <i>Magnetic Resonance in Medicine</i> , 2017, 78, 1168-1173.	1.9	7
20	High-Resolution CBV-fMRI Allows Mapping of Laminar Activity and Connectivity of Cortical Input and Output in Human M1. <i>Neuron</i> , 2017, 96, 1253-1263.e7.	3.8	255
21	Optimization of simultaneous multislice EPI for concurrent functional perfusion and BOLD signal measurements at 7T. <i>Magnetic Resonance in Medicine</i> , 2017, 78, 121-129.	1.9	24
22	Functional cerebral blood volume mapping with simultaneous multi-slice acquisition. <i>NeuroImage</i> , 2016, 125, 1159-1168.	2.1	22
23	Lamina-dependent calibrated BOLD response in human primary motor cortex. <i>NeuroImage</i> , 2016, 141, 250-261.	2.1	66
24	Baseline oxygenation in the brain: Correlation between respiratory-calibration and susceptibility methods. <i>NeuroImage</i> , 2016, 125, 920-931.	2.1	35
25	Vascular autoresizing of fMRI (VasA fMRI) improves sensitivity of population studies: A pilot study. <i>NeuroImage</i> , 2016, 124, 794-805.	2.1	33
26	Simultaneous acquisition of cerebral blood volume, blood flow, and blood oxygenation-weighted MRI signals at ultra-high magnetic field. <i>Magnetic Resonance in Medicine</i> , 2015, 74, 513-517.	1.9	9
27	Cortical lamina-dependent blood volume changes in human brain at 7 T. <i>NeuroImage</i> , 2015, 107, 23-33.	2.1	152
28	Slab-selective, BOLD-corrected VASO at 7 Tesla provides measures of cerebral blood volume reactivity with high signal-to-noise ratio. <i>Magnetic Resonance in Medicine</i> , 2014, 72, 137-148.	1.9	107
29	Using carbogen for calibrated fMRI at 7Tesla: Comparison of direct and modelled estimation of the M parameter. <i>NeuroImage</i> , 2014, 84, 605-614.	2.1	9
30	Anatomical brain imaging at 7T using two-dimensional GRASE. <i>Magnetic Resonance in Medicine</i> , 2014, 72, 1291-1301.	1.9	12
31	Regional reproducibility of calibrated BOLD functional MRI: Implications for the study of cognition and plasticity. <i>NeuroImage</i> , 2014, 101, 8-20.	2.1	18
32	Investigation of the neurovascular coupling in positive and negative BOLD responses in human brain at 7T. <i>NeuroImage</i> , 2014, 97, 349-362.	2.1	101
33	Fast accurate MR thermometry using phase referenced asymmetric spin-echo EPI at high field. <i>Magnetic Resonance in Medicine</i> , 2014, 71, 524-533.	1.9	12